

Pacific Journal of Mathematics

**ON RADICALS AND CONTINUITY OF HOMOMORPHISMS
INTO BANACH ALGEBRAS**

EDITH HIRSCH LUCHINS

ON RADICALS AND CONTINUITY OF HOMOMORPHISMS INTO BANACH ALGEBRAS

EDITH H. LUCHINS

1. Introduction. All Banach algebras considered are over the real field and all homomorphisms considered are algebraic (real-linear). An algebra is called semi-simple, strongly semi-simple, or strictly semi-simple, if its Jacobson radical [5], Segal radical [10], or strict radical [8], respectively, is the zero ideal; that is, if its regular maximal right ideals, its regular maximal two-sided ideals, or those of its two-sided ideals which are regular maximal right ideals, intersect in the zero ideal. Rickart [9, Corollary 6.3] proved that a semi-simple commutative Banach algebra has the property that every homomorphism of a Banach algebra into it is continuous. Call an algebra with this property an *absolute* algebra. Yood [12, Theorem 3.5] proved that every homomorphism of a Banach algebra onto a dense subset of a strongly semi-simple Banach algebra is continuous. Thus a strongly semi-simple Banach algebra is "almost" absolute. The question arose: Is a (noncommutative) semi-simple or strongly semi-simple Banach algebra necessarily absolute? A negative answer is furnished in the present note. It is shown that in order for a Banach algebra to be absolute it is sufficient that it be strictly semi-simple and necessary that it have zero as its only nilpotent element. The latter condition is shown to be sufficient for some special Banach algebras to be absolute.

2. Necessary condition for a Banach algebra to be absolute.

THEOREM 1. *An absolute Banach algebra has no nonzero nilpotent elements.*

Proof. Suppose the Banach algebra B contains a nonzero nilpotent element. Then there exists a nonzero $v \in B$ such that $v^2 = 0$. Let A be an infinite dimensional Banach algebra such that $A^2 = (0)$. Since A is an infinite dimensional complete vector space, there exists a discontinuous linear functional on A ; denote it by $f(x)$. Let $\pi(x) = f(x)v$. Since $f(x)$ is linear and $v^2 = 0$, π is seen to be a homomorphism of A into B .

Let $\|y\|$ be a Banach norm for B . Then $\|\pi(x)\| = |f(x)| \|v\|$ since $f(x)$ is a scalar. Since $f(x)$ is discontinuous $|f(x)|$ is not bounded and

Received February 5, 1959. This report was prepared while the author held the New York State Fellowship, 1957-8, of the American Association of University Women. It is based in part on the author's doctoral dissertation written at the University of Oregon under the sponsorship of Professor Bertram Yood. Thanks are offered to the referee for helpful suggestions.

hence π is not bounded. Thus π is discontinuous so that B is not absolute.

It is known that if a Banach algebra B is semi-simple and has a unique norm then a homomorphism of a Banach algebra onto B is necessarily continuous [9, Theorem 6.2]; that the resulting proposition is false if the word *into* is substituted for *onto* follows from our Theorem 1. Indeed, this theorem shows that for a Banach algebra to be absolute it is not sufficient that it have the properties of being simple, semi-simple, strongly semi-simple and having an identity and a unique norm. Thus, these properties are possessed by the algebra of all 2 by 2 matrices over the reals, under a Banach norm, and yet, since this algebra has nonzero nilpotent elements, Theorem 1 shows that it is not absolute.

3. Sufficient condition for a Banach algebra to be absolute. In [8] there was introduced the concept of a strictly semi-simple algebra. It was shown [8, Theorems 2 and 3] that a Banach algebra B is strictly semi-simple if and only if it is isomorphic to a subalgebra of $C(X, Q)$, the algebra of quaternion-valued functions continuous, and vanishing at ∞ , on a locally compact Hausdorff space X .

THEOREM 2. *A strictly semi-simple Banach algebra B is absolute.*

Proof. Let A be a Banach algebra with a homomorphism T into $B \subset C(X, Q)$. Let $T_x(a) = T(a)(x)$. The kernel of T_x is closed since Q is simple, and therefore T_x is continuous, whence T_x is of bound 1. That T is continuous can now be shown by the 6-line argument of Loomis [7, p. 77]. One could also use [12, Theorem 3.5].

COROLLARY 1 (Rickart). *A semi-simple commutative Banach algebra is absolute.*

4. Concerning some special Banach algebras. For each subset S of a Banach algebra B , let $S_L(S_R)$ denote the set of all left (right) annihilators of S . B is called an annihilator algebra [3] if $B_R = 0 = B_L$ and if $I_L \neq 0$ ($I_R \neq 0$) for each proper closed right (left) ideal I , where 0 denotes the zero ideal.

Lemma 1 is due to Forsythe and McCoy [4, p. 524].

LEMMA 1. *In a ring without nonzero nilpotent elements every idempotent is in the center.*

THEOREM 3. *That a Banach algebra B have zero as its only nilpotent element is both a necessary and a sufficient condition for B to be either strictly semi-simple or absolute, provided any of the following conditions is satisfied:*

- (a) B is finite-dimensional.
- (b) B satisfies the descending chain condition on right ideals.
- (c) B is a semi-simple annihilator algebra.

Proof. If B is strictly semi-simple, then $B \subset C(X, Q)$ by [8] and hence has only zero as a nilpotent element. If B is absolute, then zero is its only nilpotent element by Theorem 1. Conversely, suppose B has no nonzero nilpotent elements.

Suppose condition (a) or (b) holds. Then B has a nilpotent radical and therefore is semi-simple; also B is then a direct sum of division algebras and therefore has the property that every left (or right) ideal is two-sided [2, p. 463]. Thus B is strictly semi-simple and therefore absolute by Theorem 2.

Suppose condition (c) holds. Let M be any regular maximal right ideal in B . Bonsall and Goldie [3, pp. 155-6] show that for any semi-simple annihilator algebra B , $M_L = Be$ where e is a nonzero idempotent, B is a minimal (closed) left ideal, eB a minimal (closed) right ideal, $(eB)_L$ a maximal left ideal, and $(Be)_R = M$.

If B has no nonzero nilpotent elements, then e is in the center by Lemma 1 so that $Be = eB$ is a two-sided ideal. But the left and right annihilators of a closed two-sided ideal are identical [3, p. 159] so that $(eB)_L = (Be)_R = M$.

Since $(eB)_L$ is a left ideal, M , which was any regular maximal right ideal in B , has been shown to be a left ideal. Thus B is strictly semi-simple since it is semi-simple by hypothesis, and therefore absolute by Theorem 2.

COROLLARY 2. *An H^* algebra B is commutative if and only if any of the following properties is satisfied:*

- (a) B has no nonzero nilpotent elements.
- (b) B is strictly semi-simple.
- (c) B is absolute.

Proof. An H^* algebra is the closure of the direct sum of matrix algebras M_σ [1, pp. 379-380]. If condition (a) holds, then each M_σ must have zero as its only nilpotent element and therefore must be one-dimensional. Hence each M_σ is generated by an idempotent e_σ which, by Lemma 1, is in the center. For $u, v \in \Sigma M_\sigma$, $u = \sum r_k e_k$, $v = \sum s_t e_t$, r_k, s_t scalars, $uv = vu$ so that ΣM_σ is commutative and therefore so is its closure, B . Thus condition (a) implies that B is commutative.

Suppose B is commutative. Since an H^* algebra is semi-simple, if commutative it is strictly semi-simple and $\subset C(X, Q)$ by [8], so that zero is its only nilpotent element. Hence condition (a) prevails.

The remainder of the corollary follows immediately from Theorem 3 since an H^* algebra is a semi-simple annihilator algebra [6, p. 697].

BIBLIOGRAPHY

1. W. Ambrose, *Structure theory for a special class of Banach algebras*, Trans. Amer. Math. Soc. **57** (1945), 364-386.
2. R. Arens and I. Kaplansky, *Topological representation of algebras*, Trans. Amer. Math. Soc. **63** (1948), 457-481.
3. F. F. Bonsall and A. W. Goldie, *Annihilator algebras*, Proc. London Math. Soc. **5** (1954), 154-167.
4. A. Forsythe and N. H. McCoy, *On the commutativity of certain rings*, Bull. Amer. Math. Soc. **52** (1946), 523-526.
5. N. Jacobson, *The radical and semi-simplicity for arbitrary rings*, Amer. J. Math. **67** (1945), 300-320.
6. I. Kaplansky, *Dual rings*, Ann. of Math. **49** (1948), 689-701.
7. L. H. Loomis, *An introduction to abstract harmonic analysis*, New York: Van Nostrand, 1953.
8. E. H. Luchins, *On strictly semi-simple Banach algebras*, Pacific J. Math. (to appear.)
9. C. E. Rickart, *The uniqueness of norm problem in Banach algebras*, Ann. of Math. **51** (1950), 615-628.
10. I. E. Segal, *The group algebra of a locally compact group*, Trans. Amer. Math. Soc. **61** (1947), 69-105.
11. B. Yood, *Topological properties of homomorphisms between Banach algebras*, Amer. J. Math. **76** (1954), 155-167.
12. ———, *Homomorphisms on normed algebras*, Pacific J. Math. **8** (1958), 373-381.

UNIVERSITY OF OREGON

AND

UNIVERSITY OF MIAMI

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DAVID GILBARG

Stanford University
Stanford, California

R. A. BEAUMONT

University of Washington
Seattle 5, Washington

A. L. WHITEMAN

University of Southern California
Los Angeles 7, California

L. J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

C. E. BURGESS

E. HEWITT

A. HORN

V. GANAPATHY IYER

R. D. JAMES

M. S. KNEBELMAN

L. NACHBIN

I. NIVEN

T. G. OSTROM

H. L. ROYDEN

M. M. SCHIFFER

E. G. STRAUS

G. SZEKERES

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA

OREGON STATE COLLEGE

UNIVERSITY OF OREGON

OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE COLLEGE

UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

CALIFORNIA RESEARCH CORPORATION

HUGHES AIRCRAFT COMPANY

SPACE TECHNOLOGY LABORATORIES

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 2120 Oxford Street, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chivoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Errett Albert Bishop, <i>A minimal boundary for function algebras</i>	629
John W. Brace, <i>The topology of almost uniform convergence</i>	643
Cecil Edmund Burgess, <i>Chainable continua and indecomposability</i>	653
L. Carlitz, <i>Multiplication formulas for products of Bernoulli and Euler polynomials</i>	661
Eckford Cohen, <i>A class of residue systems (mod r) and related arithmetical functions. II. Higher dimensional analogues</i>	667
Shaul Foguel, <i>Boolean algebras of projections of finite multiplicity</i>	681
Richard Robinson Goldberg, <i>Averages of Fourier coefficients</i>	695
Seymour Goldberg, <i>Ranges and inverses of perturbed linear operators</i>	701
Philip Hartman, <i>On functions representable as a difference of convex functions</i> ...	707
Milton Vernon Johns, Jr. and Ronald Pyke, <i>On conditional expectation and quasi-rings</i>	715
Robert Jacob Koch, <i>Arcs in partially ordered spaces</i>	723
Gregers Louis Krabbe, <i>A space of multipliers of type $L^p(-\infty, \infty)$</i>	729
John W. Lamperti and Patrick Colonel Suppes, <i>Chains of infinite order and their application to learning theory</i>	739
Edith Hirsch Luchins, <i>On radicals and continuity of homomorphisms into Banach algebras</i>	755
T. M. MacRobert, <i>Multiplication formulae for the E-functions regarded as functions of their parameters</i>	759
Michael Bahir Maschler, <i>Classes of minimal and representative domains and their kernel functions</i>	763
William Schumacher Massey, <i>On the imbeddability of the real projective spaces in Euclidean space</i>	783
Thomas Wilson Mullikin, <i>Semi-groups of class (C_0) in L_p determined by parabolic differential equations</i>	791
Steven Orey, <i>Recurrent Markov chains</i>	805
Ernest Tilden Parker, <i>On quadruply transitive groups</i>	829
Calvin R. Putnam, <i>On Toeplitz matrices, absolute continuity, and unitary equivalence</i>	837
Helmut Heinrich Schaefer, <i>On nonlinear positive operators</i>	847
Robert Seall and Marion Wetzel, <i>Some connections between continued fractions and convex sets</i>	861
Robert Steinberg, <i>Variations on a theme of Chevalley</i>	875
Olga Taussky and Hans Zassenhaus, <i>On the similarity transformation between a matrix and its transpose</i>	893
Emery Thomas, <i>The suspension of the generalized Pontrjagin cohomology operations</i>	897
Joseph L. Ullman, <i>On Tchebycheff polynomials</i>	913
Richard Steven Varga, <i>Orderings of the successive overrelaxation scheme</i>	925
Orlando Eugenio Villamayor, Sr., <i>On weak dimension of algebras</i>	941