

Pacific Journal of Mathematics

HIGHER DIMENSIONAL CYCLIC ELEMENTS

JOHN MITCHELL GARY

HIGHER DIMENSIONAL CYCLIC ELEMENTS

JOHN GARY

Introduction. Whyburn, in 1934, introduced the higher dimensional cyclic elements [5]. He gave an analysis of the structure of the homology groups of a space in terms of its cyclic elements. His results were for finite dimensional spaces, and he used the integers modulo two as the coefficient group. Puckett generalized some of Whyburn's results to compact metric spaces [3]. Simon has shown that if E is a closed subset of a compact space M , which contains all the $(r-1)$ -dimensional cyclic elements of M , then $H^r(E) \approx H^r(M)$ [4]. He also obtained a direct sum decomposition of $H^r(M)$ using the cyclic elements of M . We will extend some of these results.

The properties of zero-dimensional cyclic elements in locally connected spaces, and the relation of these cyclic elements to monotone mappings, is basic in the applications of zero-dimensional cyclic element theory. We shall give some counter-examples concerning the generalization of these properties to higher dimensional cyclic elements.

1. Preliminaries. Throughout this paper M will always denote a compact Hausdorff space. We shall use the augmented Cech homology and cohomology with a field as coefficient group. Results stated in terms of cohomology may be given a dual expression in terms of homology by means of the dot product duality for the Cech theory.

DEFINITION 1.1. A T_r set in M is a closed subset T of M such that $H^r(K) = 0$, for all closed subsets K of T .

DEFINITION 1.2. An E_r set in M is a non-degenerate subset of M which is maximal with respect to the property that it can not be disconnected by a T_r set of M .

The proofs of Lemmas 1.3 through 1.9 can be found in the papers by Whyburn [5] and Simon [4]. The proofs given by Whyburn are for subsets of Euclidean space, but they can be carried over to our case without difficulty.

LEMMA 1.3. *Let K be a subset of M which can not be disconnected by a T_r set. If $M = M_1 \cup M_2$, T_r -separated (by this we mean M_1 and M_2 are proper closed subsets and $M_1 \cap M_2$ is a T_r set), then $K \subset M_1$ (or, $K \subset M_2$).*

Received December 24, 1958. The results of this paper are contained in the authors doctoral dissertation, University of Michigan, 1956. The author wishes to thank Professor R. L. Wilder for his advice and encouragement.

LEMMA 1.4. *If K is an E_r set, then K is closed and connected.*

LEMMA 1.5. *If K_1 and K_2 are both E_r sets and $K_1 \neq K_2$, then $K_1 \cap K_2$ is a T_r set. Any T_r set is also a T_{r+1} set.*

LEMMA 1.6. *If K is a non-degenerate subset of M , which can not be disconnected by a T_r set, then K is contained in a unique E_r set in M .*

DEFINITION 1.7. If $\gamma^r \in H^r(M)$ and D is a minimal, closed subset of M such that $i^*(\gamma^r) \neq 0$ (where $i^*: H^r(M) \rightarrow H^r(D)$ is the inclusion map), then D is called a *floor* for γ^r .

LEMMA 1.8. *If $\gamma^r \in H^r(M)$ and $\gamma^r \neq 0$, then there exists a floor for γ^r .*

LEMMA 1.9. *If D is a floor for γ^r , then D can not be disconnected by a T_{r-1} set.*

LEMMA 1.10. *If $\{E^1, \dots, E^n\}$ is a finite collection of E_{r-1} sets in M , with $M \neq \bigcup_{i=1}^n E^i$, then there exist proper, closed subsets, M_1 and M_2 , of M such that (1) $M = M_1 \cup M_2$, (2) $M_1 \cap M_2$ is the union of a finite number of T_{r-1} sets (therefore, $M_1 \cap M_2$ is a T_r set), (3) $M_1 \supset \bigcup_{i=1}^n E^i$.*

Proof. The proof will be by induction on n . The case $n = 1$ follows from Lemma 1.3.

Assume the lemma is true up to $n - 1$. Since M is not an E_{r-1} set, we have $M = M_1 \cup M_2$, T_{r-1} -separated. Let $E = \bigcup_{i=1}^n E^i$. If $(M - E) \cap (M - (M_1 \cap M_2)) = \phi$, then the desired T_{r-1} -separation of M could be obtained by using the boundary of an open set in $M_1 \cap M_2$. Therefore, we can assume $(M - E) \cap (M - M_1) \neq \phi$. By Lemma 1.3, we can assume $\bigcup_{i=1}^s E^i \subset M_2$ and $\bigcup_{i=s+1}^n E^i \subset M_1$, where $1 \leq s < n$. We must have $E^i \subset (\overline{M - M_1})$, for $1 \leq i \leq s$. Otherwise, we could separate E^i by the T_{r-1} set $(\overline{M - M_1}) \cap (M_1 \cap M_2)$. Since $(M - E) \cap (M - M_1) \neq \phi$, $(\overline{M - M_1}) \neq \bigcup_{i=1}^s E^i$. Thus, by the induction assumption, $(\overline{M - M_1}) = M_4 \cup M_5$, where $\bigcup_{i=1}^s E^i$ is contained in M_4 and $M_4 \cap M_5$ is the union of a finite number of T_{r-1} sets. If we let $\tilde{M}_1 = M_1 \cup M_4$ and $\tilde{M}_2 = M_5$, then

- (1) $M = \tilde{M}_1 \cup \tilde{M}_2$,
- (2) $\tilde{M}_1 \cap \tilde{M}_2$ is the union of a finite number of T_{r-1} sets,
- (3) $\bigcup_{i=1}^n E^i \subset \tilde{M}_1$,
- (4) \tilde{M}_1 and \tilde{M}_2 are proper closed subsets of M .

2. Cyclic elements and the structure of M .

DEFINITION 2.1. A closed subset A of M is called a L_r set if every E_{r-1} set, whose intersection with A is not a T_r set, is contained in A . The proofs of the following theorems are given below.

THEOREM 2.2. *If A is a L_r set, then $i^*: H^r(M) \rightarrow H^r(A)$ is onto. Thus, by duality, $i_*: H_r(A) \rightarrow H_r(M)$ is one-to-one.*

THEOREM 2.3. *Let A be a closed subset with the following property: if E is an E_{r-1} set and $H^r(E) \neq 0$, then E is contained in A . Then the map $i^*: H^r(M) \rightarrow H^r(A)$ is one-to-one and, by duality, $i_*: H_r(A) \rightarrow H_r(M)$ is onto.*

THEOREM 2.4. *Suppose there are only a finite number, say $\{E^1, \dots, E^n\}$, of E_{r-1} sets such that $H^r(E^i) \neq 0$. Let $A = \bigcup_{i=1}^n E^i$. Then the mappings $i^*: H^r(M) \rightarrow H^r(A)$ and $i_*: H_r(A) \rightarrow H_r(M)$ are isomorphisms.*

REMARK. Theorem 2.4 can not be generalized to an infinite number of E_{r-1} sets, as the following example shows. In Euclidean space let $M = D \cup [\bigcup_{i=1}^{\infty} C_i]$, where $D = \{(x, y, z) | z = 0, x^2 + y^2 \leq 1\}$ and $C_i = \{(x, y, z) | z = 1/i, x^2 + y^2 = 1\}$. We do not have $H_1(\bigcup_{i=1}^{\infty} C_i) \approx H_1(M)$, under the inclusion mapping.

THEOREM 2.5. *Let $\gamma^r \in H^r(M)$ and suppose U is an open set, such that if D is a floor for γ^r , then D is contained in U (see Definition 1.7). Then there exists a $\gamma_u^r \in H^r(M, M - U)$ such that $\gamma^r = j^*(\gamma_u^r)$, where $j^*: H^r(M, M - U) \rightarrow H^r(M)$.*

THEOREM 2.6. *Assume E is an E_{r-1} set in M and N is a closed subset of M , where $N \cap E = \emptyset$. Then the composite mapping $j_* i_*: H_r(E) \rightarrow H_r(M, N)$ is one-to-one. Here, $i_*: H_r(E) \rightarrow H_r(M)$ and $j_*: H_r(M) \rightarrow H_r(M, N)$ are the natural mappings.*

LEMMA 2.7. *Let (M, A) be a compact pair with $\gamma^r \in H^r(A)$. If $\delta^*(\gamma^r) \neq 0$, where $\delta^*: H^r(A) \rightarrow H^{r+1}(M, A)$, then there is a minimal closed set B such that $B \subset A$, and $\delta_B^*(\gamma_B^r) \neq 0$. Here, $\delta_B^*: H^r(B) \rightarrow H^{r+1}(M, B)$ and $\gamma_B^r = i^*(\gamma^r)$, where $i^*: H^r(A) \rightarrow H^r(B)$.*

LEMMA 2.8. *Let B be a minimal set defined in Lemma 2.7. There exists a minimal closed set N such that $\delta^*(\gamma_B^r) \neq 0$, where $\delta^*: H^r(B) \rightarrow H^{r+1}(N, B)$.*

Proof. The proof of these lemmas is obtained from the continuity of the Cech theory and Zorn's lemma.

LEMMA 2.9. *The set N , in Lemma 2.8, can not be disconnected by a T_{r-1} set.*

Proof. Suppose $N = N_1 \cup N_2$, where $N_1 \cap N_2$ is a T_{r-1} set. We will show this to be impossible, unless $N = N_1$. Let B be as defined in Lemma 2.8, and define $B_i = N_i \cap B$ ($i = 1, 2$). We will show that the mapping induced by inclusion

$$K^*: H^{r+1}(N, B) \rightarrow H^{r+1}(N_1, B_1) \oplus H^{r+1}(N_2, B_2)$$

is an isomorphism. We use the relative Mayer-Vietoris sequence given below; note that $T = N_1 \cap N_2$ is a T_{r-1} set [2].

$$\begin{array}{ccc} H^{r+1}(N_2, B_2) \oplus H^{r+1}(N_1, B_1) & \xrightarrow{K^*} & H^{r+1}(N, B) \\ \uparrow i_1^* & & \uparrow i^* \\ H^r(N, N) \rightarrow H^{r+1}(N, N_1 \cup B) + H^{r+1}(N, N_2 \cup B) & \xrightarrow{\bar{K}^*} & H^{r+1}(N, B \cup T) \\ & & \rightarrow H^{r+1}(N, N) . \end{array}$$

The mappings i_1^* and i_2^* are isomorphisms by excision, the map \bar{K}^* by exactness. Using the three exact sequences given below we see that i^* is an isomorphism.

$$\begin{array}{l} H^{s-1}(B \cap T) \rightarrow H^s(B \cup T) \rightarrow H^s(B) \oplus H^s(T) \rightarrow H^s(B \cap T) \\ H^s(B \cup T) \rightarrow H^s(B) \rightarrow H^{s+1}(B \cup T, B) \rightarrow H^{s+1}(B \cup T) \rightarrow H^{s+1}(B) \\ H^r(B \cup T, B) \rightarrow H^{r+1}(N, B \cup T) \rightarrow H^{r+1}(N, B) \rightarrow H^{r+1}(B \cup T, B) . \end{array}$$

The first is a Mayer-Vietoris sequence, the second is a sequence for a pair, the third is a sequence for a triple. Thus K^* is an isomorphism.

In the diagram below, since $\delta_N^*(\gamma_B^r) \neq 0$, we may assume $\delta_{N_1}^* \phi_1^*(\gamma_B^r) \neq 0$.

$$\begin{array}{ccc} H^{r+1}(N, B) & \xrightarrow{K^*} & H^{r+1}(N_1, B_1) \oplus H^{r+1}(N_2, B_2) \\ \uparrow \delta_N^* & \nearrow i_1^* & \uparrow \delta_{N_1}^* \quad \uparrow \delta_{N_2}^* \\ & H^{r+1}(M, B_1) & \\ & \nwarrow \delta_1^* & \\ H^r(B) & \xrightarrow{\phi_1^* \oplus \phi_2^*} & H^r(B_1) \oplus H^r(B_2) \end{array}$$

We now have $\delta_1^* \phi_1^*(\gamma_B^r) \neq 0$, since $i_1^* \delta_1^* \phi_1^*(\gamma_B^r) = \delta_{N_1}^* \phi_1^*(\gamma_B^r) \neq 0$. This implies $B_1 = B$, by the definition of B . Therefore, $\phi_1^*(\gamma_B^r) = \gamma_B^r$ and $\delta_{N_1}^* \phi_1^*(\gamma_B^r) = \delta_N^*(\gamma_B^r) \neq 0$. Since N is minimal, we must have $N_1 = N$. Thus, N can not be disconnected by a T_{r-1} set.

Proof of Theorem 2.2. We will show $\delta^*(\gamma^r) = 0$, for all $\gamma^r \in H^r(A)$, where $\delta^*: H^r(A) \rightarrow H^r(M, A)$. Suppose not; then choose N and B according to Lemma 2.8. Then there exists an E_{r-1} set containing N , by Lemma 1.6. Let E denote this E_{r-1} set. Since E contains N , we have $E \cap A \supset B$. Since $H^r(B) \neq 0$, B is not a T_r set. Therefore $E \subset A$, because A is an L_r set. This implies that N is contained in A . But this is impossible, as the diagram below shows. By the definition of the pair (N, B) , $\delta^*i^*(\gamma^r) \neq 0$.

$$\begin{array}{ccc} H^r(A) & \xrightarrow{\delta^*} & H^{r+1}(A, B) \\ \downarrow i^* & & \downarrow \\ H^r(B) & \xrightarrow{\delta^*} & H^{r+1}(N, B) . \end{array}$$

Proof of Theorem 2.3. Consider the exact sequence:

$$H^r(M, A) \xrightarrow{j^*} H^r(M) \xrightarrow{i^*} H^r(A) .$$

Suppose $j^*(\gamma^r) \neq 0$, where $\gamma^r \in H^r(M, A)$. By Lemmas 1.9 and 1.6 there is an E_{r-1} set which contains a floor for $j^*(\gamma^r)$. Let E be this E_{r-1} set. Since E contains a floor for $j^*(\gamma^r)$, $H^r(E) \neq 0$. Therefore, $E \subset A$; which implies $i^*j^*(\gamma^r) \neq 0$, since E contains a floor for $j^*(\gamma^r)$. Therefore j^* is a trivial map and i^* is one-to-one.

Proof of Theorem 2.4. By Theorem 2.3, $i^*: H_r(A) \rightarrow H_r(M)$ is onto. If $i_*(Z_r) = 0$, for some $Z_r \in H_r(A)$; then there is a minimal set K such that

(1) $K \supset A$, and

(2) $i_*^K(Z_r) = 0$, where $i_*^K: H_r(A) \rightarrow H_r(K)$ [2]. If $K \neq A$; then, by Lemma 1.10, we have $K = K_1 \cup K_2$, T_r -separated. The Mayer-Vietoris sequence below implies $i_*^{K_1}(Z_r) = 0$, where $i_*^{K_1}: H_r(A) \rightarrow H_r(K_1)$.

$$H_r(K_1 \cap K_2) \rightarrow H_r(K_1) \oplus H_r(K_2) \rightarrow H_r(K) .$$


Therefore, $K = A$ and $Z_r = 0$, or i_* is one-to-one.

Proof of Theorem 2.5. Consider the exact sequence,

$$H^r(M, M - U) \xrightarrow{j^*} H^r(M) \xrightarrow{i^*} H^r(M - U) .$$

We will show $i^*(\gamma^r) = 0$, where γ^r is the element of $H^r(M)$ given in the theorem. Suppose $i^*(\gamma^r) \neq 0$; then, by Lemma 1.8, there exists a floor for $i^*(\gamma^r)$ contained in $M - U$. If D is this floor, then D is a floor for γ^r , since $i_D^* = i_{DU}^*i^*$. Here, $i_D^*: H^r(M) \rightarrow H^r(D)$ and $i_{DU}^*: H^r(M - U) \rightarrow H^r(D)$ are inclusion mappings. Therefore, by the definition of U , D is contained in U . This is impossible, hence $i^*(\gamma^r) = 0$.

Proof of Theorem 2.6. Let $\phi_* = j_* i_*$, and suppose $\phi_*(Z_r) = 0$, for some $Z_r \in H_r(E)$. Then there exists a minimal closed set K in M such that $K \supset E$ and $\phi_*^K(Z_r) = 0$, where $\phi_*^K: H_r(E) \rightarrow H_r(K, K \cap N)$ is analogous to ϕ_* defined above. This follows from Zorn's lemma and continuity. We will assume $K \neq E$. Since E is an E_{r-1} set, we can write $K = K_1 \cup K_2$, T_{r-1} -separated. Also, we can assume $E \subset K_1$. Consider the following commutative diagram:



The two vertical sequences are Mayer-Vietoris sequences. Also, the two horizontal sequences are exact. We have

$$H_{r-1}(K_1 \cap K_2) = H_{r-1}(K_1 \cap K_2 \cap N) = H_r(K_1 \cap K_2) = H_r(K_1 \cap K_2 \cap N) = 0,$$

since $K_1 \cap K_2$ is a T_{r-1} set. Since $\phi_*^K(Z_r) = j_*^K i_*^K(Z_r) = 0$, there exists a $Z_r^3 \in H_r(K \cap N)$ such that $i_*^3(Z_r^3) = i_*^K(Z_r)$. There exists

$$(Z_r^1, Z_r^2) \in H_r(K_1 \cap N) \oplus H_r(K_2 \cap N)$$

such that $\psi_*^1(Z_r^1, Z_r^2) = Z_r^3$. By commutativity,

$$\psi_*^2(i_*^1(Z_r^1), i_*^2(Z_r^2)) i_*^3 \psi_*^1(Z_r^1, Z_r^2) = i_*^3(Z_r^3) = i_*^K(Z_r),$$

and

$$\psi_*^*(i_{*1}^K(Z_r), 0) = i_*^K(Z_r).$$

By exactness, ψ_*^2 is an isomorphism, hence $i_*^1(Z_r^1) = i_{*1}^K(Z_r)$. Therefore, $j_{*1}^K i_{*1}^K(Z_r) = j_*^K i_*^1(Z_r^1) = 0$. But this is impossible, since K is minimal. Thus, $K = E$ and ϕ_* is one-to-one.

3. **Cyclic elements in locally connected spaces.** The zero-dimensional cyclic elements in a locally connected continuum have several useful properties. For example, if the continuum M is locally connected, then the zero-dimensional cyclic elements of M are also locally connected and these cyclic elements form a null sequence. Also, the simple 0-links (definition below) are identical with the E_0 sets in an lc^0 space [6]. The examples below show that these properties do not generalize.

DEFINITION 3.1. A non-degenerate subset K of M is called a simple r -link of M , if K is maximal with respect to the following property: if $M = M_1 \cup M_2$, T_r -separated, then $K \subset M_1$ (or $K \subset M_2$). In other words, K is a maximal subset which can not be separated by a T_r set that also separates M .

LEMMA 3.2. All simple r -links in M are closed. If K_1 and K_2 are two distinct simple r -links in M , then $K_1 \cap K_2$ is a T_r set. If L is a non-degenerate subset of M that is not disconnected by any T_r set which also disconnects M , then L is contained in a simple r -link of M .

Proof. The proof is similar to those for the corresponding lemmas for cyclic elements.

EXAMPLE. We will construct an lc^r space M in which the collection of E_r sets does not form a null sequence. This example will also show that, in an lc^r space, the simple r -links need not be the same as the E_r sets.

For each positive integer n , let R_n be a solid, three dimensional rod of height one and diameter $1/2^n$. In Euclidean three-space, define I by $I = \{(x, y, z) | x = 0, y = 0, 0 \leq z \leq 1\}$. Imbed R_n in three-space so that R_n is tangent to R_{n+1} and the sequence of sets R_n converges to I (i.e. $R_n = \{(x, y, z) | x^2 + (y - 3/2^{n+1})^2 \leq 1/2^{2n+2}, 0 \leq z \leq 1\}$). Let M be the set $[\bigcup_{n=1}^{\infty} R_n] \cup I$. Then M is a compact lc^1 space, each R_n is an E_1 set in M , but the collection $\{R_n\}$ is not a null sequence. Also, I is a simple 1-link, but is not an E_1 set.

THEOREM 3.3. If M is s -lc and E is an E_r set of M , where $s \geq r$, then E is s -lc.

Proof. Given any $x \in E$, and an open set U^0 of E containing x , then there exists an open set U of M such that $U \cap E = U^0$. Since M is s -lc, there exists an open set V , containing x , such that $\bar{V} \subset U$ and any compact s -cycle in V bounds on a compact subset of U . Let Z_s be a compact cycle on $V \cap E = V^0$. Then there exists a minimal

closed set K in M such that $\bar{V}^0 \subset K \subset U$, and Z_s bounds on K . By using the Mayer-Vietoris sequence, as it was used in the proof of Theorem 2.4, we can show $K \subset U^0$. Therefore Z_s bounds in U^0 and E is $s - lc$.

EXAMPLE. We will construct a compact lc^r space which contains an E_r set which is not lc^r . Consider the following curve in three-space:

$$x = 0, y = t, z = \sin(\pi/t), \text{ for } 0 < t \leq 1.$$

Expand this curve slightly so that it becomes a solid, three dimensional figure, which oscillates as it approaches the origin. Let N be this space, along with its limiting line segment on the z -axis. Let $P = \{(x, y, z) | x = 0, 0 \leq y \leq 1, -1 \leq z \leq 1\}$; then define $M = P \cup N$. Thus N is an E_1 set in M and M is lc^1 but N is not $0 - lc$.

4. Cyclic elements and monotone mappings. A very basic property of the zero-dimensional cyclic element theory is the following: if $f: M \rightarrow N$ is a monotone mapping (*i.e.* the inverse image of any point is connected), M and N are lc^0 , and E_N is an E_0 set in N ; then there is an E_0 set in M whose image under f contains E_N . This result does not hold in higher dimensions, as the example below demonstrates. The best result we have obtained in this direction is Theorem 4.2.

DEFINITION 4.1. A mapping $f: M \rightarrow N$ is r -monotone, if $H^s(f^{-1}(y)) = 0$, for all $y \in N$ and $0 \leq s \leq r$.

THEOREM 4.2. Let f be an $(r - 1)$ -monotone mapping of M onto N , where M and N are compact Hausdorff spaces. If D_N is a floor for $\gamma_N^r \in H^r(N)$, then there exists a floor D_M for $f^*(\gamma_N^r)$ such that $f(D_M) = D_N$.

Proof. Since f is $(r - 1)$ -monotone, $f^*: H^r(N) \rightarrow H^r(M)$ is a one-to-one mapping [1]. Therefore, $f^*(\gamma_N^r) \neq 0$. Consider the commutative diagram below. The vertical mappings are inclusion mappings; and D_M is defined below.

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ \uparrow i_M & & \uparrow i_N \\ f^{-1}(D_N) & \xrightarrow{f_1} & D_N \\ \uparrow j_M & & \uparrow j_N \\ D_M & \xrightarrow{f_2} & f(D_M) \end{array} .$$

The mapping f_1 is the restriction of f to $f^{-1}(D_N)$. Therefore, f_1 is $(r - 1)$ -monotone. Since D_N is a floor for $\gamma_N^r, i_N^*(\gamma_N^r) \neq 0$. Since

$f_1^*: H^r(D_N) \rightarrow H^r(f^{-1}(D))$ is one-to-one, $i_M^* f^*(\gamma_N^r) = f_1^* i_N^*(\gamma_N^r) \neq 0$. Therefore, $f^{-1}(D_N)$ contains a floor for $f^*(\gamma_N^r)$. Denote this floor by D_M and let f_2 be the restriction of f to D_M . By the definition of a floor, $j_M^* i_M^* f^*(\gamma_M^r) \neq 0$. Since $j_M^* i_M^* f^*(\gamma_N^r) = f_2^* j_N^* i_N^*(\gamma_N^r)$, we have $j_N^* i_N^*(\gamma_N^r) \neq 0$. This implies $f(D_M) = D_N$, since D_N is a floor for γ_N^r .

We shall omit the proofs of Lemmas 4.3 and 4.5.

LEMMA 4.3. *Let N_1 and N_2 be subsets of M which can not be disconnected by a T_r set. Suppose that $\bar{N}_1 \cup \bar{N}_2$ is not a T_r set. Then $\bar{N}_1 \cup \bar{N}_2$ can not be disconnected by a T_r set.*

LEMMA 4.4. *Let $f: M \rightarrow N$, and suppose $T \subset N$ is a T_r set such that $f^{-1}(T)$ is also a T_s set. Also, assume f is a homeomorphism of $M - f^{-1}(T)$ onto $N - T$. Then, if T^N is a T_r set in N , $f^{-1}(T^N)$ is a T_r set in M .*

Proof. Let K be a closed subset of $f^{-1}(T^N)$. Denote $f^{-1}(T)$ by T^{-1} . In the commutative diagram below f_1^* is an isomorphism, by excision. Therefore, by exactness, $H^r(K) = 0$.

$$\begin{CD} H^r(K, K \cap T^{-1}) @>>> H^r(K) @>>> H^r(K \cap T^{-1}) \\ @V f_1^* VV @VV f^* V @VV f^* V \\ H^r(f(K), f(K \cap T^{-1})) @>>> H^r(f(K)) @>>> H^r(f(K) \cap T^{-1}) \end{CD}$$

LEMMA 4.5. *Assume f is a mapping of M onto N such that the inverse image of any T_r set in N is a T_r set in M . If $K \subset M$ can not be disconnected by a T_r set in M , then $f(K)$ can not be disconnected a T_r set in N .*

EXAMPLE. If f is an r -monotone mapping of M onto N , where M and N are lc^∞ spaces and E^N is an E_r set in N ; there may not be an E_r set, E^M , in M such that $f(E^M) \supset E^N$.

We will construct the example in three space. Consider the following solid cylinders:

$$\begin{aligned} M_1 &= \{(x, y, z) \mid x^2 + y^2 \leq 1, 0 \leq z \leq 1\} \\ M_2 &= \{(x, y, z) \mid x^2 + (y - 2)^2 \leq 1, 0 \leq z \leq 1\} . \end{aligned}$$

The cylinders M_1 and M_2 are tangent along $I = \{(x, y, z) \mid x = 0, y = 1, 0 \leq z \leq 1\}$. Let M_3 be an arc joining the endpoints of I , which does not meet $M_1 \cup M_2$ except at these endpoints. Let $M = \bigcup_{i=1}^3 M_i$. We will define a decomposition of M , and will let $f: M \rightarrow N$ be the decomposition mapping.

To form N , identify all the points in M_3 into a single point. Then the mapping $f: M \rightarrow N$ is r -monotone for all r and the restriction of f to $M - M_3$ is a homeomorphism.

We will show that N is an E_1 set. First, neither M_1 nor M_2 can be disconnected by a T_1 set. Lemmas 4.4 and 4.5 imply that neither $f(M_1)$ nor $f(M_2)$ can be disconnected by a T_1 set. By Lemma 4.3, $N = f(M_1) \cup f(M_2)$ can not be disconnected by a T_1 set, since $f(M_1) \cup f(M_2)$ contains an essential 1-cycle. If K is a closed subset of M such that $f(K) \supset N$, then $K \supset M_1 \cup M_2$. Then K can be disconnected by a T_1 set, namely $M_1 \cap M_2$. Therefore, there is no E_1 set in M whose image is N .

Note that M is obviously lc^r for all r . Therefore N is also lc^r , for all r , since f is r -monotone, for all r .

REFERENCES

1. E. G. Begle, *The Vietoris mapping theorem for bicomact spaces*, Annals of Math., (2) **51** (1950), 534-542.
2. Samuel Eilenberg and Norman Steenrod, *Foundations of Algebraic Topology*, Princeton, 1952.
3. W. J. Puckett, Jr., *A theorem concerning homologies in a compact space*, Fundamentae Mathematicae, **31-32** (1938), 32.
4. Arthur Simon, *N -Cyclic elements*, Duke Math. Journal, **24** (1957), 1-7.
5. G. T. Whyburn, *Cyclic elements of higher order*, Amer. J. Math., **56** (1934), 133-146.
6. ———, *Analytic Topology*, Amer. Math. Soc. Coll. Pub., 1942.
7. R. L. Wilder, *Topology of Manifolds*, Amer. Math. Soc. Coll. Pub. 1949.

CALIFORNIA INSTITUTE OF TECHNOLOGY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DAVID GILBARG
Stanford University
Stanford, California

A. L. WHITEMAN
University of Southern California
Los Angeles 7, California

R. A. BEAUMONT
University of Washington
Seattle 5, Washington

L. J. PAIGE
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH
C. E. BURGESS
E. HEWITT
A. HORN

V. GANAPATHY IYER
R. D. JAMES
M. S. KNEBELMAN
L. NACHBIN

I. NIVEN
T. G. OSTROM
H. L. ROYDEN
M. M. SCHIFFER

E. G. STRAUS
G. SZEKERES
F. WOLF
K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON
* * *

AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
HUGHES AIRCRAFT COMPANY
SPACE TECHNOLOGY LABORATORIES

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 2120 Oxford Street, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 9, No. 4

August, 1959

| | |
|--|------|
| Frank Herbert Brownell, III, <i>A note on Kato's uniqueness criterion for Schrödinger operator self-adjoint extensions</i> | 953 |
| Edmond Darrell Cashwell and C. J. Everett, <i>The ring of number-theoretic functions</i> | 975 |
| Heinz Otto Cordes, <i>On continuation of boundary values for partial differential operators</i> | 987 |
| Philip C. Curtis, Jr., <i>n-parameter families and best approximation</i> | 1013 |
| Uri Fixman, <i>Problems in spectral operators</i> | 1029 |
| I. S. Gál, <i>Uniformizable spaces with a unique structure</i> | 1053 |
| John Mitchell Gary, <i>Higher dimensional cyclic elements</i> | 1061 |
| Richard P. Gosselin, <i>On Diophantine approximation and trigonometric polynomials</i> | 1071 |
| Gilbert Helmsberg, <i>Generating sets of elements in compact groups</i> | 1083 |
| Daniel R. Hughes and John Griggs Thompson, <i>The H-problem and the structure of H-groups</i> | 1097 |
| James Patrick Jans, <i>Projective injective modules</i> | 1103 |
| Samuel Karlin and James L. McGregor, <i>Coincidence properties of birth and death processes</i> | 1109 |
| Samuel Karlin and James L. McGregor, <i>Coincidence probabilities</i> | 1141 |
| J. L. Kelley, <i>Measures on Boolean algebras</i> | 1165 |
| John G. Kemeny, <i>Generalized random variables</i> | 1179 |
| Donald G. Malm, <i>Concerning the cohomology ring of a sphere bundle</i> | 1191 |
| Marvin David Marcus and Benjamin Nelson Moysl, <i>Transformations on tensor product spaces</i> | 1215 |
| Charles Alan McCarthy, <i>The nilpotent part of a spectral operator</i> | 1223 |
| Kotaro Oikawa, <i>On a criterion for the weakness of an ideal boundary component</i> | 1233 |
| Barrett O'Neill, <i>An algebraic criterion for immersion</i> | 1239 |
| Murray Harold Protter, <i>Vibration of a nonhomogeneous membrane</i> | 1249 |
| Victor Lenard Shapiro, <i>Intrinsic operators in three-space</i> | 1257 |
| Morgan Ward, <i>Tests for primality based on Sylvester's cyclotomic numbers</i> | 1269 |
| L. E. Ward, <i>A fixed point theorem for chained spaces</i> | 1273 |
| Alfred B. Willcox, <i>Šilov type C algebras over a connected locally compact abelian group</i> | 1279 |
| Jacob Feldman, <i>Correction to "Equivalence and perpendicularity of Gaussian processes"</i> | 1295 |