

Pacific Journal of Mathematics

UNIONS OF CELL PAIRS IN E^3

P. H. DOYLE, III

UNIONS OF CELL PAIRS IN E^3

P. H. DOYLE

In [4] it is shown that there are pairs of cells of all dimensions possible in euclidean 3-space, E^3 , which are tame separately, but which have a wild set as their union. Such pairs can be constructed when the individual cells intersect in a single point. The present paper gives conditions that unions of some such pairs be tame sets as well as a number of other results.

LEMMA 1. *Let D_1 be a disk which is polyhedral and which lies on the boundary, ∂T , of a tetrahedron T in E^3 . If D_2 is a disk in E^3 which has a polygonal boundary and is locally polyhedral mod ∂D_2 while $D_2 \cap T = D_2 \cap D_1 = \partial D_2 \cap \partial D_1 = J$, an arc, then $D_1 \cup D_2$ is a tame disk.*

Proof. Let P_1 and P_2 be polyhedral disks in ∂T , $P_1 \cap P_2 = \square$ and $(P_1 \cup P_2) \cap D_1 = \square$. Then $\overline{\partial T \setminus (P_1 \cup P_2)}$ is a polyhedral annulus, A_1 . If Q is a polyhedral disk in $D_2 \setminus \partial D_2$, then $\overline{D_2 \setminus Q}$ is an annulus A_2 which is locally polyhedral mod ∂D_2 . By applying Lemma 5.1 of [8] to A_1 and A_2 one obtains a space homeomorphism h carrying E^3 onto E^3 while $h(D_1 \cup D_2)$ is a polyhedral set. This completes the proof of Lemma 1.

LEMMA 2. *Let D_1 be the disk of Lemma 1 while D_2 is a tame disk in E^3 such that $D_2 \cap T = D_2 \cap D_1 = \partial D_2 \cap \partial D_1 = J$, an arc. Then $\partial T \cup \partial D_2$ is tame.*

Proof. By Theorem 2 of [3] $\partial D_1 \cup \partial D_2$ is locally tame and hence tame by [1] or [8]. Let a be a point of ∂J and J' be an interval of ∂D_1 having a as an end point and $J' \cap \partial D_2 = a$. We choose a polygonal disk M on ∂T with $(J'/\partial J')$ in its interior while $\partial D_1 \cap M = J'$. By a swelling [5] of M toward the component of $E^3 \setminus \partial T$ which meets ∂D_2 we obtain a disk M' which is locally polyhedral mod ∂M and $M' \cap \partial T = \partial M = \partial M'$. The sphere $S = M' \cup (\partial T \setminus M)$ is tame by [8] and S is pierced at a by a tame arc lying on $\partial(D_1 \cup D_2)$. Hence by [7] $\partial D_2 \cup S$ is locally tame at a . We select an arc P in $(S \setminus M') \cup a$ which is locally polyhedral except at the point a . There is an arc A on ∂D_2 which lies in the exterior of S except for its end point a . The arc $A \cup P$ is tame since $S \cup \partial D_2$ is tame. Let the arc P be swollen into a 3-cell C^3 with P in its interior such that C^3 is locally polyhedral mod a , $C^3 \cap S$ is a disk while $C^3 \cap M = a$. Then ∂C^3 is pierced at a by $A \cup P$ and so $A \cup P \cup \partial C^3$ is tame by [7]. Evidently there is an arc P' on ∂C^3 so

Received April 27, 1959. The work on part of this paper was supported by the National Science Foundation Grant G-2793.

that $A \cup P'$ pierces ∂T at a . Again by [7] $\partial D_2 \cup \partial T$ is locally tame at a . A similar argument applies to the other end point of ∂J . Hence $\partial D_2 \cup \partial T$ is tame. This proves Lemma 2.

THEOREM 1. *Let D_1 and D_2 be two tame disks in E^3 such that $D_1 \cap D_2 = \partial D_1 \cap \partial D_2 = J$, an arc. Then $D_1 \cup D_2$ is a tame disk.*

Proof. Since D_1 is tame there is a homeomorphism h_1 of E^3 onto E^3 such that $h_1(D_1)$ is a plane triangle. The disk $h_1(D_1)$ is to be swollen so that a 3-cell e^3 is formed such that

- (i) $h_1(D_1) \subset \partial e^3$,
- (ii) e^3 is tame,
- (iii) and $e^3 \cap h_1(D_2) = h_1(J)$.

That such a cell e^3 exists follows from Lemma 5.1 of [5] and Theorem 9.3 of [8].

There is a homeomorphism h_2 of E^3 onto E^3 which carries ∂e^3 and $h_1(D_1)$ onto the boundary of a tetrahedron and a polyhedral disk, respectively. By Lemma 2 $h_2(e^3) \cup h_2 h_1(\partial D_2)$ is a tame set. By Theorem 2 of [6] we can insist that $h_2 h_1(D_2)$ be locally polyhedral mod $h_2 h_1(\partial D_2)$, while $h_2 h_1(\partial D_2)$ is polygonal. Hence by Lemma 1 $h_2 h_1(D_1 \cup D_2)$ is tame and so $D_1 \cup D_2$ is tame.

The following result gives a characterization of tame 1-dimensional complexes in E^3 . By a 1_n -star we mean a homeomorphic image of a 1-dimensional simplicial complex K with a vertex x whose star is K and x is the common end point of the n segments meeting only in x .

THEOREM 2. *If N is a 1_n -star in E^3 such that $(n - 1)$ of the branches of N lie on a disk D which meets the remaining branch J at x only and if each arc in N is tame, then N is tame.*

Proof. By [2] we may assume that D is locally polyhedral mod N . An application of the method in Theorem 1 of [3] makes it possible to select a subset D' of D which is a disk consisting of $(n - 1)$ tame disks which contain arcs with x as an end point of all branches of N except J . An argument almost identical with that of Theorem 2 of [3] suffices to show that $J \cup D'$ is tame and hence N is tame by [1] or [8].

COROLLARY 1. *Let G be a graph in E^3 such that the star of each vertex of G meets the conditions of Theorem 2, then G is tame. The conditions are evidently necessary as well.*

COROLLARY 2. *Let D be a tame disk and J a tame arc in E^3 . If $D \cap J = \partial D \cap J = p$, an end point of J , and if $\partial D \cup J$ is tame, then $D \cup J$ is tame.*

Proof. Since D is tame there is a space homeomorphism h which

carries D onto a face of a tetrahedron T , $[h(J)\setminus h(p)] \subset E^3 \setminus T$. Let P be a segment on $h(\partial D)$ with $h(p)$ as an end point. We enclose P in a polyhedral disk M in ∂T such that P spans M and $h(\partial D) \cap M = P$. We swell M as in Lemma 2 to obtain a tame disk M' such that $\partial M' = \partial M$, and $M' \setminus \partial M' \subset E^3 \setminus T$. Then $h(J) \cup h(\partial D)$ contains a tame arc which pierces the tame sphere [8] $S = M' \cup (\partial T \setminus M)$ at $h(p)$ and so $S \cup h(J)$ is tame by [7]. The construction of an arc P' as in Lemma 2 completes the proof.

In Example 1.4 of [4] an arc A which is the union of two tame arcs is shown. Although A has an open 3-cell complement in compactified E^3 , it is nevertheless wild. A similar example can be obtained from Example 1.4 of two tame disks which meet at a point on the boundary of each and which have a wild union. In this connection we give the following result.

THEOREM 3. *Let D_1 and D_2 be disks in E^3 such that each arc in D_1 and D_2 is tame and $D_1 \cap D_2 = \partial D_1 \cap \partial D_2 = J$, an arc. Then $D_1 \cup D_2$ is a disk such that each arc in $D_1 \cup D_2$ is tame.*

Proof. Let J' be an arc in $D_1 \cup D_2$. If $\partial J'$ does not lie in $\partial D_1 \cup \partial D_2$ we extend J' so that this is the case, obtaining $J'' \supset J'$, $\partial J'' \subset \partial D_1 \cup \partial D_2$ and $J'' \subset D_1 \cup D_2$. By [2] there is a disk D such that $\partial D = \partial(D_1 \cup D_2)$, $J \cup J'' \subset D$ and D is locally polyhedral mod $J \cup J'' \cup \partial D$. The arc J in D is the intersection of two disks in D , D'_1 and D'_2 , such that $D'_1 \cup D'_2 = D$. Consider any point x of J'' in $D'_1 \setminus \partial D'_1$. In [3] a method is given for enclosing x in the interior of a tame subdisk of D'_1 . Hence D'_1 is locally tame at each of its interior points and $\partial D'_1$ is tame. By [8] D'_1 is tame. A similar argument can be applied to D'_2 . Hence $D'_1 \cup D'_2$ is a tame disk by Theorem 2. Then J'' is tame and so J' is tame. Since J' was arbitrarily chosen $D_1 \cup D_2$ is a disk in which each arc is tame.

COROLLARY 1. *Let L_1 and L_2 be tame disks which intersect in a single point on the boundary of each. If $L_1 \cup L_2$ lies on a disk in which each arc is tame, then $L_1 \cup L_2$ is tame.*

Proof. Let $L_1 \cup L_2$ lie on a disk D such that each arc in D is tame. By Theorem 2 $\partial L_1 \cup \partial L_2$ is tame. There is a disk D' in D with a tame boundary such that $D' \cap (L_1 \cup L_2) \subset \partial L_1 \cup \partial L_2$ while $D' \cup L_1 \cup L_2$ is a disk. Then by [2] there is a disk D'' such that $\partial D'' = \partial D'$, D'' is locally polyhedral mod $\partial D''$ and $\partial D'' \cap (L_1 \cup L_2) = \partial D' \cap (L_1 \cup L_2)$. Now D'' is tame by [8] and so $D'' \cup L_1 \cup L_2$ is tame by Theorem 2. It follows that $L_1 \cup L_2$ is tame.

REFERENCES

1. R. H. Bing, *Locally tame sets are tame*, Ann. of Math. **59** (1954), 145-158.

2. R. H. Bing, *Approximating surfaces with polyhedral ones*, Ann. of Math. **65** (1957), 456-483.
3. P. H. Doyle, *Tame triods in E^3* , Proc. Amer. Math. Soc. **10** (1959), 656-658.
4. R. H. Fox and E. Artin, *Some wild cells and spheres in three-dimensional space*, Ann. of Math. **49** (1948), 979-990.
5. O. G. Harrold, H. C. Griffith, and E. E. Posey, *A Characterization of tame curves in three-space*, Trans. Amer. Math. Soc. **79** (1955), 12-34.
6. E. E. Moise, *Affine structures in 3-manifolds, V. The triangulation theorem and Hauptvermutung*, Ann. of Math. **56** (1952), 96-114.
7. ———, *Affline structures in 3-manifolds, VII. Disk which are pierced by intervals*, Ann. of Math. **58** (1953), 403-408.
8. ———, *Affline structures in 3-manifolds, VIII. Invariance of the knot-types; Local tame imbedding*, Ann. of Math. **59** (1954), 159-170.

MICHIGAN STATE UNIVERSITY
UNIVERSITY OF TENNESSEE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DAVID GILBARG

Stanford University
Stanford, California

F. H. BROWNELL

University of Washington
Seattle 5, Washington

A. L. WHITEMAN

University of Southern California
Los Angeles 7, California

L. J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

T. M. CHERRY

D. DERRY

E. HEWITT

A. HORN

L. NACHBIN

M. OHTSUKA

H. L. ROYDEN

M. M. SCHIFFER

E. SPANIER

E. G. STRAUS

F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
HUGHES AIRCRAFT COMPANY
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 2120 Oxford Street, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Maynard G. Arsove, <i>The Paley-Wiener theorem in metric linear spaces</i>	365
Robert (Yisrael) John Aumann, <i>Acceptable points in games of perfect information</i>	381
A. V. Balakrishnan, <i>Fractional powers of closed operators and the semigroups generated by them</i>	419
Dallas O. Banks, <i>Bounds for the eigenvalues of some vibrating systems</i>	439
Billy Joe Boyer, <i>On the summability of derived Fourier series</i>	475
Robert Breusch, <i>An elementary proof of the prime number theorem with remainder term</i>	487
Edward David Callender, Jr., <i>Hölder continuity of n-dimensional quasi-conformal mappings</i>	499
L. Carlitz, <i>Note on Alder's polynomials</i>	517
P. H. Doyle, III, <i>Unions of cell pairs in E^3</i>	521
James Eells, Jr., <i>A class of smooth bundles over a manifold</i>	525
Shaul Foguel, <i>Computations of the multiplicity function</i>	539
James G. Glimm and Richard Vincent Kadison, <i>Unitary operators in C^*-algebras</i>	547
Hugh Gordon, <i>Measure defined by abstract L_p spaces</i>	557
Robert Clarke James, <i>Separable conjugate spaces</i>	563
William Elliott Jenner, <i>On non-associative algebras associated with bilinear forms</i>	573
Harold H. Johnson, <i>Terminating prolongation procedures</i>	577
John W. Milnor and Edwin Spanier, <i>Two remarks on fiber homotopy type</i>	585
Donald Alan Norton, <i>A note on associativity</i>	591
Ronald John Nunke, <i>On the extensions of a torsion module</i>	597
Joseph J. Rotman, <i>Mixed modules over valuations rings</i>	607
A. Sade, <i>Théorie des systèmes demosiens de groupoi des</i>	625
Wolfgang M. Schmidt, <i>On normal numbers</i>	661
Berthold Schweizer, Abe Sklar and Edward Oakley Thorp, <i>The metrization of statistical metric spaces</i>	673
John P. Shanahan, <i>On uniqueness questions for hyperbolic differential equations</i>	677
A. H. Stone, <i>Sequences of coverings</i>	689
Edward Oakley Thorp, <i>Projections onto the subspace of compact operators</i>	693
L. Bruce Treybig, <i>Concerning certain locally peripherally separable spaces</i>	697
Milo Wesley Weaver, <i>On the commutativity of a correspondence and a permutation</i>	705
David Van Vranken Wend, <i>On the zeros of solutions of some linear complex differential equations</i>	713
Fred Boyer Wright, Jr., <i>Polarity and duality</i>	723