

Pacific Journal of Mathematics

ON NORMAL NUMBERS

WOLFGANG M. SCHMIDT

ON NORMAL NUMBERS

WOLFGANG SCHMIDT

1. Introduction. A real number ξ , $0 \leq \xi < 1$, is said to be *normal in the scale of r* (or *to base r*), if in $\xi = 0 \cdot a_1 a_2 \dots$ expanded in the scale of $r^{(1)}$ every combination of digits occurs with the proper frequency. If $b_1 b_2 \dots b_k$ is any combination of digits, and Z_N the number of indices i in $1 \leq i \leq N$ having

$$b_1 = a_i, \dots, b_k = a_{i+k-1},$$

then the condition is that

$$(1) \quad \lim_{N \rightarrow \infty} Z_N N^{-1} = r^{-k}.$$

A number ξ is called *simply normal* in the scale of r if (1) holds for $k = 1$. A number is said to be *absolutely normal* if it is normal to every base r . It is well-known (see, for example, [6], Theorem 8.11) that almost every number ξ is absolutely normal.

We write $r \sim s$, if there exist integers n, m with $r^n = s^m$. Otherwise, we put $r \not\sim s$.

In this paper we solve the following problem. *Under what conditions on r, s is every number ξ which is normal to base r also normal to base s ?* The answer is given by

THEOREM 1. *A Assume $r \sim s$. Then any number normal to base r is normal to base s .*

B If $r \not\sim s$, then the set of numbers ξ which are normal to base r but not even simply normal to base s has the power of the continuum.

The A-part of the Theorem is rather trivial, but I shall sketch a proof of it, since I could not find one in the literature.

Next, let I be an interval of length $|I|$ contained in the unit-interval $U = [0, 1]$. We write $M_N(\xi, r, I)$ for the number of indices i in $1 \leq i \leq N$ such that the fractional part $\{r^i \xi\}$ of $r^i \xi$ lies I . A sequence $\xi, r\xi, r^2\xi, \dots$ has *uniform distribution modulo 1* if

$$R_N(\xi, r, I) = M_N(\xi, r, I) - N|I| = o(N)$$

for any I . It was proved by Wall [8] (the most accessible proof in [6], Theorem 8.15) that ξ is normal to base r if and only if $\xi, r\xi, r^2\xi, \dots$ has uniform distribution modulo 1.

Write $T_{s,t}$, where $1 < t < s$, for the following mapping in U : If $\xi = 0 \cdot a_1 a_2 \dots$ in the scale of t , then $T_{s,t}\xi = 0 \cdot a_1 a_2 \dots$ in the scale of s .

Received June 2, 1959.

¹ In case of ambiguity we take the representation with an infinity of a_i less than $r - 1$. But this does not affect the property of ξ to be normal or not.

THEOREM 2. *Assume $r \not\sim s$. Then there exists a constant $\alpha_1 = \alpha_1(r, s, t) > 0$ such that for almost every ξ there exists a $N_0(\xi)$ with*

$$(2) \quad R_N(T_{s,t}\xi, r, I) \leq N^{1-\alpha_1}$$

for every $N \geq N_0(\xi)$ and any I .

Thus $T_{s,t}\xi$ is normal to base r for almost all ξ . Since $T_{s,t}\xi$ is not simply normal to base s part B of Theorem 1 follows. It does not follow immediately for $s = 2$, but instead of $T_{2,t}$, which does not exist, we may take $T_{4,t}$.

We can interpret our results as follows. Write $C_{s,t}$ for the image set $T_{s,t}U$ of the unit-interval U under the mapping $T_{s,t}$. $C_{s,t}$ is essentially a Cantor set. In $C_{s,t}$ we define a measure $\mu_{s,t}$ by

$$(3) \quad \int_{C_{s,t}} f(\xi) d\mu_{s,t} = \int_0^1 f(T_{s,t}\xi) d\xi,$$

where $f(\xi)$ is any real-valued function such that the integral on the right hand side of (3) exists. Then it follows from Theorem 2 that with respect to $\mu_{s,t}$ almost every ξ in $C_{s,t}$ is normal in the scale of r .

Throughout this paper, lower case italics stand for integers. $\alpha_1 = \alpha_1(r, s, t), \alpha_2, \alpha_3, \dots$ will be positive constants depending on some or all the variables r, s, t .

1. The case $r \sim s$. First, it follows almost from the definition that any number normal to base s^m is normal to base s .

Next, assume ξ is normal to base r , we shall show it is normal in the scale of r^m . If $\xi = 0.a_1a_2\dots$ in the scale of $r, b_1\dots b_{mk}$ is any combination of mk digits and $Z_N^{(1)}$ is the number of indices i in $1 \leq i \leq N$ with $i \equiv 1 \pmod m$ satisfying

$$b_1 = a_i, \dots, b_{mk} = a_{i+mk-1},$$

then it was shown in [7] and in [3] that

$$\lim_{N \rightarrow \infty} Z_N^{(1)} N^{-1} = r^{-mk} m^{-1}$$

and hence

$$\lim_{N \rightarrow \infty} Z_{mN}^{(1)} N^{-1} = (r^m)^{-k}.$$

Thus ξ is normal to base r^m .

Combining the above remarks we obtain the A-part of Theorem 1.

2. The measure $\mu_{s,t}$. We define *numbers of order h* to be the number $0.a_1\dots a_h$ with $0 \leq a_i < t$ in the scale of s . There are t^h numbers of order h , we denote them in ascending order by $\theta_1^{(h)}, \dots, \theta_{t^h}^{(h)}$.

LEMMA 1. Let $f(\xi)$ be a step-function, having a finite number of steps. Then

$$\int_{\sigma_{s,t}} f(\xi) d\mu_{s,t} = \int_0^1 f(T_{s,t}\xi) d\xi = \lim_{h \rightarrow \infty} t^{-h} \sum_{k=1}^{t^h} f(\theta_k^{(h)}) .$$

The integrals and the limit exist and are finite.

Proof. It will be sufficient to prove the lemma for $f(\xi) = \{\xi, \gamma\}$, where $0 \leq \gamma \leq 1$ and

$$\{\xi, \gamma\} = \begin{cases} 1, & \text{if } \{\xi\} < \gamma \\ 0 & \text{otherwise.} \end{cases}$$

$\xi_k^{(h)} = \int_0^1 \{T_{s,t}\xi, \theta_k^{(h)}\} d\xi$ is the least upper bound of numbers ξ having $T_{s,t}\xi \leq \theta_k^{(h)}$. Thus if $\theta_k^{(h)} = 0 \cdot a_1 \cdots a_h$ in the scale of s , then $\xi_k^{(h)} = 0 \cdot a_1 \cdots a_h$ in the scale of t and therefore $\xi_k^{(h)} = (k - 1)t^{-h}$.

Hence if $\theta_k^{(h)} \leq \gamma \leq \theta_{k+1}^{(h)}$, or if $\theta_k^{(h)} \leq \gamma$ with $k = t^h$, then

$$\int_0^1 \{T_{s,t}\xi, \gamma\} d\xi = kt^{-h} - \varepsilon ,$$

where $0 \leq \varepsilon \leq t^{-h}$. We can rewrite this in the form

$$\int_0^1 \{T_{s,t}\xi, \gamma\} d\xi = t^{-h} \sum_{k=1}^{t^h} \{\theta_k^{(h)}, \gamma\} - \varepsilon ,$$

and Lemma 1 follows.

Particularly, for

$$\begin{aligned} \mu(\gamma, x) &= \int_0^1 \{xT_{s,t}\xi, \gamma\} d\xi \\ \mu(\gamma, x, y) &= \int_0^1 \{xT_{s,t}\xi, \gamma\} \{yT_{s,t}\xi, \gamma\} d\xi \end{aligned}$$

we have

$$(4) \quad \mu(\gamma, x) = \lim_{h \rightarrow \infty} t^{-h} \sum_{k=1}^{t^h} \{x\theta_k^{(h)}, \gamma\} ,$$

$$(5) \quad \mu(\gamma, x, y) = \lim_{h \rightarrow \infty} t^{-h} \sum_{k=1}^{t^h} \{x\theta_k^{(h)}, \gamma\} \{y\theta_k^{(h)}, \gamma\} .$$

3. Exponential sums. Write $e(\xi)$ for $e^{2\pi i \xi}$. There exist ([5], pp. 91–92, 99) for any $\gamma, 0 \leq \gamma \leq 1$, and any $\eta > 0$ functions $f_1(\xi), f_2(\xi)$ periodic in ξ with period 1, such that $f_1(\xi) \leq \{\xi, \gamma\} \leq f_2(\xi)$, having Fourier expansions

$$f_1(\xi) = \gamma - \eta + \sum'_u A_u^{(1)} e(u\xi)$$

$$f_2(\xi) = \gamma + \eta + \sum'_u A_u^{(2)} e(u\xi) ,$$

where the summation is over all $u \neq 0$ and $A_u^{(i)}$ is majorized by

$$(6) \quad |A_u| \leq \frac{1}{u^2 \eta} .$$

Applying this to (5) we obtain

$$\mu(\gamma, x, y) \leq (\gamma + \eta)^2 + \overline{\lim}_{h \rightarrow \infty} t^{-h} \sum'_{\substack{u, v \\ \neq 0, 0}} \left| A_u^{(2)} \| A_v^{(2)} \right| \sum_{k=1}^{t^h} e((ux + vy)\theta_k^{(h)}) \Big| ,$$

where we put $A_0^{(2)} = \gamma + \eta$ and take the sum over all pairs u, v of numbers not both being zero. Since

$$\left| t^{-h} \sum_{k=1}^{t^h} e((ux + vy)\theta_k^{(h)}) \right| \leq 1 ,$$

and since the double sum over u, v is uniformly convergent in h , we may change the order of limit and summation and obtain

$$\mu(\gamma, x, y) \leq (\gamma + \eta)^2 + \sum'_{u, v} |A_u^{(2)}| |A_v^{(2)}| \overline{\lim}_{h \rightarrow \infty} t^{-h} \left| \sum_{k=1}^{t^h} e((ux + vy)\theta_k^{(h)}) \right| .$$

The numbers $\theta_k^{(h)}$ are the numbers

$$\frac{a_1}{s} + \frac{a_2}{s^2} + \dots + \frac{a_h}{s^h} ,$$

where $0 \leq a_i < t$. Hence

$$\sum_{k=1}^{t^h} e(w\theta_k^{(h)}) = \prod_{j=1}^h \left(1 + e\left(\frac{w}{s^j}\right) + e\left(\frac{2w}{s^j}\right) + \dots + e\left(\frac{(t-1)w}{s^j}\right) \right) .$$

If we keep w fixed, and if j is large, then

$$\left| \left(1 + e\left(\frac{w}{s^j}\right) + \dots + e\left(\frac{(t-1)w}{s^j}\right) \right) t^{-1} - 1 \right| < \frac{t|w|}{s^i} .$$

Therefore

$$(7) \quad \Pi(s, t; w) = \prod_{j=1}^{\infty} \left| \left(1 + e\left(\frac{w}{s^j}\right) + \dots + e\left(\frac{(t-1)w}{s^j}\right) \right) t^{-1} \right|$$

exists and

$$(8) \quad \mu(\gamma, x, y) \leq (\gamma + \eta)^2 + \sum'_{u, v} |A_u^{(2)}| |A_v^{(2)}| \Pi(s, t; ux + vy) .$$

The next three sections will be devoted to finding bounds for sums like

$$\sum_{N_1 < n, m \leq N_2} \Pi(s, t; ur^n + vr^m) .$$

4. Two lemmas on digits.

LEMMA 2. Write $w = c_j \cdots c_2 c_1$ in the scale of s . Assume there are at least z pairs of digits $c_{i+1}c_i$ with

$$(9) \quad 1 \leq c_{i+1}c_i \leq s^2 - 2.$$

(Here $c_{i+1}c_i = sc_{i+1} + c_i$). Then

$$H(s, t; w) \leq \alpha_2^z,$$

where $\alpha_2 = \alpha_2(s, t)$, $0 < \alpha_2 < 1$.

Proof. There are at least z numbers i having

$$\frac{1}{s^2} \leq \left\{ \frac{w}{s^i} \right\} \leq 1 - \frac{1}{s^2}.$$

For such an i we have

$$\left| 1 + e\left(\frac{w}{s^i}\right) + \cdots + e\left(\frac{(t-1)w}{s^i}\right) \right| \leq \left| 1 + e\left(\frac{1}{s^2}\right) \right| + t - 2 = t\alpha_2$$

and the Lemma is proved.

There exists an $\alpha_3(s)$, $0 < \alpha_3 < 1/4$, such that

$$\frac{(s^2 - 2)^{\alpha_3} 2^{1/2 - \alpha_3}}{(2\alpha_3)^{\alpha_3} (1 - 2\alpha_3)^{1/2 - \alpha_3}} < 2^{3/4}.$$

LEMMA 3. If k is large, $k > \alpha_3(s)$, then the number of combinations of digits $c_k c_{k-1} \cdots c_1$ in the scale of s with less than $\alpha_3(s)k$ indices i satisfying (9) is not greater than $2^{(3/4)k}$.

Proof. It will be sufficient to show that the number of combinations with less than $\alpha_3(s)k$ indices i satisfying both (9) and $i \equiv 1 \pmod{2}$ is not greater than $2^{(3/4)k}$. We first assume k is even. There exist

$$\binom{k}{2} \binom{k}{l} (s^2 - 2)^l 2^{k/2 - l}$$

combinations $c_k \cdots c_1$ with exactly l indices i having both (9) and $i \equiv 1 \pmod{2}$. Hence the number of combinations with less than $\alpha_3(s)k$ indices i satisfying (9) and $i \equiv 1 \pmod{2}$ does not exceed

$$k \binom{k}{[\alpha_3 k]} (s^2 - 2)^{[\alpha_3 k]} 2^{(k/2) - [\alpha_3 k]}.$$

Using Stirling's formula for the binomial coefficient we obtain for large enough k the upper bound

$$\alpha_5(s)k \frac{(s^2 - 2)^{\alpha_3 k} 2^{((1/2) - \alpha_3)k}}{(2\alpha_3)^{\alpha_3 k} (1 - 2\alpha_3)^{((1/2) - \alpha_3)k}} < 2^{(3/4)k} .$$

Actually, the expression on the left hand side is $< 2^{\alpha_6 k}$, where $\alpha_6 < 3/4$. This permits us to extend the result to odd k .

5. The order of r modulo p^k as a function of k .

LEMMA 4. *Assume p is a prime with $p \nmid r$. Then the order $o(r, p^k)$, of r modulo p^k satisfies*

$$o(r, p^k) \geq \alpha_7(r, p)p^k .$$

COROLLARY. *Let n run through a residue system modulo p^k . Then at most $\alpha_8(r, p)$ of the numbers r^n will fall into the same residue class modulo p^k .*

Proof. Write

$$g = g(p) = \begin{cases} p - 1, & \text{if } p \text{ is odd} \\ 2, & \text{if } p = 2. \end{cases}$$

There exists an $\alpha_9 = \alpha_9(r, p)$ such that

$$(10) \quad r^g \equiv 1 + qp^{\alpha_9 - 1} \pmod{p^{\alpha_9}} ,$$

where $q \not\equiv 0 \pmod{p}$. We have necessarily $\alpha_9 > 1$ and even $\alpha_9 > 2$ if $p = 2$. It follows from (10) by standard methods (see, for instance, [4], § 5.5) that

$$r^{g p^e} \equiv 1 + qp^{\alpha_9 - 1 + e} \pmod{p^{\alpha_9 + e}}$$

for any $e \geq 0$. Thus for $k \geq \alpha_9$ we have

$$r^{g p^{k - \alpha_9}} \equiv 1 + qp^{k - 1} \pmod{p^k}$$

and

$$o(r, p^k) \geq gp^{k - \alpha_9} = \alpha_7(r, p)p^k .$$

Assume $r \not\sim s$. Write

$$\begin{aligned} r &= p_1^{d_1} p_2^{d_2} \cdots p_h^{d_h} \\ s &= p_1^{e_1} p_2^{e_2} \cdots p_h^{e_h} , \end{aligned}$$

where we may assume that never both $d_i = 0, e_i = 0$. We also may assume that the primes p_1, \dots, p_h are ordered in such a way that

$$\frac{e_1}{d_1} \geq \frac{e_2}{d_2} \geq \dots \geq \frac{e_h}{d_h},$$

where we put $(e_i/d_i) = +\infty$ if $d_i = 0$. Since $r \not\sim s$, we have

$$r_1 = \frac{r^{e_1}}{s^{d_1}} > 1.$$

From now on, $p = p_1(r, s)$ is the prime defined above. We have $p \mid s$ but $p \nmid r_1$. For any $x \neq 0$, $y > 1$ we define two new numbers x_y and x'_y by $x = x_y x'_y$, where x_y is a power of y and $y \nmid x'_y$.

LEMMA 5. A. Assume $r \not\sim s$, $v \neq 0$. Let m run through a system $K(s^k)$ of non-negative representatives modulo s^k . Then at most

$$\alpha_{10}(r, s) \left(\frac{s}{2}\right)^k v_p$$

of the numbers

$$v(r^m)'_s$$

are in the same residue class modulo s^k .

B. Assume $r \not\sim s$, furthermore $p \nmid r$. Suppose $u \neq 0$, $v \neq 0$, n are fixed. Then, if m runs through $K(s^k)$, at most

$$\alpha_{11}(r, s) \left(\frac{s}{2}\right)^k v_p$$

of the numbers

$$ur^n + vr^m$$

will fall into the same residue class modulo s^k .

Proof. A. Write $m = m_1 e_1 + m_2$, $0 \leq m_2 < e_1$. Then $r^m = r^{m_1 e_1 + m_2} = s^{m_1 d_1} r_1^{m_1} r^{m_2}$ and $v(r^m)'_s = v r_1^{m_1} (r^{m_2})'_s$. The equation

$$r_1^{m_1} \equiv a \pmod{p^k}$$

has for fixed a at most $e_1 \alpha_8(r_1, p)$ solutions in $m = m_1 e_1 + m_2$, if m runs through a system $K(p^k)$ of residues modulo p^k . This follows from the corollary of Lemma 4. The equation

$$av(r^{m_2})'_s \equiv b \pmod{p^k}$$

has for fixed b, m_2 at most

$$\text{g.c.d.}(v(r^{m_2})'_s, p^k) \leq v_p r^{m_2}$$

solutions in a . Hence the number of solutions of

$$vr_1^{m_1}(r^{m_2})'_s \equiv b \pmod{p^k}$$

in $m = m_1e_1 + m_2 \in K(p^k)$ does not exceed

$$e_1\alpha_8v_p(1 + r + \dots + r^{e_1-1}) = \alpha_{10}(r, s)v_p .$$

But this implies that the number of solutions of

$$vr_1^{m_1}(r^{m_2})'_s \equiv b \pmod{s^k}$$

in $m = m_1e_1 + m_2 \in K(s^k)$ is not greater than

$$\alpha_{10}(r, s)v_p\left(\frac{s}{p}\right)^k \leq \alpha_{10}(r, s)\left(\frac{s}{2}\right)^k v_p .$$

B. The equation

$$ur^m + vr^m \equiv b \pmod{p^k}$$

has according to the corollary of Lemma 4 at most

$$\alpha_8(r, p)v_p$$

solutions in $m \in K(p^k)$. The result follows as before.

The following conjecture seems related to our results: *Assume $r \not\sim s$. Then for any ε and k almost all the numbers r, r^2, \dots are (ε, k) -normal to the base s in the sense of Besicovitch [1]; that is, the number of $n \leq N$ for which r^n is not (ε, k) -normal is $o(N)$ as $N \rightarrow \infty$ for fixed ε and k .*

6. Bounds for exponential sums.

LEMMA 6. A. *Let r, s, v be as in Lemma 5A. Then*

$$\sum_{m \in K(s^k)} \Pi(s, t; vr^m) \leq \alpha_{13}v_p s^{(1-\alpha_{13})k}$$

B. *Let r, s, u, v, n be as in Lemma 5B. Then*

$$\sum_{m \in K(s^k)} \Pi(s, t; ur^m + vr^m) \leq \alpha_{14}v_p s^{(1-\alpha_{15})k} .$$

Proof. A. Write $v(r^m)'_s = c_\nu \dots c_k \dots c_1$ in the scale of s . Lemma 5A implies that any digit combination $c_k c_{k-1} \dots c_1$ will occur at most $\alpha_{10}(r, s)(s/2)^k v_p$ times. According to Lemma 3, there are for large k not more than $2^{(3/4)k}$ digit-combinations $c_k \dots c_1$ with less than $\alpha_3 k$ indices i satisfying (9). Thus of all the numbers $v(r^m)'_s, m \in K(s^k)$, and hence of all the numbers vr^m there will be at most

$$\alpha_{10}(r, s)(s/2)^k v_p 2^{(3/4)k} = \alpha_{10}(r, s)v_p (s/2^{1/4})^k = \alpha_{10}(r, s)v_p s^{(1-\alpha_{16})k}$$

having less than $\alpha_3 k$ digits c_i in their expansion in the scale of s satisfying (9). Thus Lemma 2 yields

$$\Pi(s, t; vr^m) \leq \alpha_2^{k\alpha_3}$$

for all but at most

$$\alpha_{10}(r, s)v_p s^{(1-\alpha_{16})k}$$

numbers $m \in K(s^k)$. This gives

$$\sum_{m \in K(s^k)} \Pi(s, t; vr^m) \leq s^k \alpha_2^{k\alpha_3} + \alpha_{10} v_p s^{(1-\alpha_{16})k} \leq \alpha_{12} v_p s^{(1-\alpha_{13})k} .$$

B is proved similarly, using Lemma 5B.

LEMMA 7. A. Assume $r \not\sim s, v \neq 0$. Then

$$(11) \quad \sum_{N_1 < n \leq N_2} \Pi(s, t; vr^m) \leq \alpha_{17}(N_2 - N_1)^{1-\alpha_{18}} v_p .$$

B. Assume $r \not\sim s, u \neq 0, v \neq 0$. Then

$$(12) \quad \sum_{N_1 < n, m \leq N_2} \Pi(s, t; ur^n + vr^m) \leq \alpha_{19}(N_2 - N_1)^{2-\alpha_{20}} \max(u_p, v_p) .$$

Proof. A. There exists a k having $s^{2k} \leq N_2 - N_1 < s^{2(k+1)}$, hence there exists a w satisfying $s^k w \leq N_2 - N_1 < s^k(w + 1)$, where $s^k \leq w < s^{k+2}$. Thus if m runs from N_1 to N_2 , then m runs through w systems $K(s^k)$ of residue classes modulo s^k and at most s^k other numbers. Hence by Lemma 6A

$$\sum_{N_1 < m \leq N_2} \Pi(s, t; vr^m) \leq w \alpha_{12} v_p s^{(1-\alpha_{13})k} + s^k \leq \alpha_{17}(N_2 - N_1)^{1-\alpha_{18}} v_p .$$

B. If $p \nmid r$, then we proceed as in part A. We first take the sum over m and use Lemma 6B.

If $p \mid r$, then our argument is as follows. Consider, for example, the part of the sum with $n \leq m$. Changing the notation in n, m , we see that this part of the sum (12) equals

$$\sum_{n=0}^{N_2-N_1-1} \sum_{m=N_1+1}^{N_2-n} \Pi(s, t; (ur^n + v)r^m) .$$

Except for possibly one exceptional n we have $(ur^n)_p \neq v_p$ and therefore $(ur^n + v)_p \leq v_p \leq \max(u_p, v_p)$. If n is not exceptional, then the already proved Lemma 7A can be applied to the inner sum and we obtain the bound

$$\alpha_{17}(N_2 - N_1 - n)^{1-\alpha_{18}} \max(u_p, v_p) .$$

Taking the sum over n we obtain (12).

7. A fundamental lemma. Generalizing $M_N(\xi, r, I)$ we write ${}_{N_1}M_{N_2}(\xi, r, I)$ for the number of indices i in $N_1 < i \leq N_2$ such that $\{r^i\xi\}$ lies in I . We put

$${}_{N_1}R_{N_2}(\xi, r, I) = {}_{N_1}M_{N_2}(\xi, r, I) - (N_2 - N_1)|I|.$$

Fundamental lemma. Assume $r \not\sim s$. Then

$$\int_0^1 {}_{N_1}R_{N_2}^2(T_{s,t}\xi, r, I) d\xi \leq \alpha_{21}(N_2 - N_1)^{2-\alpha_{22}}.$$

Proof. It is enough to prove this for intervals of the type $I = [0, \gamma)$. Then

$${}_{N_1}M_{N_2}(\xi, r, I) = \sum_{N_1 < n \leq N_2} \{r^n\xi, \gamma\}$$

and

$$(13) \quad \int_0^1 {}_{N_1}M_{N_2}(T_{s,t}\xi, r, I) d\xi = \sum_{N_1 < n \leq N_2} \mu(\gamma, r^n)$$

$$(14) \quad \int_0^1 {}_{N_1}M_{N_2}^2(T_{s,t}\xi, r, I) d\xi = \sum_{N_1 < n, m \leq N_2} \mu(\gamma, r^n, r^m).$$

Now we combine (8) and Lemma 7. We obtain, together with (6),

$$\begin{aligned} \sum_{N_1 < n, m \leq N_2} \mu(\gamma, r^n, r^m) &\leq (\gamma + \eta)^2(N_2 - N_1)^2 \\ &+ 2(\gamma + \eta) \sum_{v \neq 0} \frac{v_p}{\eta v^2} \alpha_{17}(N_2 - N_1)^{2-\alpha_{18}} \\ &+ \sum_{u \neq 0} \sum_{v \neq 0} \frac{\max(u_p, v_p)}{\eta u^2 \eta v^2} \alpha_{19}(N_2 - N_1)^{2-\alpha_{20}}. \end{aligned}$$

Since the sums

$$\sum_{v \neq 0} \frac{v_p}{v^2}, \quad \sum_{u \neq 0} \sum_{v \neq 0} \frac{\max(u_p, v_p)}{u^2 v^2}$$

are convergent, and since η was arbitrary, we have

$$\sum_{N_1 < n, m \leq N_2} \mu(\gamma, r^n, r^m) - (N_2 - N_1)^2 \gamma^2 \leq \alpha_{23}(N_2 - N_1)^{2-\alpha_{24}}.$$

In the same fashion we can prove

$$\begin{aligned} \left| \sum_{N_1 < n, m \leq N_2} \mu(\gamma, r^n, r^m) - (N_2 - N_1)^2 \gamma^2 \right| &\leq \alpha_{23}(N_2 - N_1)^{1-\alpha_{24}} \\ \left| \sum_{N_1 < n \leq N_2} \mu(\gamma, r^n) - (N_2 - N_1) \gamma \right| &\leq \alpha_{25}(N_2 - N_1)^{1-\alpha_{26}}. \end{aligned}$$

These two inequalities, together with (13) and (14), give the Fundamental Lemma.

8. Proof of the theorems. Once the Fundamental Lemma is shown, we can prove Theorem 2 by the standard method developed in [2].

By J_B , $B > 0$, we denote the set of intervals $[\beta, \gamma)$, $0 \leq \beta < \gamma < 1$ of the type $\beta = a2^{-b}$, $\gamma = (a + 1)2^{-b}$, where $0 \leq b \leq \alpha_{22}B/2$. By P_B we denote the set of all pairs of integers N_1, N_2 having $0 \leq N_1 < N_2 \leq 2^B$ of the type $N_1 = a2^b$, $N_2 = (a + 1)2^b$ for integers a and $b \geq 0$.

LEMMA 8. *Assume $r \neq s$. Then*

$$\sum_{(N_1, N_2) \in P_B} \sum_{I \in J_B} \int_0^1 R_{N_1}^2(T_{s, t\xi}, r, I) d\xi \leq \alpha_{27} 2^{2B(1-\alpha_{28})} .$$

Proof. Because of the Fundamental Lemma the left hand side is not greater than

$$\alpha_{21} 2^{\alpha_{22}B/2+1} \Sigma ,$$

where $2^{\alpha_{22}B/2+1}$ is an upper bound for the number of intervals in J_B and

$$(15) \quad \Sigma = \sum_{(N_1, N_2) \in P_B} (N_2 - N_1)^{2-\alpha_{22}} .$$

In (15) each value of $N_2 - N_1 = 2^b$ occurs 2^{B-b} times, so that

$$\Sigma = \sum_{b=0}^B 2^{B-b+b(2-\alpha_{22})} \leq \alpha_{29} 2^{2B(1-\alpha_{22}/2)} .$$

Hence Lemma 8 is true with $\alpha_{28} = \alpha_{22}/4$.

LEMMA 9. *For large B there exists a set E_B of measure not greater than $2^{-\alpha_{30}B}$ such that*

$$(16) \quad R_N(T_{s, t\xi}, r, I) \leq 2^{B(1-\alpha_{31})}$$

for all I , $N \leq 2^B$ and all ξ in $[0, 1)$ but not in E_B .

Proof. We define E_B to be the set consisting of all ξ in $[0, 1)$ for which it is not true that

$$(17) \quad \sum_{(N_1, N_2) \in P_B} \sum_{I \in J_B} N_1 R_{N_2}^2(T_{s, t\xi}, r, I) \leq 2^{2B(1-\alpha_{28}/2)} .$$

Lemma 8 assures that the measure of E_B does not exceed

$$\alpha_{27} 2^{-2B\alpha_{28}/2} < 2^{-\alpha_{30}B}$$

for large B . We have to show that (16) is a consequence of (17).

We first assume I to be of the type $I = [0, \gamma)$, $\gamma = a2^{-b}$, where $0 \leq b \leq \alpha_{22}B/2$. Then the interval $[0, \gamma)$, is the sum of at most $b < B$ intervals I , $I \in J_B$, as may be seen by expressing a in the binary scale.

Expressing N in the binary scale we see that the interval $[0, N)$ can be expressed as a union of at most B intervals $[N_1, N_2)$, where the pair $N_1, N_2 \in P_B$. Hence we can write $R_N(T_{s,t}\xi, r, I)$ as a sum of $_{N_1}R_{N_2}(T_{s,t}\xi, r, I)$ over at most B^2 sets N_1, N_2, I , where $N_1, N_2 \in P_B, I \in J_B$:

$$R_N(T_{s,t}\xi, r, I) = \Sigma_{N_1} R_{N_2}(T_{s,t}\xi, r, I).$$

Hence by (17) and Cauchy's inequality,

$$R_N^2(T_{s,t}\xi, r, I) \leq B^2 2^{2B(1-\alpha_{28}/2)} < 2^{2B(1-\alpha_{32})}$$

for large B .

Next, let $I = [0, \gamma)$ be of the type $a2^{-b} \leq \gamma \leq (a+1)2^{-b}$, where $\alpha_{22}B/4 < b \leq \alpha_{22}B/2$. Then

$$\begin{aligned} |R_N(T_{s,t}\xi, r, [0, \gamma))| &= |M_N(T_{s,t}\xi, r, [0, \gamma)) - \gamma N| \\ &\leq |R_N(T_{s,t}\xi, r, [0, (a+1)2^{-b}))| + |R_N(T_{s,t}\xi, r, [0, a2^{-b}))| + 2^{-b}N \\ &\leq 2 \cdot 2^{B(1-\alpha_{32})} + 2^{(1-\alpha_{22}/4)B} < 2^{B(1-\alpha_{33})}. \end{aligned}$$

The Lemma now follows from

$$|R_N(\cdot, \cdot, [\beta, \gamma))| \leq |R_N(\cdot, \cdot, [0, \beta))| + |R_N(\cdot, \cdot, [0, \gamma))|.$$

Proof of Theorem 2. Since $\Sigma 2^{-\alpha_{30}B}$ is convergent, there exists for almost all ξ a $B_0 = B_0(\xi)$ such that $\xi \notin E_B$ for $B \geq B_0$. If $N \geq 2^{B_0}$, then we can find a $B \geq B_0$ satisfying $2^{B-1} < N \leq 2^B$ and Lemma 9 yields

$$R_N(T_{s,t}\xi, r, I) < 2^{B(1-\alpha_{31})} < 2N^{1-\alpha_{31}} < N^{1-\alpha_1}$$

for large enough N .

REFERENCES

1. A. S. Besicovitch, *The asymptotic distribution of numerals in the decimal representation of squares of natural numbers*, Math. Zeit. **39** (1934), 146-156.
2. J. W. S. Cassels, *Some metrical theorems in Diophantine approximation III*, Proc. Camb. Phil. Soc. **46** (1951), 219-225.
3. ———, *On a paper of Niven and Zuckerman*, Pacific J. Math, **2** (1952), 555-557.
4. H. Hasse, *Vorlesungen über Zahlentheorie*, Springer Grundlehren LIX (1950).
5. J. F. Koksma, *Diophantische Approximationen*, Ergebnisse der Math. u. ihrer Grenzgebiete 4, **4** (1936).
6. I. Niven, *Irrational numbers*, The Carus Math. Monographs **11** (1956).
7. I. Niven and H. S. Zuckerman, *On the definition of normal numbers*, Pacific J. Math. **1** (1951), 103-110.
8. D. D. Wall, *Normal numbers*, Ph. D. thesis (1949), University of California, Berkeley.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DAVID GILBARG
Stanford University
Stanford, California

F. H. BROWNELL
University of Washington
Seattle 5, Washington

A. L. WHITEMAN
University of Southern California
Los Angeles 7, California

L. J. PAIGE
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH
T. M. CHERRY
D. DERRY

E. HEWITT
A. HORN
L. NACHBIN

M. OHTSUKA
H. L. ROYDEN
M. M. SCHIFFER

E. SPANIER
E. G. STRAUS
F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
HUGHES AIRCRAFT COMPANY
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 2120 Oxford Street, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 10, No. 2

October, 1960

Maynard G. Arsove, <i>The Paley-Wiener theorem in metric linear spaces</i>	365
Robert (Yisrael) John Aumann, <i>Acceptable points in games of perfect information</i>	381
A. V. Balakrishnan, <i>Fractional powers of closed operators and the semigroups generated by them</i>	419
Dallas O. Banks, <i>Bounds for the eigenvalues of some vibrating systems</i>	439
Billy Joe Boyer, <i>On the summability of derived Fourier series</i>	475
Robert Breusch, <i>An elementary proof of the prime number theorem with remainder term</i>	487
Edward David Callender, Jr., <i>Hölder continuity of n-dimensional quasi-conformal mappings</i>	499
L. Carlitz, <i>Note on Alder's polynomials</i>	517
P. H. Doyle, III, <i>Unions of cell pairs in E^3</i>	521
James Eells, Jr., <i>A class of smooth bundles over a manifold</i>	525
Shaul Foguel, <i>Computations of the multiplicity function</i>	539
James G. Glimm and Richard Vincent Kadison, <i>Unitary operators in C^*-algebras</i>	547
Hugh Gordon, <i>Measure defined by abstract L_p spaces</i>	557
Robert Clarke James, <i>Separable conjugate spaces</i>	563
William Elliott Jenner, <i>On non-associative algebras associated with bilinear forms</i>	573
Harold H. Johnson, <i>Terminating prolongation procedures</i>	577
John W. Milnor and Edwin Spanier, <i>Two remarks on fiber homotopy type</i>	585
Donald Alan Norton, <i>A note on associativity</i>	591
Ronald John Nunke, <i>On the extensions of a torsion module</i>	597
Joseph J. Rotman, <i>Mixed modules over valuations rings</i>	607
A. Sade, <i>Théorie des systèmes démosiens de groupoïdes</i>	625
Wolfgang M. Schmidt, <i>On normal numbers</i>	661
Berthold Schweizer, Abe Sklar and Edward Oakley Thorp, <i>The metrization of statistical metric spaces</i>	673
John P. Shanahan, <i>On uniqueness questions for hyperbolic differential equations</i>	677
A. H. Stone, <i>Sequences of coverings</i>	689
Edward Oakley Thorp, <i>Projections onto the subspace of compact operators</i>	693
L. Bruce Treybig, <i>Concerning certain locally peripherally separable spaces</i>	697
Milo Wesley Weaver, <i>On the commutativity of a correspondence and a permutation</i>	705
David Van Vranken Wend, <i>On the zeros of solutions of some linear complex differential equations</i>	713
Fred Boyer Wright, Jr., <i>Polarity and duality</i>	723