Pacific Journal of Mathematics

CONCERNING CERTAIN LOCALLY PERIPHERALLY SEPARABLE SPACES

L. BRUCE TREYBIG

Vol. 10, No. 2

October 1960

CONCERNING CERTAIN LOCALLY PERIPHERALLY SEPARABLE SPACES

L. B. TREYBIG

In 1954, F. Burton Jones raised the question [2] "Is every connected, locally peripherally separable [3], metric space separable?" In this paper it will be shown that there exists a connected, semi-locally-connected, space Σ satisfying R. L. Moore's axioms 0 and C^1 , in which every region has a separable boundary, every pair of points is a subset of some separable continuum², and the set of all points at which Σ is not locally separable is separable. It will also be shown that every compactly connected, locally peripherally separable, metric space is completely separable.

Part 1

Let S' denote the set of all points of the Euclidean plane E. A square disk in E will be said to be horizontal if it has two horizonta sides. A point set in E will be called an *H*-disk only if that set is a horizontal square disk. By the *width* of a square disk will be meant the length of one of its sides.

Let K denote a definite H-disk of width d. Let $R_0(K)$ denote the H-disk of width d/4 whose center is on the vertical line that contains the center of K, and whose upper side lies at a distance of d/16 below the upper side of K. Let $R_{00}(K)$ and $R_{01}(K)$ denote the H-disks of width d/8 whose upper sides are at a distance of d/32 above the lower side of $R_0(K)$ and whose centers are on the vertical lines containing the left and right sides, respectively, of $R_0(K)$.

In general, for each positive integer n let $U_n(K)$ denote a collection of 2^n mutually exclusive congruent *H*-disks such that

(1) $R_{01}(K)$ and $R_{00}(K)$ are the elements of $U_1(K)$,

(2) if n is a positive integer and y is an element of $U_n(K)$, and x and z are H-disks of width $d/4(2)^{n+1}$ whose centers lie on the same vertical lines as the left and right sides of y, respectively, and whose upper sides lie at a distance of $d/32(2)^n$ above the lower side of y, then x and z are elements of $U_{n+1}(K)$.

If n is a positive integer and $R_{x_1x_2\cdots x_n}(K)$ is an elements of $U_n(K)$, then let the elements x and y of $U_{n+1}(K)$ whose centers lie on the same

Presented to the American Mathematical Society, June 15, 1957; received by the editors September 4, 1958. This paper is part of a dissertation submitted to the Graduate School of the University of Texas in partial fulfillment of the requirements for the Ph. D. degree.

 $^{^{1}}$ The proof that every space which satisfies axioms 0 and C is metric is due to R. L. Moore.

² A continuum is a connected, closed set.

vertical lines as the left and right sides of $R_{x_1x_2\cdots x_n}(K)$, respectively, be denoted by $R_{x_1x_2\cdots x_n}(K)$ and $R_{x_1x_2\cdots x_n}(K)$, respectively. Let C(K) be a collection to which x belongs if and only if x is $R_0(K)$ or in one of the collections $U_1(K)$, $U_2(K)$, \cdots .

Let L(K) denote the *H*-disk of width d/8 whose center is on the same vertical line as the center of *K*, and whose lower side is at a distance of 3d/16 above the lower side of *K*. Let $P_i(K)$ and $P_r(K)$ denote the left and right-hand end points, respectively, of the lower side of L(K). Let M(K) denote the point set such that a point *P* belongs to it if and only if *P* is a point of the interval $P_i(K)P_r(K)$ such that there is no nonnegative integer *p* and positive integer *q* such that $PP_i(K)/P_i(K)P_r(K) = p/2^q$. Let I(K) denote the collection to which *x* belongs if and only if *x* is a vertical interval containing a point of M(K), and with both end points on the boundary of L(K). Let an interval *i* of I(K) be denoted by $i_x(I(K))$ if and only if it is true that if *P* is the lowest point of *i*, then $P_i(K)P_r(K) = x$.

Let R denote some definite H-disk. Let $R_0(R)$ be denoted by Q_0 ; let $R_{00}(R)$ and $R_{01}(R)$ be denoted by Q_{00} and Q_{01} , respectively. Let $R_{000}(R)$, $R_{001}(R)$, $R_{010}(R)$, and $R_{011}(R)$ be denoted by Q_{000} , Q_{001} , Q_{010} , and Q_{011} , respectively, and so forth. Let C(R) be denoted by C_1 and let I(R) be denoted by I_0 .

Let C_2 denote the collection to which x belongs if and only if x is an element of C(y), for some element y of C_1 distinct from Q_0 . Let $R_0(Q_{00})$ be denoted by $Q_{00,0}$; let $R_{01}(Q_{00})$ be denote by $Q_{00,01}$. In general, let $R_x(Q_y)$ be denoted by $Q_{y,x}$. Also, if Q_x is in C_1 and $x \neq 0$, let $I(Q_x)$ be denoted by I_x .

In general, let C_{n+1} denote the collection to which x belongs if and only if x is an element of C(y), for some elements y of C_n , which, in case x_n is 0, is distinct from $Q_{x_1,x_2,...,x_n}$. Let the element $R_{x_{n+1}}[R_{x_n}[R_{x_{n-1}}[\cdots [R_{x_1}(R)]\cdots]]]$ of C_{n+1} be denoted by $Q_{x_1,x_2,...,x_{n+1}}$. Also if w is the element $Q_{x_1,x_2,...,x_n}$ of C_n and $x_n \neq 0$, then let I(w) be denoted by $I_{x_1,x_2,...,x_n}$. For each n let I_n be the collection to which x belongs if and only if there is an element $Q_{x_1,x_2,...,x_n}$ of C_n such that $x_n \neq 0$ and xis in $I(Q_{x_1,x_2,...,x_n})$.

Let W denote the point set to which a point P belongs if and only if P belongs to C_n^{*3} for each positive integer n. For each positive integer n let B_n denote the collection of all the boundaries of the elements of C_n . The boundary of Q_{x_1,x_2,\ldots,x_n} will be denoted by J_{x_1,x_2,\ldots,x_n} .

Let S denote $[I_0^* + I_1^* + \cdots] + [B_1^* + B_2^* + \cdots] + W$.

Let C' be a collection to which w belongs if and only if w is R or I(w) is a subset of S and there is a positive integer n such that w is in C_n .

³ C_n^* Means the sum of all the point sets of the collection C_n .

For each positive integer n let H_n denote a collection to which x belongs if and only if x is the common part of S and the interior of some square of $[B_n + B_{n+1} + \cdots]$. For each element $Q_{x_1, x_2, \cdots, x_n}$ of C_n , let the set of all points of S in the interior of $J_{x_1, x_2, \cdots, x_n}$ be denoted by $r_{x_1, x_2, \cdots, x_n}$.

For each positive integer n let K_n denote a collection to which x belongs if and only if, either (1) x is a segment of an arc lying on some square J of $(B_1 + B_2 + \cdots)$, having length less than $1/4^n$ times the perimeter of J, and intersecting no square of the collection $(B_1 + B_2 + \cdots)$ except J, or (2) x is the sum of two straight line segments p and q intersecting at their midpoints and lying on different squares J_p and J_q of $(B_1 + B_2 + \cdots)$, such that p and q each have length less than $1/4^n$ times the perimeters of J_p and J_q , respectively, and such that neither p nor q intersects three squares of $(B_1 + B_2 + \cdots)$.

Suppose x is a positive number such that $i_x[I_{j_1,j_2,...,j_n}]$ is an interval of $I_{j_1,j_2,...,j_n}$. For each positive integer n there exists a unique pair (k_n, x_n) such that k_n is a non-negative integer, x_n is a positive number less than one, and $x = (k_n + x_n)/2^n$. By $i_n[i_x(I_{j_1,j_2,...,j_n})]$ will be meant the vertical interval $i_{x_n}(I(y))$, where y is the H-disk of $U_n[Q_{j_1,j_2,...,j_n}]$ with only k_n disks of $U_n(Q_{j_1,j_2,...,j_n})$ to the left of it.

Suppose, for some y in C', P is the highest point of the interval $i_x(I(y))$. By $R_n(P)$ will be meant the sum of all the sects z such that either

(1) for some positive integer d greater than or equal to n, z is the subset of $i_a[i_x(I(y))]$ with length $1/2^n$ times the length of $i_a[i_x(I(y))]$ that contains the lowest point of $i_a[i_x(I(y))]$, or

(2) z is the subset of $i_x[I(y)]$ with length $1/2^n$ times the length of $i_x(I(y))$ that contains the highest point of $i_x(I(y))$.

For each positive integer n let L_n denote a collection such that x belongs to it if and only if there exists a positive integer d greater than or equal to n, an element y of C', and an interval of the collection I(y) such that if P denotes the highest point of that interval, then $x = R_a(P)$.

For each positive integer n let N_n denote a collection to which x belongs if and only if either

(1) for some element y of C' there exists an interval i of the collection I(y) such that x is a segment of i of length less than $1/2^n$ times the length of i, or

(2) for some element y of C' there exists an element i of I(y) such that x is a sect lying in i, containing the lowest point of i and of length less than $1/2^n$ times the length of i.

For each positive integer n let G_n denote a collection to which a belongs if and only if it lies in $H_n + K_n + L_n + N_n$. S is the set of

all points of Σ . A subset r of S is a region in Σ if and only if r belongs to G_1^4 .

R. L. Moore's axioms 0 and C are as follows:

Axiom 0. Every region is a point set.

Axiom C. There exists a sequence G_1, G_2, \cdots such that

(1) for each positive integer n, G_n is a collection such that each element of G_n is of region and G_n covers S,

(2) for each n, G_{n+1} is a subcollection of G_n ,

(3) if A is a point, B is a point and R is a region containing A, then there exists a positive integer n such that if x is a region of G_n containing A and y is a region of G_n intersecting x, then

(a) y is a subset of R and

(b) if B is not A, y does not contain B,

(4) if M_1, M_2, \cdots is a sequence of closed point sets such that for each *n* there exists a region g_n of G_n such that M_n is a subset of \overline{g}_n and for each $n M_n$ contains M_{n+1} , then there is a point common to all the point sets of this sequence.

It is obvious that in the space Σ each region has a countable, and therefore separable, boundary, and that the sequence G_1, G_2, \cdots defined for the space Σ satisfies conditions (1) and (2) of axiom C. It will be shown that it also satisfies conditions (3) and (4) of this axiom.

Suppose that P is a point of W, that $r = r_{x_1,x_2,...,x_n}$ is a region of H_n containing P, and that Q is a point of r distinct from P. If q is a region containing a point of W, then q must belong to H_1 . Since each element of C_{n+1} which contains P has a side of length less than or equal 1/4 times the length of a side of $Q_{x_1,x_2,...,x_n}$, and each element of C_{n+2} which contains P has a side of length less than or equal $1/4^2$ times the length of side of $Q_{x_1,x_2,...,x_n}$, and so forth; it is obvious that there is a d > n such that if q is a region of H_a which contains P, then \bar{q} does not intersect Q and is a subset of r. Suppose that x and y are two intersecting regions of G_{n+1} such that x contains P. x belongs to H_{n+1} and is therefore a subset of r. Every region of G_{n+1} which intersects x is a subset of r, so clearly, y is a subset of r.

Now suppose that P is a point of $J_{x_1,x_2,...,x_n}$ of B_n and r is a region containing P, and Q is a point of r distinct from P. There exists a circle J in E with center at P such that every point of S in the interior of J belongs to r, but Q is not in the interior of J. There exists a positive integer d such that $1/4^a$ times the perimeter of any square of $(B_1 + B_2 + \cdots)$ to which P belongs is less than the radius of J, and such that no region of H_a contains P. If R^1 is a region of G_{a+1} containing P, then \overline{R}^1 does not contain Q and is a subset of r. If n > d+2

⁴ The collection G_1 of regions is a basis for the space Σ .

and x and y are two intersecting regions of G_n such that x contains P, then x + y is a subset of r.

Now suppose that P is a point of $i_x(I(y))$, for y in C', and that r is a region containing P and that Q is a point of r distinct from P.

Case 1. Suppose P is not the highest point of $i_x(I(y))$. There exists a segment t containing P, or a sect in case P is the lowest point of $i_x(I(y))$, such that t is a subset of r and does not contain Q nor the highest point of $i_x(I(y))$. There exists a positive number ε such that every point of $i_x(I(y))$ which is at a distance from P of less than ε lies in t. There exists a positive integer d such that

(1) no region of L_a intersects t and no region of H_a intersects $i_x(I(y))$, and

(2) $1/2^{a}$ times the length of $i_{x}(I(y))$ is less than ε . Therefore, if k is a region of G_{a+1} containing P, then \overline{k} is a subset of r and does not contain Q. Also, if x and y are two intersecting regions of G_{a+2} such that x contains P, then x + y is a subset of r.

Case 2. Suppose P is the highest point of $i_x(I(y))$. Whether Q belongs to $i_x(I(y))$ or there is a positive integer p such that Q belongs to $i_p[i_x(I(y))]$ or r is in H_1 and Q does not belong to $i_x(I(y)) + i_1[i_x(I))] + i_2[i_x(I(y))] + \cdots$, there is a positive integer d such that

(1) $R_d(P)$ does not contain Q and is a subset of r, and

(2) no region of H_a contains P. If k is a region of G_{a+1} containing P, then \overline{k} is a subset of r and does not contain Q. Also, if x and y are two intersecting regions of G_{a+3} such that x contains P, then x + y is a subset of r.

Therefore G_1, G_2, \cdots satisfies the third part of axiom C.

Suppose that M_1, M_2, \cdots is a sequence of closed point sets such that

(1) for each $n M_n$ contains M_{n+1} , and

(2) for each n there is a region g_n of G_n such that M_n is a subset of \overline{g}_n .

In case, for each n, g_n is in H_n , then by definition of W, there is a point common to M_1, M_2, \cdots because some point of W can be easily shown to be a limit point or point of M_n for each n.

In case there is a positive integer j such that g_j belongs to K_j , then for n > j, g_n belongs to K_n . But M_j , M_{j+1} , \cdots is a sequence of closed and compact point sets such that for $n \ge j$ M_n contains M_{n+1} . So there is a point common to M_j , M_{j+1} , \cdots and thus common to M_1 , M_2 , \cdots .

In case there is a positive integer j such that g_j belongs to N_j , then for n > j, g_n belongs to N_n . So, for the same reason as in the L. B. TREYBIG

previous case, there is a point common to M_1, M_2, \cdots .

The only case not considered is the one where there is a positive integer j_1 such that, for $n \ge j_1$, g_n belongs to L_n . In this case g_{j_1} must be $R_{x_1}(P)$ for some point P and positive integer x_1 . There is a positive integer $j_2 > j_1$ such that $g_{j_2} = R_{x_2}(P)$, where $x_2 > x_1$. There is a positive integer $j_3 > j_2$ such that $g_{j_3} = R_{x_3}(P)$, for $x_3 > x_2$, and so forth. P is common to the sets $R_{x_1}(P)$, $R_{x_2}(P)$, \cdots . But if P does not belong to each of the sets M_{j_1}, M_{j_2}, \cdots then there is a positive integer d such that $\overline{R}_{x_d}(P)$ contains no point of M_{x_j} for any j. But $R_{x_d}(P)$ contains $M_{j_{d+1}}$. So P is common to the sets M_{j_1}, M_{j_2}, \cdots and thus common to M_{j_1}, M_{j_2}, \cdots .

Thus, Σ satisfies the fourth part of axiom C.

In order to show that Σ is connected, an indirect argument will be used. Suppose that S is the sum of two mutually separated sets H and K. Since $W + (B_1^* + B_2^* + \cdots)$ is connected, let H' be the one of the sets H and K that contains this set and let K' be the other. There exists an element y of C' such that for some $x i_x[I(y)]$ is a subset of K'. But there exists a positive integer d_1 such that for $n \ge d_1$, $i_n[i_x(I(y))]$, belongs to K'. There exists a positive integer d_2 such that for $n \ge d_2 i_n[i_{a_1}(i_x(I(y)))]$ belongs to K'. So, obviously, there is a positive integer sequence, d_1, d_2, \cdots such that if j is a positive integer and $n \ge d_j$, then $i_n(i_{a_{j-1}}(i_{a_{j-2}}(\cdots i_{a_1}(i_x(I(y)))\cdots)))$ belongs to K'. But from this fact it is easily seen that some point of W is a limit point of K'. So Σ is connected.

It has been shown that in any space satisfying axioms 0 and C(1) if M is a separable point set, M is completely separable, and (2) if M is separable, any subset of M is separable.

In order to show that any two points of S lie in a separable continuum, suppose first that P and Q are two points of S. Obviously, $(B_1^* + B_2^* + \cdots)$ is separable and connected, and therefore $W + (B_1^* + B_2^* + \cdots)$ is a separable continuum. In case P and Q both lie in $W + (B_1^* + B_2^* + \cdots)$, this continuum has the desired properties. In case P does not belong to this set, P belongs to $i_x[I(y)]$ for some y in C'. Let M_P be the set to which point R belongs if and only if, either

(1) there is a finite positive integer sequence x_1, x_2, \dots, x_n such that R belongs to $i_{x_1}[i_{x_2}[\dots i_{x_n}[i_x(I(y))]\dots]]$, or

(2) there is a positive integer q such that R belongs to $i_q[i_x(I(y))]$, or

(3) R belongs to $i_x[I(y)]$. $M_P + (B_1^* + B_2^* + \cdots) + W$ is a separable continuum. If Q does not belongs to this set, let M_Q be a set related to Q like M_P was related to P. The continuum $M_P + M_Q + (B_1^* + B_2^* + \cdots) + W$ is separable.

The statement that Σ is locally separable at the point P means that there is a region R containing P such that R is separable. Alexandroff [1] has shown that if β is a connected, locally completely separable,

space satisfying axioms 0 and C, then β is completely separable. It is interesting to note that Σ is locally separable, and therefore locally completely separable, at each point except those of a separable set, and yet, Σ is not separable.

 Σ is obviously locally separable at all points not belonging to W. Since every region that contains a point of W contains uncountably many mutually exclusive domains, Σ is not locally separable at any point of W. Furthermore $(B_1^* + B_2^* + \cdots)$ is separable, and so $(\overline{B_1^* + B_2^* + \cdots})$ is separable, and thus, since W is a subset of the latter, W is separable.

 Σ is said to be *semi-locally-connected* [5] at point *P* if and only if it is true that if *R* is a region containing *P*, *R* contains a region *R'* containing *P* such that S - R does not intersect infinitely many components of S - R'. Σ is said to be semi-locally-connected if and only if Σ is semi-locally-connected at each point.

The space Σ is obviously semi-locally-connected because S minus any region has only a finite number of components.

Part 2

Suppose that Σ is a space satisfying the conditions specified on the first page of this paper.

For each positive integer j let G_j denote the collection of all open sets which have diameter less that j^{-1} .

Let P denote some definite point, and suppose n is a positive integer such that no countable subcollection of G_n covers S. Let R_n be some region of G_n which contains P, let $H_1 = \{R_n\}$, and let K_1 be the boundary of R_n .

For each point Q of S let $\Delta(Q)$ be the least integer j > n such that some region R(Q) of G_n contains every region of G_j that intersects a region of G_j that contains Q.

It has been shown that in a space satisfying these axioms if L is a separable point set and G is a collection of open sets covering L, then some countable subcollection of G covers L. Therefore, there is a countable point set T_1 dense in K_1 such that the collection H_2 of all R(Q)'s, for Q's in T_1 , covers K_1 . Let K_2 be the sum of the boundaries of all the sets in $H_1 + H_2$. There is a countable point set T_2 dense in K_2 such that the collection H_3 of all R(Q)'s for Q's in T_2 , covers K_2 . Let K_3 be the sum of the boundaries of the sets in $H_1 + H_2 + H_3$, and so forth.

There is a point B not in the closure of $H = (H_1 + H_2 + \cdots)^*$. Let M be a compact continuum containing P and B.

Case 1. Suppose some point A of $M - M \cdot H$ is a limit point of $K = K_1 + K_2 + \cdots$.

Let R'_1 be a region of G_n containing A, let Q_1 be a point of $T = T_1 + T_2 + \cdots$ in R'_1 , and let x_1 be the largest integer i such that R'_1 belongs to G_i . Let R'_2 be a region of $C_{x_{l+1}}$ containing A such that $\bar{R'_2}$ lies in $R'_1 - Q_1$. Let Q_2 be a point of T in R'_2 and let x_2 be the largest integer i such that R'_2 is in G_i . Obtain R'_3, Q_3 , and x_3 similarly, and so forth. $n \leq x_1 < x_2 < x_3 < \cdots$. For each $i, \Delta(Q_i) > x_i$. Otherwise, for some $i, R(Q_i)$ would contain R'_i , and thus A. However, there is a positive integer t > n such that if x, y, and z are regions of G_i such that $x \cdot y$ and $y \cdot z$ exist and x contains A, then R_n contains x + y + z. For some $s > t, \Delta(Q_s) > t$. But R_n contains every region of G_i that intersects a region of G_i that contains G_s . So $\Delta(Q_s) \leq t$, which is a contradiction.

Case 2. Suppose no point of $M - M \cdot H$ is a limit point of K. For each point Q of $M - M \cdot H$ let g_q be a region containing Q such that g_q contains no point of K + P. Some finite subcollection C of the g_q 's covers this set of limit points. Let $D = H - H \cdot \overline{C}^*$. Let C_1 be the component of $M - M \cdot \overline{D}$ which contains B. Some point z of $M \cdot \overline{D}$ is a limit point of C_1 . But z lies in a region r of H, and therefore C_1 would intersect the boundary of r, and thus contain a limit point of K. This yields a contradiction.

Since, for each n, some countable subcollection of G_n covers S, Σ is completely separable.

References

2. F. B. Jones, On a property related to separability in metric spaces, Journal of the Elisha Mitchell Scientific Society, **70** (1954), 30-33.

THE UNIVERSITY OF TEXAS

^{1.} Paul Alexandroff, Uber die Metrization der im Kleinen Kompakten topologischen Raume, Mathematische Annalen, **92** (1924) 294-301.

^{3.} _____, A theorem concerning locally peripherally separable spaces, Bull. Amer. Math. Soc. **41** (1935), 437-439.

^{4.} R. L. Moore, Foundations of point set theory, Amer. Math. Soc. Colloq. Publ., vol 13 1932.

^{5.} G. T. Whyburn, Semi-locally-connected sets, Amer. J. Math. 61 (1939), 733-749.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DAVID GILBARG

Stanford University Stanford, California

F. H. BROWNELL

University of Washington Seattle 5, Washington

1015

A. L. WHITEMAN University of Southern California Los Angeles 7, California

L. J. PAIGE

University of California Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH	E. HEWITT	M. OHTSUKA	E. SPANIER
T. M. CHERRY	A. HORN	H. L. ROYDEN	E. G. STRAUS
D. DERRY	L. NACHBIN	M. M. SCHIFFER	F. WOLF

SUPPORTING INSTITUTIONS

STANFORD UNIVERSITY UNIVERSITY OF BRITISH COLUMBIA UNIVERSITY OF TOKYO CALIFORNIA INSTITUTE OF TECHNOLOGY UNIVERSITY OF CALIFORNIA UNIVERSITY OF UTAH WASHINGTON STATE COLLEGE MONTANA STATE UNIVERSITY UNIVERSITY OF WASHINGTON UNIVERSITY OF NEVADA NEW MEXICO STATE UNIVERSITY AMERICAN MATHEMATICAL SOCIETY OREGON STATE COLLEGE UNIVERSITY OF OREGON CALIFORNIA RESEARCH CORPORATION HUGHES AIRCRAFT COMPANY OSAKA UNIVERSITY SPACE TECHNOLOGY LABORATORIES UNIVERSITY OF SOUTHERN CALIFORNIA NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 2120 Oxford Street, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics Vol. 10, No. 2 October, 1960

Maynard G. Arsove, The Paley-Wiener theorem in metric linear spaces	365
Robert (Yisrael) John Aumann, Acceptable points in games of perfect	
information	381
A. V. Balakrishnan, Fractional powers of closed operators and the semigroups	
generated by them	419
Dallas O. Banks, Bounds for the eigenvalues of some vibrating systems	439
Billy Joe Boyer, On the summability of derived Fourier series	475
Robert Breusch, An elementary proof of the prime number theorem with	
remainder term	487
Edward David Callender, Jr., <i>Hölder continuity of n-dimensional</i> quasi-conformal mappings	499
L. Carlitz. Note on Alder's polynomials	517
P. H. Doyle, III, Unions of cell pairs in E^3	521
James Eells, Jr., A class of smooth bundles over a manifold	525
Shaul Foguel. <i>Computations of the multiplicity function</i>	539
James G. Glimm and Richard Vincent Kadison <i>Unitary operators in</i>	007
C^* -algebras	547
Hugh Gordon. Measure defined by abstract L _n spaces	557
Robert Clarke James, Separable conjugate spaces	563
William Elliott Jenner. On non-associative algebras associated with bilinear	
forms	573
Harold H. Johnson, Terminating prolongation procedures	577
John W. Milnor and Edwin Spanier, <i>Two remarks on fiber homotopy type</i>	585
Donald Alan Norton, A note on associativity	591
Ronald John Nunke, On the extensions of a torsion module	597
Joseph J. Rotman, Mixed modules over valuations rings	607
A. Sade, Théorie des systèmes demosiens de groupoï des	625
Wolfgang M. Schmidt, On normal numbers	661
Berthold Schweizer, Abe Sklar and Edward Oakley Thorp, <i>The metrization of statistical metric spaces</i>	673
John P. Shanahan <i>On uniqueness questions for hyperbolic differential</i>	0.0
equations	677
A. H. Stone, Sequences of coverings	689
Edward Oakley Thorp, <i>Projections onto the subspace of compact operators</i>	693
L. Bruce Treybig, <i>Concerning certain locally peripherally separable</i> <i>spaces</i>	697
Milo Wesley Weaver, On the commutativity of a correspondence and a	
permutation	705
David Van Vranken Wend, On the zeros of solutions of some linear complex differential equations	712
Fred Bover Wright Ir Polarity and duality	713
	123