

Pacific Journal of Mathematics

**MINIMUM PROBLEMS IN THE THEORY OF
PSEUDO-CONFORMAL TRANSFORMATIONS AND THEIR
APPLICATION TO ESTIMATION OF THE CURVATURE OF
THE INVARIANT METRIC**

JEREMIAH MILTON STARK

MINIMUM PROBLEMS IN THE THEORY OF PSEUDO- CONFORMAL TRANSFORMATIONS AND THEIR APPLICATION TO ESTIMATION OF THE CURVATURE OF THE INVARIANT METRIC

J. M. STARK

1. Introduction. Resume of some previous results.¹ Let B be a domain in the z_1, z_2 -space² possessing a Bergman kernel function $K^{(B)}(z_1, z_2; \bar{t}_1, \bar{t}_2), (z_1, z_2) \in B, (t_1, t_2) \in B$. By identifying the arguments $(t_1, t_2) = (z_1, z_2)$ one obtains the function $K^{(B)} \equiv K^{(B)}(z_1, z_2) \equiv K^{(B)}(z_1, z_2; \bar{z}_1, \bar{z}_2)$ which plays an essential role in the theory of pseudo-conformal transformations. An important application to this theory is the theorem proved by S. Bergman stating that the metric

$$(1.1) \quad ds^2 = \sum_{m, n=1}^2 T_{m\bar{n}}^{(B)} dz_m d\bar{z}_n, \quad T_{m\bar{n}} = \frac{\partial^2 \log K^{(B)}}{\partial z_m \partial \bar{z}_n}$$

is invariant under pseudo-conformal transformations (B. [1], p. 52). From this follows that all measures of geometric objects in B which are based on the metric (1.1) are also invariant under pseudo-conformal transformations.

In the present paper we are concerned in particular with the *Riemann Curvature of (1.1) in an analytic direction* (see definition in section 3). Since the second derivatives of the function $\log K^{(B)}(z_1, z_2)$ are the main constituent in the definition of the curvature, we at first discuss bounds for their distortion under pseudo-conformal transformation (see Theorem 1). For this purpose, *Bergman's method of the minimum integral* is used (B. [3], p. 48; K. [1]; S. [1]):

Relations among various solutions of minimum problems of the type

$$(1.2) \quad \int_B |f(z)|^2 d\omega \equiv \min = \lambda_B \quad (d\omega = \text{volume element}),$$

are studied (see Theorems 2 and 3). Here $f(z)$ are analytic functions,

Received September 28, 1959. The author wishes to thank Mr. M. Maschler for valuable help in preparing the manuscript for publication.

¹ Square brackets refer to the bibliography at the end of the paper. We use the abbreviations B. = Bergman, F. = Fuchs, K. = Kobayashi, S. = Stark.

² In the present paper we consider only domains in the space of two complex variables. The generalization of the methods to the space of more complex variables involves difficulties of technical nature only.

regular in B and subject to certain auxiliary conditions. By varying these conditions, one obtains different λ_B 's. (Upper and lower indices on λ_B indicate the auxiliary conditions, as described, e. g., at the end of this section.) The method of the minimum integral, which is applied in order to obtain bounds and distortion theorems for various quantities having a geometrical meaning, is based on the fact that these λ_B 's depend monotonically on the domain B (see (1.6), (1.7)). Indeed, if, for instance, one can express these quantities and/or their distortion in terms of the λ_B 's, and if one knows that there exist "domains of comparison" I and A such that $I \subset B \subset A$, then, using the relations among the λ 's, one can estimate the geometrical quantities and/or their distortion in terms of the λ_I 's and the λ_A 's. In general, I and A are required to be domains for which the kernel function can be expressed in a closed form; therefore, the various λ 's can be estimated if one knows how to express them in terms of the kernel function. This is done in B. [2], pp. 41-43, (see (1.5)), and in § 2, (see (2.2)).

Using the method of the minimum integral, Fuchs [1] has obtained an expression for the curvature in analytic direction R , in terms of certain λ_B 's, (see (3.6)). From this expression a bound for R is derived in terms of the corresponding λ_I 's and λ_A 's, where I and A are domains of comparison, $I \subset B \subset A$, (see (3.7)). It is shown in Theorem 4 and in the example which follows that this bound can be sharpened if a bound for the volume of B is given, and if a finite number of orthogonal functions in B and certain integrals over B with weighting functions depending only on A are known.

In order to prove some of the relations among the various λ 's, we use some results which were obtained in S. [1] and B. [4] p. 97 ff. These results and the definitions of the λ 's, used in S. [1], will be stated now for the convenience of the reader:

Consider the following general minimum problem: Let $\{\varphi^{(\nu)}(z)\}, \nu = 1, 2, \dots$, be a system of functions orthogonal in a domain B^3 and complete for the class $\mathcal{L}^2(B)$. Let $\alpha_{qp}, q = 1, 2, \dots, n, p = 1, 2, \dots$, be a system of complex numbers such that $\sum_{\nu=1}^{\infty} |\alpha_{q\nu}|^2 < \infty$ for $q = 1, 2, \dots, n$. Let X_1, \dots, X_n be complex numbers. Finally, let λ represent the minimum of the integral

$$(1.3) \quad \int_B |f|^2 d\omega = \sum_{\nu=1}^{\infty} A_{\nu} \bar{A}_{\nu}, \quad A_{\nu} = \int_B f \cdot \overline{\varphi^{(\nu)}} d\omega,$$

for functions $f \in \mathcal{L}^2(B)$ and satisfying

$$(1.4) \quad \sum_{\nu=m}^{\infty} A_{\nu} \alpha_{q\nu} = X_q, \quad q = 1, 2, \dots, n;$$

then ([Cf. B. (2), pp. 41-43; S. (1), (2.13)])

³ In the sense that $\int_B \varphi^{(\mu)}(z) \cdot \overline{\varphi^{(\nu)}(z)} d\omega = \delta_{\mu\nu}$, where $\delta_{\mu\nu} = 0$ for $\mu \neq \nu$, $\delta_{\nu\nu} = 1$. $\mathcal{L}^2(B)$ is the class of functions $f(z)$ which are regular in B and for which $\int_B |f(z)|^2 d\omega < \infty$.

$$(1.5) \quad \lambda = - \left| \begin{array}{cc} 0 & (\bar{X})' \\ (X) & (D) \end{array} \right| \div |(D)|,$$

where (X) is the column matrix of n rows having X_r as elements in the r th row, $(\bar{X})'$ is the transpose of (X) , conjugated; (D) is the square matrix of n rows having $\sum_{\nu=m}^{\infty} \alpha_{r\nu} \bar{\alpha}_{s\nu}$ as element in the r th row, s th column, and $|(D)|$ is the determinant of (D) .

Denote by (1)-(8) the auxiliary conditions

- (1) $f(t) = 1$
- (2) $f(t) = 0$
- (3) $f_{z_1}(t) = 1$
- (4) $f_{z_1}(t) = 0$
- (5) $f_{z_2}(t) = 1$
- (6) $f_{z_2}(t) = 0$
- (7) $\int_B f d\omega = 0$
- (8) $u_1(\partial f / \partial z_1)_t + u_2(\partial f / \partial z_2)_t = 1$, u_1, u_2 complex numbers;

and let

- | | | | |
|-------------------------|--|-------------------------|-------------------------|
| (a) λ_B^1 | (b) λ_B^{01} | (c) λ_B^{001} | (d) λ_B^{*1} |
| (e) $\lambda_B^{* * 1}$ | (f) $\lambda_B^{0 * 1}$ | (g) $\lambda_B^{* 0 1}$ | (h) $\lambda_B^{1 * 0}$ |
| (i) λ_B^{010} | (j) $\lambda_B^{* 10}$ | (k) λ_B^{10} | (l) $\lambda_B^{(2)}$ |
| (m) $\lambda_B^{(4)}$; | $(\lambda_B \equiv \lambda_B(t), t \in B)$, | | |

be the minima of the integral (1.2), for functions $f \in \mathcal{L}^2(B)$ which are normalized at $t \in B$ by the respective auxiliary conditions

- (a): (1);
- (b): (2) and (3);
- (c): (2), (4) and (5);
- (d): (3);
- (e): (5);
- (f): (2) and (5);
- (g): (4) and (5);
- (h): (1) and (6);
- (i): (2), (3) and (6);
- (j): (3) and (6);
- (k): (1) and (4);
- (l): (2) and (8);
- (m): (8).

Let G be a domain containing a domain B , $B \subset G$; we denote by

- | | | | |
|--------------------------|---|--------------------------|----------------------------|
| (n) λ_{GB}^1 | (o) λ_{GB}^{01} | (p) λ_{GB}^{001} | (q) $\lambda_{GB}^{0 * 1}$ |
| (r) λ_{GB}^{010} | (s) $\lambda_{GB}^{(2)}$; $(\lambda_B \equiv \lambda_B(t), t \in B)$, | | |

the minima of the integral

$$\int_G |f|^2 d\omega,$$

for functions $f \in \mathcal{L}^2(G)$ and normalized at $t \in B$ by the conditions

- (n): (1) and (7); (o): (2), (3) and (7);
 (p): (2), (4), (5) and (7); (q): (2), (5) and (7);
 (r): (2), (3), (6) and (7); (s): (2), (7) and (8) .

It follows from the definitions of the various λ 's that

$$(1.6) \quad \lambda_I \leq \lambda_B \leq \lambda_A \text{ for domains } I, B, A \text{ such that } I \subset B \subset A$$

$$(1.7) \quad \lambda_{AB} \geq \lambda_{BB} \quad \text{for domains } A, B \text{ such that } A \supset B$$

(See S. [1], (3.7a). (3.7b)). From these inequalities the following result can be deduced⁴:

LEMMA. *Let B be any domain with finite Euclidian volume, in the (z_1, z_2) -space, such that $\text{Vol } B \leq V < \infty$. Then if I and A are any domains $I \subset B \subset A$, we have*

$$(1.8) \quad (1/\lambda_B^1) \geq (1/\lambda_{AB}^1) + (1/V)$$

$$(1.9) \quad (1/\lambda_B^{01}) \geq \{1 - (\lambda_I^1/V)\} (1/\lambda_{AB}^{01}) + (\lambda_I^1/\lambda_A^{*1})(1/\lambda_A^{*1})$$

$$(1.10) \quad (1/\lambda_B^{001}) \geq \{1 - (\lambda_I^{10}/V)\} (1/\lambda_{AB}^{001}) + (\lambda_I^{10}/V)(1/\lambda_A^{*01})$$

$$(1.11) \quad (1/\lambda_B^{0*1}) \geq \{1 - (\lambda_I^1/V)\} (1/\lambda_{AB}^{0*1}) + (\lambda_I^1/V)(1/\lambda_A^{**1})$$

$$(1.12) \quad (1/\lambda_B^{010}) \geq \{1 - (\lambda_I^{*0}/V)\} (1/\lambda_{AB}^{010}) + (\lambda_I^{*0}/V)(1/\lambda_A^{*10})$$

$$(1.13) \quad (1/\lambda_B^{(2)}) \geq \{1 - (\lambda_I^1/V)\} (1/\lambda_{AB}^{(2)}) + (\lambda_I^1/V)(1/\lambda_A^{(4)})$$

at $t \equiv (t_1, t_2) \in I$.

2. Distortion theorems under some assumptions about the structure of the domain. If integrals over a domain B of the type

$$(2.1) \quad \int_B K^{(A)}(\zeta, \bar{t}) d\omega_\zeta, \quad \int_B (\partial K^{(A)}/\partial \bar{z}_\nu)_{(\zeta, \bar{t})} d\omega_\zeta, \\ \int_B \int_B K^{(A)}(\zeta, \bar{\xi}) d\omega_\zeta d\omega_\xi, \quad \nu = 1, 2,$$

are known, where A is a domain which contains the domain B , then in (1.8)–(1.13) the terms involving the λ_{AB} can be evaluated. For, if λ_{AB} is any one of the λ 's with double subindex in (1.8)–(1.13), then the relation between λ_{AB} and the λ_A which has the same upper indices is described as follows: Let $\psi^{(\nu)}(z)$, $z \equiv (z_1, z_2) \in A$, $\nu=1, 2, \dots$, be a complete orthonormal system of functions for the class $\mathcal{L}^2(A)$, then each function $f(z)$ of this class can be represented in the form: $f(z) = \sum_{\nu=1}^{\infty} A_\nu \varphi^{(\nu)}(z)$, and the series converges absolutely and uniformly in any closed subdomain of A . Therefore, each of the λ_A 's is a special case of the general minimum problem described in (1.3), (1.4)⁵. Thus it follows from (1.5)

⁴ This is Theorem 2 of (S. [1]).

⁵ In these formulas replace B by A .

that each λ_A can be written in the form $\lambda_A = -|(N)| \div |(D)|$, where $|(D)|$ is as in (1.5) and (N) is the matrix whose determinant occurs in the numerator of (1.5).

Since $K^{(A)}(z, \bar{\xi}) = \sum \psi^{(\nu)}(z) \overline{\psi^{(\nu)}(\bar{\xi})}$, the matrices (N) and (D) depend only upon $K^{(A)}(z, \bar{\xi})$ and its derivatives at the points $z = t$, $\bar{\xi} = \bar{t}$, and in the case of $\lambda_A^{(2)}$ also upon u_1, u_2 .

LEMMA. If $\lambda_A = -|(N)| \div |(D)|$, where (N) and (D) are used as explained above, then the corresponding λ_{AB} , i. e., the λ_{AB} which has the same upper indices as λ_A , can be expressed in the form

$$(2.2) \quad \lambda_{AB} = - \left| \begin{pmatrix} (N) & (\bar{U})' \\ (U) & E \end{pmatrix} \right| \div \left| \begin{pmatrix} (D) & (\bar{W})' \\ (W) & E \end{pmatrix} \right|.$$

Here

$$E \equiv \int_B \int_B K^{(A)}(\zeta, \bar{\xi}) d\omega_\zeta d\omega_{\bar{\xi}}$$

and

$$\begin{aligned} (W) &\equiv (W_1, W_2, \dots, W_n), \\ (U) &\equiv (0, W_1, W_2, \dots, W_n) \end{aligned}$$

are row matrices⁶, and $(\bar{U})'$ is the transpose of the row matrix whose elements are the conjugates of the elements of (U) ; the same rule applies to $(W)'$. The elements W_i depend only upon the expressions (2.1), and in the case of $\lambda_{AB}^{(2)}$ also upon u_1, u_2 .

E. g., for

$$\lambda_{AB} = \lambda_{AB}^1, \lambda_{AB}^{01}, \lambda_{AB}^{001}, \lambda_{AB}^{(2)},$$

we have, using the notation

$$K_{\nu\mu uv}^{(A)}(\zeta, \bar{t}) \equiv [\partial^{\nu+} \mu + u + v K^{(A)}(\zeta, \bar{t}) / \partial \zeta_1^{\nu} \partial \zeta_2^{\mu} \partial \bar{t}_1^u \partial \bar{t}_2^v],$$

$$\zeta \equiv (\zeta_1, \zeta_2), \quad \bar{t} \equiv (\bar{t}_1, \bar{t}_2),$$

$$(W) = \left(\int_B K^{(A)}(\zeta, \bar{t}) d\omega_\zeta \right),$$

$$(W) = \left(\int_B K^{(A)}(\zeta, \bar{t}) d\omega_\zeta, \int_B K_{0010}^{(A)}(\zeta, \bar{t}) d\omega_\zeta \right),$$

$$(W) = \left(K^{(A)}(\zeta, \bar{t}) d\omega_\zeta, \int_B K_{0010}^{(A)}(\zeta, \bar{t}) d\omega_\zeta, \int_B K_{0001}^{(A)}(\zeta, \bar{t}) d\omega_\zeta \right),$$

$$(W) = \left(K^{(A)}(\zeta, \bar{t}) d\omega_\zeta, \bar{u}_1 \int_B K_{0010}^{(A)}(\zeta, \bar{t}) d\omega_\zeta + \bar{u}_2 \int_B K_{0001}^{(A)}(\zeta, \bar{t}) d\omega_\zeta \right),$$

respectively.

⁶ Note that (X) in (1.5) is a column matrix.

Proof. We shall prove (2.2) only for the case of λ_{AB}^{01} . The proof for the other cases can be carried out along the same lines.

Let $\varphi_\nu(z)$ be a complete orthogonal system for the class $\mathcal{L}^2(A)$. Choosing $n = 2, m = 1, \alpha_{1\nu} = \varphi_\nu(t)$,

$$\alpha_{2\nu} = \left. \frac{\partial \varphi_\nu(z_1, z_2)}{\partial z_1} \right|_{z=t}, \quad X_1 = 0, X_2 = 1,$$

the general minimum problem of § 1 (B is replaced by A) becomes the minimum problem for λ_A^{01} .

The elements of the matrix (D) in (1.5) become values of the partial derivatives of the kernel function and (1.5) is reduced to

$$\lambda_A^{01} = - \frac{|S|}{|S_{1I}|}$$

where

$$S = \begin{pmatrix} 0 & 0 & 1 \\ 0 & K_{0000}^{(A)} & K_{0010}^{(A)} \\ 1 & K_{1000}^{(A)} & K_{1010}^{(A)} \end{pmatrix}$$

and S_{1I} is the matrix which one obtains by deletion of the first row and the first column from the matrix S . Here

$$K_{\gamma\mu uv}^{(A)} = \partial^{+\mu+u+v} K^{(A)}(z, \bar{t}) / \partial z_1^\gamma \partial z_2^\mu \partial \bar{t}_1^u \partial \bar{t}_2^v \Big|_{(z_1, z_2) = (t_1, t_2)}.$$

If we choose $m = 1, n = 3, \alpha_{1\nu} = \varphi_\nu(t), \alpha_{2\nu} = \partial \varphi_\nu(z_1, z_2) / \partial z_1|_{z=t}$

$$\alpha_{3\nu} = \int_B \varphi_\nu(\xi) d\omega_\xi, \quad X_1 = 0, X_2 = 1, X_3 = 0$$

then the same general minimum problem becomes the minimum problem for λ_{AB}^{01} , and (1.5) becomes $\lambda_{AB}^{01} = - |T| / |T_{1I}|$ where

$$T = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & K_{0000}^{(A)} & K_{0010}^{(A)} & \int_B K_{0000}^{(A)}(t, \bar{\xi}) d\omega_\xi \\ 1 & K_{1000}^{(A)} & K_{1010}^{(A)} & \int_B K_{1000}^{(A)}(t, \bar{\xi}) d\omega_\xi \\ 0 & \int_B K_{0000}^{(A)}(\xi, \bar{t}) d\omega_\xi & \int_B K_{0010}^{(A)}(\xi, \bar{t}) d\omega_\xi & \int_B \int_B K_{0000}^{(A)}(\xi, \bar{\xi}) d\omega_\xi d\omega_\xi \end{pmatrix}$$

and T_{1I} is the matrix which one obtains by deletion of the first row and the first column from the matrix T . Since $K^{(A)}(z, \bar{t}) = K^{(A)}(t, \bar{z})$, we have that this expression for λ_{AB}^{01} is equivalent to (2.2).

Other assumptions about B permit further estimates: E.g., let $\{\varphi_0^{(\sigma)}(z_1, z_2)\}$, $\sigma = 1, 2, \dots, r-1$, $\varphi_0^{(1)}(z_1, z_2) = \text{constant}$, be a set of independent functions of the class $\mathcal{L}^2(B)$. Let $\{\varphi^{(\sigma)}(z_1, z_2)\}$, $\sigma = 1, 2, \dots, r-1$, be functions obtained by orthonormalizing over B the set $\{\varphi_0^{(\sigma)}(z_1, z_2)\}$, $\sigma = 1, 2, \dots, r-1$. If we assume that integrals over B of the type

$$(2.3) \quad \begin{aligned} U_n &= \int_B \overline{\varphi_0^{(n)}(\xi)} K^{(A)}(\xi, \bar{t}) d\omega_\xi \\ P_{mn} &= \int_B \int_B \varphi_0^{(m)}(\xi) \overline{\varphi_0^{(n)}(\bar{\xi})} K^{(A)}(\xi, \bar{\xi}) d\omega_\xi d\omega_{\bar{\xi}} \\ m, n &= 1, 2, \dots, r-1 \end{aligned}$$

are known, then we can use the relation

$$(2.4) \quad (1/\lambda_B^1) \geq (1/\lambda_{ABr}^1) + \sum_{\sigma=1}^{r-1} |\varphi^{(\sigma)}|^2,$$

where

$$(2.5) \quad \lambda_{ABr}^1 = |(P_{mn})| \div \begin{vmatrix} K^{(A)} & (U_n) \\ (\bar{U}_n)' & (P_{mn}) \end{vmatrix}.$$

Here (P_{mn}) is the square matrix whose elements P_{mn} are given by (2.3); $|(P_{mn})|$ is the determinant of (P_{mn}) , (U_n) is the row matrix $(U_1 U_2 \dots U_{r-1})$, and $(\bar{U}_n)'$ is the transpose of (U_n) conjugated.⁷ Notice that $\lambda_{AB2}^1 = \lambda_{AB}^1$.

Let

$$\begin{aligned} &\{\varphi_0^{(\sigma)}(z_1, z_2)\}, \quad \sigma = 1, 2, \dots, r-1, \\ &\{\alpha_0^{(\nu)}(z_1)\}, \nu = 1, 2, \dots, p-1, \{\beta_0^{(\mu)}(z_2)\}, \mu = 1, 2, \dots, q-1 \end{aligned}$$

be sets of functions⁸ such that each set consists of independent functions and each function belongs to $\mathcal{L}^2(B)$. Further, let $\{\varphi^{(\cdot)}\}$, $\{\alpha^{(\nu)}\}$, $\{\beta^{(\mu)}\}$, $\nu = 1, 2, \dots$, be sets of orthonormal functions such that each set is complete for $\mathcal{L}^2(B)$, and such that the first functions of each sequence are obtained respectively by orthonormalizing over B the sets $\{\varphi_0^{(\sigma)}\}$, $\{\alpha_0^{(\nu)}\}$ and $\{\beta_0^{(\mu)}\}$. For any domain G , $G \supset B$, we define λ_{GB**q}^{01} , λ_{GBr}^{*1} , λ_{GB**p}^{0*1} , λ_{GBr}^{**1} , in the same manner as we defined λ_{GB}^{01} , λ_{GB}^{*1} , λ_{GB}^{0*1} , λ_{GB}^{**1} , except that auxiliary condition (7) of § 1 is replaced respectively by the conditions

$$(2.6) \quad \begin{cases} \int_B \overline{\beta_0^{(\mu)}} f d\omega = 0, \mu = 1, 2, \dots, q-1; \\ \int_B \overline{\varphi_0^{(\sigma)}} f d\omega = 0, \sigma = 1, 2, \dots, r-1; \\ \int_B \overline{\alpha_0^{(\nu)}} f d\omega = 0, \nu = 1, 2, \dots, p-1; \text{ and} \\ \int_B \overline{\beta_0^{(\mu)}} f d\omega = 0, \mu = 1, 2, \dots, q-1. \end{cases}$$

⁷ A formula similar to (2.4) is proved in B. [1, 4] for the case in which the domains are two-dimensional. (Extension to the case of four-dimensional domains offers no difficulty.)

⁸ E. g., $\{\alpha_0^{(\nu)}\} \equiv \{1, z_1, \dots, z_1^{p-2}\}$, $\{\beta_0^{(\mu)}\} \equiv \{1, z_2, \dots, z_2^{q-2}\}$, $\{\varphi_0^{(\sigma)}\} \equiv \{1, z_1, z_2, \dots, z_1^a \cdot z_2^b\}$.

REMARK. Conditions (2.6) do not change if one replaces $\beta_0^{(\mu)}, \varphi_0^{(\sigma)}, \alpha_0^{(\nu)}$, by other linearly independent functions which are respectively linear combinations of the previous functions. In particular, these functions can be replaced by $\beta^{(\mu)}, \varphi^{(\sigma)}, \alpha^{(\nu)}$, respectively.

THEOREM 1. *Let B^* be the image of a domain B under a pseudo-conformal transformation $z_k^* = z_k^*(z_1, z_2)$, $k = 1, 2$, normalized at t by $(\partial z_k^* / \partial z_i)_t = \delta_{ki}$. Here $t \equiv (t_1, t_2) \in B$, $z_k^*(t_1, t_2) \equiv t_k^*$, $\delta_{\nu\nu} = 1$, and $\delta_{\nu\mu} = 0$ for $\nu \neq \mu$. Further, let I and A be domains of comparison for B such that $t \in I \subset B \subset A$. Finally, let $\{\varphi^{(\sigma)}(z_1, z_2)\}$, $\sigma = 1, 2, \dots, r-1$, $\{\alpha^{(\nu)}(z_1)\}$, $\nu = 1, 2, \dots, p-1$, and $\{\beta^{(\mu)}(z_2)\}$, $\mu = 1, 2, \dots, q-1$, be three sets of functions possessing the properties*

$$(2.7) \quad \int_B \varphi^{(\sigma)} \overline{\varphi^{(k)}} d\omega = \delta_{\sigma k}, \int_B \alpha^{(\nu)} \overline{\alpha^{(k)}} d\omega = \delta_{\nu k}, \int_B \beta^{(\mu)} \overline{\beta^{(k)}} d\omega = \delta_{\mu k}.$$

Then⁹

$$(2.8a) \quad \begin{aligned} & \left\{ \lambda_I^1 \left(1 - \lambda_I^1 \sum_{\mu=1}^{q-1} |\beta^{(\mu)}|^2 \right) (1/\lambda_{AB^*}^{01} * q) \right. \\ & \quad + (\lambda_I^1)^2 \sum_{\mu=1}^{q-1} |\beta^{(\mu)}|^2 \left[(1/\lambda_{AB^*}^{*1}) + \sum_{\sigma=1}^{r-1} |\varphi_{z_1}^{(\sigma)}|^2 \right] \Big\}_{z=t} \\ & \leq [\partial^2 \log K^{(B^*)}(z^*, \bar{t}^*) / \partial z_1^* \partial \bar{t}_2^*]_{z^*=t^*} \\ & \leq \left\{ (1/\lambda_I^{01}) \left[(1/\lambda_{AB^*}^1) + \sum_{\sigma=1}^{r-1} |\varphi^{(\sigma)}|^2 \right] \right\}_{z=t}^{10} \end{aligned}$$

and

$$(2.8b) \quad \begin{aligned} & \left\{ \lambda_I^1 \left(1 - \lambda_I^1 \sum_{\nu=1}^{p-1} |\alpha^{(\nu)}|^2 \right) (1 - \lambda_{AB^*}^{0*1}) \right. \\ & \quad + (\lambda_I^1)^2 \sum_{\nu=1}^{p-1} |\alpha^{(\nu)}|^2 \left[(1/\lambda_{AB^*}^{*1}) + \sum_{\sigma=1}^{r-1} |\varphi_{z_2}^{(\sigma)}|^2 \right] \Big\}_{z=t} \\ & \leq [\partial^2 \log K^{(B^*)}(z^*, \bar{t}^*) / \partial z_2^* \partial \bar{t}_1^*]_{z^*=t^*} \\ & \leq \left\{ (1/\lambda_I^{0*1}) \left[(1/\lambda_{AB^*}^1) + \sum_{\sigma=1}^{r-1} |\varphi^{(\sigma)}|^2 \right] \right\}_{z=t}. \end{aligned}$$

The λ 's bearing multiple subscripts are functions of

$$K^{(A)}(t, \bar{t}), K_{\nu\mu\bar{u}\bar{v}}^{(A)} \equiv [\partial^{\nu+\mu+u+v} K^{(A)}(z, \bar{t}) / \partial z_1^\nu \partial z_2^\mu \partial \bar{t}_1^u \partial \bar{t}_2^v]_{z=t}$$

$\nu + \mu + u + v = 1, 2$, and a finite number of integrals over B with weighting functions depending only upon A .

REMARKS: For two given domains, the theorem gives necessary conditions in terms of various properties that one of the domains can

⁹ We use the abbreviations $z \equiv (z_1, z_2)$, $\bar{t} \equiv (\bar{t}_1, \bar{t}_2)$.

¹⁰ Concerning the symbols $\lambda_{AB^*}^1$ see pp. 6 and 14.

be mapped onto the other by a transformation of the type described. The middle terms in (2.8a), (2.8b) depend only upon B^* , t^* .

For our proof we need certain relations between the λ 's which we formulate in the

THEOREM 2. *The following relations hold:*

$$(2.9) \quad (1/\lambda_B^{01}) = (1/\lambda_{BB**q}^{01}) + \lambda_B^1 \cdot \sum_{\mu=1}^{q-1} |\beta^{(\mu)}|^2 \cdot \{(1/\lambda_B^{*1}) - (1/\lambda_{BB**q}^{01})\},$$

$$(2.10) \quad (1/\lambda_B^{0*1}) = (1/\lambda_{BB*p}^{0*1}) + \lambda_B^1 \cdot \sum_{\nu=1}^{p-1} |\alpha^{(\nu)}|^2 \cdot \{(1/\lambda_B^{**1}) - (1/\lambda_{BB*p}^{0*1})\},$$

$$(2.11) \quad (1/\lambda_B^{*1}) = (1/\lambda_{BBr}^{*1}) + \sum_{\sigma=1}^{r-1} |\varphi_{z_1}^{(\sigma)}|^2,$$

$$(2.12) \quad (1/\lambda_B^{**1}) = (1/\lambda_{BBr}^{**1}) + \sum_{\sigma=1}^{r-1} |\varphi_{z_2}^{(\sigma)}|^2,$$

$$(2.13) \quad (1/\lambda_B^{01}) \geq \left(1 - \lambda_I^1 \cdot \sum_{\mu=1}^{q-1} |\beta^{(\mu)}|^2\right) (1/\lambda_{AB**q}^{01}) \\ + \lambda_I^1 \cdot \sum_{\mu=1}^{q-1} |\beta^{(\mu)}|^2 \left\{ (1/\lambda_{ABr}^{*1}) + \sum_{\sigma=1}^{r-1} |\varphi_{z_1}^{(\sigma)}|^2 \right\},$$

$$(2.14) \quad (1/\lambda_B^{0*1}) \geq \left(1 - \lambda_I^1 \cdot \sum_{\nu=1}^{p-1} |\alpha^{(\nu)}|^2\right) (1/\lambda_{AB*p}^{0*1}) \\ + \lambda_I^1 \cdot \sum_{\nu=1}^{p-1} |\alpha^{(\nu)}|^2 \left\{ (1/\lambda_{ABr}^{**1}) + \sum_{\sigma=1}^{r-1} |\varphi_{z_2}^{(\sigma)}|^2 \right\},$$

Proof. To establish (2.9) we evaluate λ_{BB**q}^{01} , λ_B^{01} , λ_B^{*1} , and λ_B^1 using (1.5) by taking for $(m, n, X_1, X_2, \alpha_{1\nu}, \alpha_{2\nu})$ the values $(q, 2, 0, 1, \beta^{(\nu)}, \beta_{z_1}^{(\nu)})$, $(1, 2, 0, 1, \beta^{(\nu)}, \beta_{z_1}^{(\nu)})$, $(1, 1, 1, -, \beta_{z_1}^{(\nu)}, -)^{11}$ and $(1, 1, 1, -, \beta^{(\nu)}, -)$ respectively, where $\beta^{(\nu)}$ and $\beta_{z_1}^{(\nu)}$ are evaluated at $t = (t_1, t_2)$. Now all the λ 's of (2.9) are expressed in terms of $\{\beta^{(\nu)}\}$, $\nu = 1, 2, \dots$, and their derivatives at t , and the relation between these λ 's is easily verified to be (2.9). Equation (2.10)-(2.12) are established in the same manner by using in (1.5) for λ_{BB*p}^{0*1} , λ_B^{0*1} , λ_B^{**1} , λ_B^1 , λ_{BBr}^{*1} , λ_B^{*1} , λ_{BBr}^{**1} , λ_B^{**1} , values of $(m, n, X_1, X_2, \alpha_{1\nu}, \alpha_{2\nu})$ respectively as follows: $(p, 2, 0, 1, \alpha^{(\nu)}, \alpha_{z_2}^{(\nu)})$, $(1, 2, 0, 1, \alpha^{(\nu)}, \alpha_{z_2}^{(\nu)})$, $(1, 1, 1, -, \alpha_{z_2}^{(\nu)}, -)$, $(1, 1, 1, -, \alpha^{(\nu)}, -)$, $(r, 1, 1, -, \varphi_{z_1}^{(\nu)}, -)$, $(1, 1, 1, -, \varphi_{z_1}^{(\nu)}, -)$, $(r, 1, 1, -, \varphi_{z_2}^{(\nu)}, -)$, and $(1, 1, 1, -, \varphi_{z_2}^{(\nu)}, -)$.

From the relations $(1/\lambda_B^1) \geq \sum_{\mu=1}^{\infty} |\beta^{(\mu)}|^2$, $(1/\lambda_B^{*1}) \geq \sum_{\nu=1}^{\infty} |\alpha^{(\nu)}|^2$, it follows that the coefficients before the braces in (2.9) and (2.10) are each less than or equal to 1. By essentially the same reasoning used to derive (1.7) (see S. [1], (3.7b)), we obtain.

$$(2.15) \quad \lambda_{ABr} \geq \lambda_{BBr}, \lambda_{AB*p} \geq \lambda_{BB*p}, \lambda_{AB**q} \geq \lambda_{BB**q}, A \supset B.$$

¹¹ “—” means no special value is required for this quantity.

Equations (2.9) and (2.10) are of the form¹²

$$(2.16) \quad (1/\lambda_B) = (1/\lambda_{BB(S)}) + \lambda_B^\infty \cdot g \cdot \{(1/\lambda_B^{(*)}) - (1/\lambda_{BB(S)})\} ,$$

where $\lambda_B^\infty \cdot g \leq 1$, and where the auxiliary condition associated with $\lambda_B^{(*)}$ are among the auxiliary conditions associated with $\lambda_{BB(S)}$. Hence $\lambda_{BB(S)} \geq \lambda_B^{(*)}$, and the brace in (2.16) is non-negative. By use of (2.15), and since

$$\begin{aligned} (1/\lambda_B) &\geq (1/\lambda_{BB(S)}) + \lambda_I^\infty \cdot g \{(1/\lambda_B^{(*)}) - (1/\lambda_{BB(S)})\} \\ &= (1 - \lambda_I^\infty \cdot g)(1/\lambda_{BB(S)}) + \lambda_I^\infty \cdot g(1/\lambda_B^{(*)}) , \end{aligned}$$

we have

$$(2.17) \quad (1/\lambda_B) \geq (1 - \lambda_I^\infty \cdot g)(1/\lambda_{AB(S)}) + \lambda_1^\infty \cdot g(1/\lambda_A^{(*)}) .$$

Using (2.15) in (2.11) and (2.12), and substituting the resulting inequalities into (2.17), we obtain (2.13) and (2.14). This completes the proof of Theorem 2 and we begin with the proof of the Theorem 1.

Since

$$T_{\mu\nu}^{*-} = \sum_{p,q=1}^2 T_{p\bar{q}}^- \frac{\partial z_p}{\partial z_\mu^*} \frac{\partial \bar{z}_q}{\partial \bar{z}_\nu^*}$$

(see (1.1)), it follows from the normalization that

$$\begin{aligned} T_{11}^{(B)}(t_1, t_2) &= T_{11}^{(B^*)}(t_1^*, t_2^*) , \\ T_{22}^{(B)}(t_1, t_2) &= T_{22}^{(B^*)}(t_1^*, t_2^*) . \end{aligned}$$

From the relation

$$K^{(B)}(z_1, z_2; \bar{z}_1, \bar{z}_2) = K^{(B^*)}(z_1^*, z_2^*; \bar{z}_1^*, \bar{z}_2^*) \left| \frac{\partial(z_1^*, z_2^*)}{\partial(z_1, z_2)} \right|^2$$

(see B. [2]), it follows that the last two equalities are equivalent to

$$(\lambda_B^1/\lambda_B^{01})_t = T_{11}^{(B^*)} \equiv [\partial^2 \log K^{(B^*)}(z_1^*, z_2^*)/\partial z_1^* \partial \bar{z}_1^*]_{(t_1^*, t_2^*)}$$

and

$$(\lambda_B^1 \lambda_B^{0*1})_t = T_{22}^{(B^*)} \equiv [\partial^2 \log K^{(B^*)}(z_1^*, z_2^*)/\partial z_2^* \partial \bar{z}_2^*]_{(t_1^*, t_2^*)} .$$

Using the bounds for λ_B^1 , λ_B^{01} , and λ_B^{0*1} as given by (2.4), (2.13), and (2.14), we obtain (2.8a) and (2.8b).

To complete the proof of Theorem 1 we need only show that in (2.8a) and (2.8b) the λ 's bearing multiple subscripts are expressible in

¹² (2.16) becomes (2.9) if we set $\lambda_B = \lambda_B^{01}$, $\lambda_{BB(S)} = \lambda_{BB^{**}q}^{01}$, $\lambda_B^\infty = \lambda_B^1$, $\lambda_B^{(*)} = \lambda_B^{*1}$ and $g = \sum_{\mu=1}^{q-1} |\beta^{(\mu)}|^2$; (2.16) becomes (2.10) if we set $\lambda_B = \lambda_B^{0*1}$, $\lambda_{BB(S)} = \lambda_{BB^{**}p}^{0*1}$, $\lambda_B^\infty = \lambda_B^1$, $\lambda_B^{(*)} = \lambda_B^{*1}$, and $g = \sum_{\nu=1}^{p-1} |\alpha^{(\nu)}|^2$. The symbol (S) in $\lambda_{BB(S)}$ means that some additional conditions are superimposed in addition to the conditions (n) - (s), see p. 6.

terms of $K^{(A)}$, $K_{\nu\mu\bar{u}\bar{v}}^{(A)}$, $\nu + \mu + u + v = 1, 2$, and a finite number of integrals over B with certain weighting functions independent of B . That this is true for λ_{ABr}^1 is shown by (2.4). Let $\{\psi^{(\nu)}\}$, $\nu = 1, 2, \dots$, be a system of orthonormal functions over A complete for $\mathcal{L}^2(A)$. λ_{ABr}^1 is the minimum of

$$\sum_{\nu=1}^{\infty} A_{\nu} \bar{A}_{\nu}, \quad A_{\nu} = \int_A f \overline{\psi^{(\nu)}} d\omega,$$

for functions $f \in \mathcal{L}^2(A)$ and satisfying

$$\begin{aligned} f_{z_1}(t) &= \sum_{\nu=1}^{\infty} A_{\nu} \psi_{z_1}^{(\nu)}(t) = 1, \quad \int_B \overline{\varphi_0^{(\mu)}}(\zeta) \cdot f(\zeta) d\omega_{\zeta} \\ &= \sum_{\nu=1}^{\infty} A_{\nu} \int_B \overline{\varphi_0^{(\mu)}}(\zeta) \cdot \psi^{(\nu)}(\zeta) d\omega_{\zeta} = 0 \quad \mu = 1, 2, \dots, r-1. \end{aligned}$$

Thus λ_{ABr}^{*1} can be evaluated using (1.5) and taking $m = X_1 = 1, n = r, X_2 = \dots = X_r = 0, \alpha_{1\nu} = \psi_{z_1}^{(\nu)}, \alpha_{k\nu} = \int_B \overline{\varphi_0^{(k-1)}}(\zeta) \cdot \psi^{(\nu)}(\zeta) d\omega_{\zeta}, 2 \leq k \leq r$. Likewise $\lambda_{ABr}^{**1}, \lambda_{AB**q}^{01}$ and λ_{AB**p}^{0*1} are evaluated in the form required by the theorem by substituting in (1.5) the values

$$\begin{aligned} (2.18) \quad m &= 1, n = r, X_1 = 1, X_2 = \dots = X_r = 0, \\ \alpha_{1\nu} &= \psi_{z_2}^{(\nu)}, \alpha_{k\nu} = \int_B \overline{\varphi_0^{(k-1)}}(\zeta) \psi^{(\nu)}(\zeta) d\omega_{\zeta}, 2 \leq k \leq r; \end{aligned}$$

$$\begin{aligned} (2.19) \quad m &= 1, n = q + 1, X_1 = 0, X_2 = 1, X_3 = \dots \\ &= X_{q+1} = 0, \alpha_{1\nu} = \psi_{z_1}^{(\nu)}, \alpha_{2\nu} = \psi_{z_1}^{(\nu)}, \alpha_{k\nu} \\ &= \int_B \overline{\beta_0^{(k-2)}}(\zeta_2) \psi^{(\nu)}(\zeta) d\omega_{\zeta}, 3 \leq k \leq q + 1; \end{aligned}$$

$$\begin{aligned} (2.20) \quad m &= 1, n = p + 1, X_1 = 0, X_2 = 1, X_3 = \dots \\ &= X_{p+1} = 0, \alpha_{1\nu} = \psi_{z_1}^{(\nu)}, \alpha_{2\nu} = \psi_{z_2}^{(\nu)}, \alpha_{k\nu} \\ &= \int_B \overline{\alpha_0^{(k-2)}}(\zeta_1) \psi^{(\nu)}(\zeta) d\omega_{\zeta}, 3 \leq k \leq p + 1; \end{aligned}$$

respectively. This completes the proof of the theorem.

3. Curvature in an analytic direction. In this paragraph we consider the Riemann curvature of Bergman's metric.

$$(3.1) \quad ds_B^2 = \sum_{m,n=1}^2 T_{m\bar{n}}^{(B)} dz_m d\bar{z}_n, \quad T_{m\bar{n}}^{(B)} \equiv T_{m\bar{n}} = \frac{\partial^2 \log K^{(B)}}{\partial z_m \partial \bar{z}_n}$$

is the metric defined in a domain B , where the formal operations are carried out as if $z_1, z_2, \bar{z}_1, \bar{z}_2$ were independent coordinates. The components of the fundamental tensor of the Riemannian geometry defined by (3.1) are then

$$\begin{aligned} g_{11} &= g_{12} = g_{22} = g_{1\bar{1}} = g_{1\bar{2}} = g_{2\bar{2}} = 0, \\ g_{i\bar{i}} &= \frac{1}{2} T_{i\bar{i}}, \quad g_{i\bar{j}} = \frac{1}{2} T_{i\bar{j}}, \quad g_{2\bar{1}} = \frac{1}{2} T_{2\bar{1}}, \quad g_{2\bar{2}} = \frac{1}{2} T_{2\bar{2}}, \end{aligned}$$

where now

$$(3.2) \quad \sum_{\mu, \nu=1, 2, \bar{1}, \bar{2}} g_{\mu\nu} dz_{\mu} dz_{\nu} = \sum_{m, n=1}^2 T_{m\bar{n}} dz_m d\bar{z}_n^{13}$$

Taking the usual formula for the Riemannian curvature of the metric $\sum_{\mu, \nu=1, 2, \bar{1}, \bar{2}} g_{\mu\nu} dz_{\mu} dz_{\nu}$ in the plane defined by the vectors $\{u_{\alpha}\}, \{v_{\alpha}\}, \alpha = 1, 2, \bar{1}, \bar{2} (v_{\bar{\alpha}} = \bar{u}_{\alpha}, v_{\alpha} = \bar{v}_{\alpha})$ we obtain

$$(3.3) \quad \frac{\sum R_{n\mu\nu k} u_n v_{\mu} u_{\nu} v_k}{\sum (g_{n\nu} g_{\mu k} - g_{nk} g_{\mu\nu}) u_n v_{\mu} u_{\nu} v_k}$$

where

$$\sum \equiv \sum_{n, \mu, \nu, k=1, 2, \bar{1}, \bar{2}}$$

and $R_{n\mu\nu k}$ are the usual Riemann symbols of the first kind.

If $\{v_{\alpha}\}$ belongs to the same analytic plane as $\{u_{\alpha}\}$ (i.e., if $v_{\alpha} = \alpha u_{\alpha}$, $v_{\bar{\alpha}} = \bar{\alpha} u_{\bar{\alpha}}$, then (3.3) becomes what is called the curvature in the analytic direction $\{u_{\alpha}\}, \alpha = 1, 2$ (B. [2], p. 54)

$$R = \frac{\sum R_{n\mu\nu k} \bar{u}_n u_{\mu} u_{\nu} \bar{u}_k}{\sum T_{n\mu} T_{\nu k} \bar{u}_n u_{\mu} u_{\nu} \bar{u}_k}$$

where

$$\begin{aligned} \sum &\equiv \sum_{n, \mu, \nu, k=1}^2, R_{\bar{\lambda}\alpha\mu\bar{\beta}} \frac{\partial^2 T_{\mu\bar{\lambda}}}{\partial z_{\alpha} \partial \bar{z}_{\beta}} + \sum_{\rho, k=1}^2 T_{k\rho} \cdot \frac{\partial T_{\mu\rho}}{\partial z_{\alpha}} \cdot \frac{\partial T_{k\bar{\lambda}}}{\partial \bar{z}_{\beta}} \\ T^{1\bar{1}} &= T_{2\bar{2}}/D, T^{1\bar{2}} = -T_{2\bar{1}}/D, T^{2\bar{1}} = -T_{1\bar{2}}/D, T^{2\bar{2}} = T_{1\bar{1}}/D, \\ D &= T_{1\bar{1}} T_{2\bar{2}} - T_{1\bar{2}} \cdot T_{2\bar{1}}. \end{aligned}$$

Using Bergman's method of the minimum integral, Fuchs [1] has obtained the following result. Let $\lambda_B^{[3]} = \lambda_B^{[3]}(t), t = (t_1, t_2) \in B$ denote the minimum of the integral

$$(3.4) \quad \int_B |f|^2 d\omega, d\omega = dx_1 dy_1 dx_2 dy_2$$

for functions $f \in \mathcal{L}^2(B), f \equiv f(z) \equiv f(z_1, z_2), z_k = x_k + iy_k, k = 1, 2$, and normalized by the auxiliary conditions

$$f(t) = f_{10}(t) = f_{01}(t) = 0, u_1^2 f_{20}(t) + 2u_1 u_2 f_{11}(t) + u_2^2 f_{02}(t) = 1$$

where $f_{mn}(t) \equiv [\partial^{m+n} f / \partial z_1^m \partial z_2^n]_t$; u_1 and u_2 are arbitrary fixed complex numbers, then

¹³ I.e., in the summation, both μ and ν take the values $1, 2, \bar{1}, \bar{2}$; $Z_{\bar{\nu}} = \bar{Z}_{\nu}$.

$$(3.5) \quad \lambda_B^{[3]} = \frac{1}{K(2-R) \sum T_{m\bar{n}} u_m \bar{u}_n}, \quad \Sigma \equiv \sum_{m, n=1}^2, \\ K \equiv K^{(B)}(t, \bar{t}), \quad T_{m\bar{n}} = T_{m\bar{n}}^{(B)}(t, \bar{t}).$$

Using (3.5), the relation $1/K = \lambda_B^1$ and the relation

$$ds^2 = \sum_{\mu, \nu=1}^2 T_{\mu\nu}^{(B)} u_\mu \bar{u}_\nu = \lambda_B^1 / \lambda_B^{(2)}$$

(see B. [2], p. 53), we have

$$(3.6) \quad 2 - R = \lambda_B^{(2)} / \lambda_B^{[3]}, \quad R = R(t).$$

Since $\lambda_B^{(2)}$ and $\lambda_B^{[3]}$ are positive, it follows from (3.6) that the curvature in an arbitrary analytic direction is less than 2. (B. [2], p. 54; F. [1]).

Let I and $A, t \in I \subset B \subset A$, be domains of comparison for the given domain B . Then from (3.6) and the monotonicity of the λ 's, we obtain

$$(3.7) \quad R \leq 2 - (\lambda_I^{(2)} / \lambda_A^{[3]}).$$

We shall show that the inequality (3.7) can be improved in certain cases if information about B of the following types is given: (1) Volume $B \leq V$, where V is a known number, (2) a few functions orthonormal over B , and (3) certain moments over B with weighting functions depending only upon I and A . We assume that

$$(3.8) \quad \text{Vol } A > V \geq \text{Vol } B, \quad I \subset B \subset A.$$

We shall show that this information leads to an improvement in (8.7) for some cases.

Define $\lambda_B^{[*3]} \equiv \lambda_B^{[3]}(t), t \in B$ to be the minimum of the integral (3.4) for functions $f \in \mathcal{L}^2(B)$ and normalized by the auxiliary conditions $f_{10}(t) = f_{01}(t) = 0, u_1^2 f_{20}(t) + 2u_1 u_2 f_{11}(t) + u_2^2 f_{02}(t) = 1$, where u_1 and u_2 are arbitrary fixed complex numbers.

Let

$$\{\alpha^{(\nu)}(z_1)\}, \nu = 1, 2, \dots, p-1, \{\beta^{(\mu)}(z_2)\},$$

$\mu = 1, 2, \dots, q-1$, be sets of functions satisfying

$$\int_B \alpha^{(\nu)} \cdot \overline{\alpha^{(k)}} d\omega = \delta_{\nu k}, \quad \int_B \beta^{(\mu)} \cdot \overline{\beta^{(k)}} d\omega = \delta_{\mu k},$$

$\delta_{\rho k} = 0, \rho \neq k, \delta_{kk} = 1$. We define $\lambda_{GB}^{[3]}, \lambda_{GB^{*p}}^{*01}, \lambda_{GB^{*q}}^{*01}$ for $B \subset G$ to be the minima of the integral

$$(3.9) \quad \int_G |f|^2 d\omega$$

for functions $f \in \mathcal{L}^2(G)$ and normalized by the respective sets of auxiliary conditions

1. $f(t) = f_{10}(t) = f_{01}(t) = 0, u_1^2 f_{20}(t) + 2u_1 u_2 f_{11}(t) + u_2^2 f_{02}(t) = 1, \int_B f d\omega = 0;$
2. $f_{10}(t) = 0, f_{01}(t) = 1, \int_B \bar{\alpha}^{(\nu)} \cdot f d\omega = 0, \nu = 1, 2, \dots, p-1;$
3. $f_{10}(t) = 0, f_{01}(t) = 1, \int_B \bar{\beta}^{(\mu)} \cdot f d\omega = 0, \mu = 1, 2, \dots, q-1.$

THEOREM 3. *The following relations hold*

$$(3.11) \quad (1/\lambda_B^{[3]}) = (1/\lambda_{BB}^{[3]}) + (F_B/\text{Vol } B) \cdot \{(1/\lambda_B^{[*3]}) - (1/\lambda_{BB}^{[3]})\}$$

where $F_B = \lambda_B^1 \lambda_B^{01} \lambda_B^{001} / \lambda_B^{*1} \lambda_B^{*01}$,

$$(3.12) \quad (1/\lambda_B^{*01}) = (1/\lambda_{BB* p}^{*01}) + \lambda_B^{*1} \left(\sum_{\nu=1}^{p-1} |\alpha_{z_1}^{(\nu)}|^2 \right) \{(1/\lambda_B^{* *1}) - (1/\lambda_{BB* p}^{*01})\}$$

$$(3.13) \quad (1/\lambda_B^{*01}) = (1/\lambda_{BB* * q}^{*01}) + \sum_{\mu=1}^{q-1} |\beta_{z_2}^{(\mu)}|^2.$$

Proof. Let $\{\psi^{(\sigma)}(z)\}$, $\sigma = 1, 2, \dots$, $\psi^{(1)} = (\text{Vol } B)^{-1/2}$, be a set of orthonormal functions complete for $\mathcal{L}^2(B)$. The minima $\lambda_B^{[3]}$, $\lambda_B^{[*3]}$ and $\lambda_{BB}^{[3]}$ are expressed in terms of u_1, u_2 and sums involving the functions $\{\psi^{(\sigma)}\}$ and their derivatives by taking in (1.5) values of $[m, n, X_1 X_2, X_3, X_4, \alpha_{1\nu}, \alpha_{2\nu}, \alpha_{3\nu}, \alpha_{4\nu}]$, respectively, as follows:

$$[1, 4, 0, 0, 0, 1, \psi^{(\nu)}(t), \psi_{10}^{(\nu)}(t), \psi_{01}^{(\nu)}(t), H],$$

$$[1, 3, 0, 0, 1, -, \psi_{10}^{(\nu)}(t), \psi_{01}^{(\nu)}(t), H, -],$$

and

$$[2, 4, 0, 0, 0, 1, \psi^{(\nu)}(t), \psi_{10}^{(\nu)}(t), \psi_{01}^{(\nu)}(t), H],$$

where

$$H = u_1^2 \psi_{20}^{(\nu)}(t) + 2u_1 u_2 \psi_{11}^{(\nu)}(t) + u_2^2 \psi_{02}^{(\nu)}(t).$$

The minima occurring in F_B (see (3.11)) are expressed in terms of u_1, u_2 and sums involving $\{\psi^{(\nu)}\}$ and their derivatives as indicated in the proof of Theorem 1. Combining the expressions for the minima so as to eliminate u_1, u_2 and the sums involving $\{\psi^{(\sigma)}\}$ and their derivatives, we obtain (3.11). Relations (3.12) and (3.13) are established in a similar manner. To express $\lambda_{BB* p}^{*01}$ and $\lambda_{BB* * q}^{*01}$ in terms of sums involving $\{\psi^{(\nu)}\}$ and their derivatives, we take in (1.5) values of $[m, n, X_1, X_2, \alpha_{1\nu}, \alpha_{2\nu}]$, respectively as follows: $[p, 2, 0, 1, \psi_{10}^{(\nu)}(t), \psi_{01}^{(\nu)}(t)]$, $[q, 2, 0, 1, \psi_{10}^{(\nu)}(t), \psi_{01}^{(\nu)}(t)]$. To find similar expressions for the other minima in (3.12) and (3.13), see the proof of Theorem 2.

THEOREM 4. *Let B be a given domain in the (z_1, z_2) -space having interior and exterior domains of comparison I and A . Let the point $t = (t_1, t_2) \in I$ and R be the Riemann curvature at t in the analytic direction (u_1, u_2) , of the Bergman metric (3.1) where $K \equiv K^{(B)}(z, \bar{z})$ is the kernel function of B . Let $\{\varphi^{(\sigma)}(z_1, z_2)\}$, $\sigma = 1, 2, \dots, r-1$, $\{\alpha^{(\nu)}(z_1)\}$, $\nu = 1, 2, \dots, p-1$, $\{\beta^{(\mu)}(z_2)\}$, $\mu = 1, 2, \dots, q-1$, be three sets of functions possessing the properties*

$$(3.14) \quad \int_B \varphi^{(\sigma)} \overline{\varphi^{(k)}} d\omega = \delta_{\sigma k}, \quad \int_B \alpha^{(\nu)} \overline{\alpha^{(k)}} d\omega = \delta_{\nu k}, \quad \int_B \beta^{(\mu)} \overline{\beta^{(k)}} d\omega = \delta_{\mu k},$$

$$\delta_{\rho k} = 0, \rho \neq k, \delta_{kk} = 1.$$

Then

$$(3.15) \quad R \leq 2 - \lambda_I^{(2)} L$$

where

$$L = \max \{ (1/\lambda_A^{[3]}), (1/\lambda_{AB}^{[3]}) + (F/V)[1/\lambda_A^{[*3]}] - (1/\lambda_{AB}^{[3]}) \} ,$$

$$V \geq \text{Vol } B, F = \lambda_I^1 \lambda_I^{01} \lambda_I^{001} F_1 F_2 ,$$

$$F_1 = \max \left\{ (1/\lambda_A^{*1}), (1/\lambda_{ABr}^{*1}) + \sum_{\sigma=1}^{r-1} |\varphi_{z_2}^{(\sigma)}|^2 \right\} ,$$

$$F_2 = \max \left\{ (1/\lambda_A^{*01}), (1/\lambda_{AB*}^{*01}) + \sum_{\mu=1}^{q-1} |\beta_{z_2}^{(\mu)}|^2 , \right.$$

$$\left. (1/\lambda_{AB*}^{*01}) + \lambda_I^{*1} \left(\sum_{\nu=1}^{p-1} |\alpha_{z_1}^{(\nu)}|^2 \right) [(1/\lambda_A^{*1}) - (1/\lambda_{AB*}^{*01})] \right\} ,$$

and where the λ 's are solutions of minimum problems depending upon the domains indicated in the subscripts, and where the λ 's bearing multiple subscripts depend only upon the kernel function of A , the first few derivatives of the kernel function of A , a finite number of integrals over B with weighting functions depending only upon A , and in the case of $\lambda_{AB}^{[3]}$ also upon u_1, u_2 .

Proof. From the definitions it is clear that

$$(3.16) \quad \lambda_B^{[*3]} \leq \lambda_B^{[3]} \leq \lambda_{BB}^{[3]}$$

Hence (3.11) implies

$$(3.17) \quad F_B / \text{Vol } B \equiv (\lambda_B^1 \cdot \lambda_B^{01} \cdot \lambda_B^{001}) / (\lambda_B^{*1} \cdot \lambda_B^{*01} \cdot \text{Vol } B) \leq 1 .$$

Using the monotonicity of the λ 's, (3.11)–(3.13) imply

$$(3.18) \quad (1/\lambda_B^{[3]}) \geq (1/\lambda_{AB}^{[3]}) + (F_1/V)[(1/\lambda_A^{[*3]}) - (1/\lambda_{AB}^{[3]})] .$$

(See details of the proof of Theorem 2 in S. [1], which is the lemma in § 1 of this paper.)

where F_1 is such that $F_B \geq F_1$,

$$(3.19) \quad (1/\lambda_B^{*01}) \geq (1/\lambda_{AB**p}^{*01}) + \lambda_I^{*1} \left(\sum_{\nu=1}^{p-1} |\alpha_{z_1}^{(\nu)}|^2 \right) [(1/\lambda_A^{*1}) - (1/\lambda_{AB**p}^{*01})]$$

$$(3.20) \quad (1/\lambda_B^{*01}) \geq (1/\lambda_{AB**q}^{*01}) + \sum_{\mu=1}^{q-1} |\beta_{z_2}^{(\mu)}|^2.$$

From (3.6) it follows that

$$(3.21) \quad R \leq 2 - \lambda_I^{(2)} \cdot L_1$$

where L_1 is a lower bound for $1/\lambda_A^{[3]}$. Combining (3.21), (3.18), (3.19), (3.20), (2.13), and (2.14), we obtain (3.15).

To express the minima bearing multiple subscripts in terms of the quantities mentioned in the theorem, we proceed as in the proof of Theorem 3.

EXAMPLE. If $L = 1/\lambda_A^{[3]}$, then (3.15) reduces to (3.7). To show that there are cases in which (3.15) is an improvement over (3.7), we proceed as follows: Let the given domain B contain the origin and have as interior an exterior domain of comparison at the origin the hyper-spheres

$$I: \sum_{k=1}^2 |z_k - \varepsilon m|^2 < m^2$$

and

$$A: \sum_{k=1}^2 |z_k - \varepsilon M|^2 < M^2, \quad \varepsilon^2 < 1/2$$

respectively.

In constructing examples we must always take m and M so related that $I \subset A$. We note that $I \subset A$ if $m < M$.

Define F_{IA} to be

$$F_{IA} = (\lambda_I^1 \lambda_I^{01} \lambda_I^{001}) / (\lambda_A^{*1} \lambda_A^{*01})$$

and to facilitate computing take $u_1 = 1$, $u_2 = yi$, y real. Using formula (1.5) (see also B. [2] p. 43) and the fact that the kernel functions of I and A are

$$K^{(I)}(z, \bar{z}) = 2m^2 / \left\{ \pi^2 \left[m^2 - \sum_{k=1}^2 (z_k - \varepsilon m)(\bar{z}_k - \varepsilon m) \right] \right\}^3,$$

$$K^{(A)}(z, \bar{z}) = 2M^2 / \left\{ \pi^2 \left[M^2 - \sum_{k=1}^2 (z_k - \varepsilon M)(\bar{z}_k - \varepsilon M) \right] \right\}^3$$

respectively, we compute the λ 's in terms of ε , m , M , y , and obtain that $(F_{IA}/V)(1/\lambda_A^{[3]})$ and $(1/\lambda_A^{[*3]})$ can be written in the forms

$$(F_{IA}/V)(1/\lambda_A^{[3]}) = \frac{1}{\rho_I} \frac{1}{\rho^{12}} \alpha \sum_{\nu=0}^7 U_{\nu}(y) \cdot \varepsilon^{2\nu},$$

$$(1/\lambda_A^{[3]}) = \alpha \sum_{\mu=0}^4 W_\mu(y) \varepsilon^{2\mu}$$

where

$$\begin{aligned} \rho_I &= V / \left(\frac{\pi^2 m^4}{2} \right) = V / (\text{Vol } I) \geq 1, \\ \alpha &= \left\{ \frac{48}{\pi^2 M^8 (1 - 2\varepsilon^2)^8} \right\}, \quad \rho = M/m > 1, \end{aligned}$$

and where the coefficients $U_\nu(y)$ and $W_\mu(y)$ are functions of y only.

Computation gives that $U_0(y) = (y + 1)^2 = W_0(y)$, and

$$U_1(y) = 4(y^4 - 2y^2 - 3) = W_1(y).$$

If we let $\eta = (1/\sqrt{2}) - \varepsilon$ we obtain

$$\sum_{\nu=2}^7 U_\nu(y) \varepsilon^{2\nu-4} = b_0(y) + g_1(y)\eta + 0(\eta^2)$$

and

$$\sum_{\mu=2}^4 W_\mu(y) \varepsilon^{2\mu-4} = b_0(y) + g_2(y)\eta + 0(\eta^2),$$

where $b_0(y) = 4(1 - 2y^2 - 3y^4)$ and where $g_1(y) > g_2(y)$ for $|y|$ sufficiently small.

To obtain our desired example, we first choose y sufficiently near zero that $g_1(y) > g_2(y)$. Then we take ε positive and near to $1/\sqrt{2}$, m near to M , and B such that $I \subset B \subset A$ and such that $V/(\text{Vol } I)$ is near 1 so that

$$(F_{IA}/V)(1/\lambda_A^{[*3]}) > (1/\lambda_A^{[3]}).$$

This then gives the desired example, for we have

$$L \geq (F/V)(1/\lambda_A^{[*3]}) \geq (F_{IA}/V)(1/\lambda_A^{[*3]}) > (1/\lambda_A^{[3]}),$$

so that (3.15) gives a better bound for R at the origin in the direction $(1, yi)$, y real and $|y|$ small, than does (3.7).

BIBLIOGRAPHY

S. Bergman,

1. *The method of the minimum integral and the analytic continuation of functions of two complex variables*, Proc. Mat. Acad. Sci., U. S. A., **27** (1941), 328–332.
2. *Sur les fonctions orthogonales de plusieurs variables complexes avec les applications à la théorie des fonctions analytiques*. Mémorial des Sciences Mathématiques, **106** (1947) Paris.
3. *Sur la fonction-noyau d'un domaine et ses applications dans la théorie des transformations pseudo-conformes*, Mémorial des Sciences Mathématiques, **108** (1948) Paris.
4. *Partial differential equations*, mimeographed notes, Brown University, Providence, R. I. (1941).

B. Fuchs,

1. *Über geodätische Mannigfaltigkeiten einer bei pseudo-konformen Abbildungen invarianten Riemannschen Geometrie*, Recueil Math., nouv. ser., **2** (44), 567–594.

S. Kobayashi,

1. *Geometry of bounded domain*, Trans. of Amer. Math. Soc., **92**, (1959), 267 ff.

J. M. Stark,

1. *On Distortion in Pseudo-Conformal Mapping*, Pacific J. Math. **6** (1956) 565–582.

LAMAR STATE COLLEGE OF TECHNOLOGY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DAVID GILBARG

Stanford University
Stanford, California

F. H. BROWNELL

University of Washington
Seattle 5, Washington

A. L. WHITEMAN

University of Southern California
Los Angeles 7, California

L. J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

T. M. CHERRY

D. DERRY

E. HEWITT

A. HORN

L. NACHBIN

M. OHTSUKA

H. L. ROYDEN

M. M. SCHIFFER

E. SPANIER

E. G. STRAUS

F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA

NEW MEXICO STATE UNIVERSITY

OREGON STATE COLLEGE

UNIVERSITY OF OREGON

OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE COLLEGE

UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

CALIFORNIA RESEARCH CORPORATION

HUGHES AIRCRAFT COMPANY

SPACE TECHNOLOGY LABORATORIES

NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 2120 Oxford Street, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 10, No. 3

November, 1960

Glen Earl Baxter, <i>An analytic problem whose solution follows from a simple algebraic identity</i>	731
Leonard D. Berkovitz and Melvin Dresher, <i>A multimove infinite game with linear payoff</i>	743
Earl Robert Berkson, <i>Sequel to a paper of A. E. Taylor</i>	767
Gerald Berman and Robert Jerome Silverman, <i>Embedding of algebraic systems</i>	777
Peter Crawley, <i>Lattices whose congruences form a boolean algebra</i>	787
Robert E. Edwards, <i>Integral bases in inductive limit spaces</i>	797
Daniel T. Finkbeiner, II, <i>Irreducible congruence relations on lattices</i>	813
William James Firey, <i>Isoperimetric ratios of Reuleaux polygons</i>	823
Delbert Ray Fulkerson, <i>Zero-one matrices with zero trace</i>	831
Leon W. Green, <i>A sphere characterization related to Blaschke's conjecture</i>	837
Israel (Yitzchak) Nathan Herstein and Erwin Kleinfeld, <i>Lie mappings in characteristic 2</i>	843
Charles Ray Hobby, <i>A characteristic subgroup of a p-group</i>	853
R. K. Juberg, <i>On the Dirichlet problem for certain higher order parabolic equations</i>	859
Melvin Katz, <i>Infinitely repeatable games</i>	879
Emma Lehmer, <i>On Jacobi functions</i>	887
D. H. Lehmer, <i>Power character matrices</i>	895
Henry B. Mann, <i>A refinement of the fundamental theorem on the density of the sum of two sets of integers</i>	909
Marvin David Marcus and Roy Westwick, <i>Linear maps on skew symmetric matrices: the invariance of elementary symmetric functions</i>	917
Richard Dean Mayer and Richard Scott Pierce, <i>Boolean algebras with ordered bases</i>	925
Trevor James McMinn, <i>On the line segments of a convex surface in E_3</i>	943
Frank Albert Raymond, <i>The end point compactification of manifolds</i>	947
Edgar Reich and S. E. Warschawski, <i>On canonical conformal maps of regions of arbitrary connectivity</i>	965
Marvin Rosenblum, <i>The absolute continuity of Toeplitz's matrices</i>	987
Lee Albert Rubel, <i>Maximal means and Tauberian theorems</i>	997
Helmut Heinrich Schaefer, <i>Some spectral properties of positive linear operators</i>	1009
Jeremiah Milton Stark, <i>Minimum problems in the theory of pseudo-conformal transformations and their application to estimation of the curvature of the invariant metric</i>	1021
Robert Steinberg, <i>The simplicity of certain groups</i>	1039
Hisahiro Tamano, <i>On paracompactness</i>	1043
Angus E. Taylor, <i>Mittag-Leffler expansions and spectral theory</i>	1049
Marion Franklin Tinsley, <i>Permanents of cyclic matrices</i>	1067
Charles J. Titus, <i>A theory of normal curves and some applications</i>	1083
Charles R. B. Wright, <i>On groups of exponent four with generators of order two</i>	1097