

Pacific Journal of Mathematics

MULTIPLICATION ON CLASSES OF PSEUDO-ANALYTIC FUNCTIONS

JOHN WILLIAM JEWETT

MULTIPLICATION ON CLASSES OF PSUEDO-ANALYTIC FUNCTIONS

JOHN JEWETT

Lipman Bers [1, 2] has formulated a theory of solutions of linear elliptic partial differential equations in terms of classes of psuedo-analytic functions on a plane domain D . The theory for each class of psuedo-analytic functions is based on the notion of a generating pair of Holder continuous complex valued functions F and G defined on D and satisfying $\Im[\overline{F(z)}G(z)] > 0$ in D .

If w is any function defined on D , then there exist two real valued functions ϕ and ψ such that w can be written uniquely as

$$(1) \quad w(z) = \phi(z)F(z) + \psi(z)G(z).$$

A function w defined on D is said to be (F, G) -psuedo-analytic (of the first kind) if a certain generalized derivative exists or equivalently if the equations

$$(2) \quad \begin{aligned} \phi_x F_1 - \phi_y F_2 + \psi_x G_1 - \psi_y G_2 &= 0 \\ \phi_y F_1 + \phi_x F_2 + \psi_y G_1 + \psi_x G_2 &= 0 \end{aligned}$$

are satisfied in D , where the subscripts x and y refer to partial derivatives with respect to x and y and the subscripts 1 and 2 refer to the real and imaginary parts of the functions F and G . If $F = 1$ and $G = i$, these equations reduce to the Cauchy-Riemann equations.

Given a generating pair (F, G) let B denote the class of all functions which are (F, G) -psuedo-analytic. If $F = 1$ and $G = i$, then B is the class of analytic functions on D , which will be referred to in this paper as A .

Any B has many of the properties of the ring of analytic functions. In particular very close analogues of the identity theorem, the Cauchy theorem, the Cauchy integral formula, the standard convergence theorems, and power series expansions have been proved.

With each class B is associated a class B' of psuedo-analytic functions of the second kind. This association is made by a mapping γ of B into B' defined by

$$\gamma(\phi F + \psi G) = \phi + i\psi.$$

On the class A of analytic functions this mapping is clearly the identity.

Each class B is a vector space with the usual definition of addition

of functions and multiplication by scalars and η is a vector space isomorphism of B onto B' . The class A is a ring under the usual pointwise multiplication of functions. Since the classes of psuedo analytic functions each bear such marked resemblances to the class A of analytic functions, the question arises as to whether there exist for other classes appropriate generalizations of the ordinary multiplications of function. We shall prove that if such a multiplication bears a certain slight resemblance to the point wise multiplication, then B is multiplicatively isomorphic to A under the mapping η and conversely.

We denote the ordinary multiplication of functions by juxtaposition. Let m denote any mapping from $B \times B$ to the set of all functions from D to the plane. In particular let m_x be the mapping defined as follows: if $w = \phi F + \psi G$ and $w' = \phi' F + \psi' G$, let

$$m_x(w, w') = (\phi\phi' - \psi\psi')F + (\phi\psi' + \psi'\phi)G .$$

THEOREM. *Let B be a system of psuedo-analytic functions on the plane domain D and let m be a multiplication on B (any mapping from $B \times B$ to the set of all functions from D to the plane). Let m be associative and bilinear with respect to addition in B . Then a necessary and sufficient condition for the mapping η to be a multiplicative isomorphism of B onto the ring A of all analytic functions on D is that there exists a non-constant w in B such that $m(w, G) = m_x(w, G)$ and $m_x(w, G) \in B$.*

The proof of this theorem will be preceded by a lemma.

LEMMA. *Suppose that for all w and w' in B , $m(w, w') = m_x(w, w')$. Then the mapping η defined above is an isomorphism of B onto the ring A of analytic functions on D if and only if $F_1 = G_2$ and $F_2 = -G_1$.*

Proof of Lemma. A simple calculation shows that η is an isomorphism of B onto B' if and only if $m = m_x$. So the condition concerning isomorphism in the lemma is that $B' = A$.

By adding and subtracting terms involving ψ the system (2) is seen to be equivalent to

$$(3) \quad \begin{aligned} F_1(\phi_x - \phi_y) - F_2(\phi_y + \psi_x) + \psi_y(F_1 - G_2) + \psi_x(F_2 + G_1) &= 0 \\ F_1(\phi_y + \psi_x) + F_2(\phi_x - \psi_y) + \psi_y(F_2 + G_1) - \psi_x(F_1 - G_2) &= 0 . \end{aligned}$$

First suppose $F_1 = G_2$ and $F_2 = -G_1$. Then this system becomes

$$(4) \quad \begin{aligned} F_1(\phi_x - \psi_y) - F_2(\phi_y + \psi_x) &= 0 \\ F_1(\phi_y + \psi_x) + F_2(\phi_x - \psi_y) &= 0 . \end{aligned}$$

It is clear that if $w^* = \phi + i\psi$ is analytic, then $\eta^{-1}(w^*)$ satisfies the system (4). Therefore $A \subset B'$

Suppose then that B' contains a function $w = \phi + i\psi$ which at some point z of D does not satisfy the Cauchy-Riemann equations. For this point the system (4) is a system of homogeneous algebraic equations with a non zero determinant whose value is

$$[\phi_x(z) - \psi_y(z)]^2 + [\phi_y(z) + \psi_x(z)]^2 .$$

Hence the only solution at z is the zero solution and thus

$$\Im[\overline{F(z)}G(z)] = F_1^2(z) + F_2^2(z) = 0$$

which contradicts the definition of generating pair. Thus $B' \subset A$ and we have proved that $A = B'$.

Conversely suppose that η is an isomorphism onto A so that $A = B'$. Let $w^* = \phi + i\psi$ be a non constant analytic function in B' . Then the system (3) becomes for this w^*

$$(5) \quad \begin{aligned} \psi_y(F_1 - G_2) + \psi_x(F_2 + G_1) &= 0 \\ \psi_y(F_2 + G_1) - \psi_x(F_1 - G_2) &= 0 . \end{aligned}$$

If for some z the equations $F_1(z) = G_2(z)$ and $F_2(z) = -G_1(z)$ do not both hold, then the determinant of this system is non-zero at z and hence by continuity of F and G the determinant is non-zero in some neighborhood of z and hence $\psi_x = \psi_y = 0$ on this neighborhood. By the identity theorem for harmonic functions ψ_x and ψ_y must then be zero everywhere so that ψ is constant. A similar argument demonstrates the constancy of ϕ so that w^* is constant contrary to assumption. This completes the proof of the lemma.

Proof of Theorem. Suppose first that η is a multiplicative isomorphism of B onto the ring A of analytic functions in D . Then as before m is identically equal to m_p so that for $w = \phi F + \psi G \in B$ we have $m(w, G) = -\psi F + \phi G$. Substituting this function for $\phi F + \psi G$ in the system (1) yields that $m(w, G)$ is in B if and only if

$$(6) \quad \begin{aligned} -\psi_x F_1 + \psi_y F_2 + \phi_x G_1 - \phi_y G_2 &= 0 \\ -\psi_y F_1 - \psi_x F_2 + \phi_y G_1 + \phi_x G_2 &= 0 . \end{aligned}$$

By the lemma $F_1 = G_2$ and $F_2 = -G_1$. Using this to substitute for the G 's in the system (6) we obtain the system (4) and this system must be satisfied because $\phi + i\psi$ is analytic. Thus if w is in B then so is $m(w, G)$ and the condition of the theorem is necessary.

Conversely suppose that there exists a non-constant w in B such that $m(w, G) = m_p(w, G) = -\psi F + \phi G$ and this function is in B . Then ϕ and ψ satisfy both (1) and (6) and since w is non-constant there must exist a z such that this system of four equations has a non-zero solution, i.e., the determinant of this system must be zero.

The determinant of this system is

$$(7) \quad [(F_1 - G_1)^2 + (F_2 + G_1)^2][(F_1 + G_2)^2 + (F_2 - G_1)^2].$$

Now $\Im(\bar{F}G) = F_1G_2 - F_2G_1$ which must be everywhere positive since F and G form a generating pair. If the second factor of (7) is zero, then it follows that

$$\Im(\bar{F}G) = -F_1^2 - F_2^2 < 0.$$

Hence the first factor must be zero and the lemma implies that η is an isomorphism of B onto A .

BIBLIOGRAPHY

1. Lipman Bers, *Theory of Pseudo-analytic functions*, Mimeographed Lecture Notes, New York University, 1953.
2. ———, *Local Theory of Pseudo-analytic functions*, Lectures on Functions of a Complex Variable, Ann Arbor, Michigan, 1955, pp. 213-244.

UNIVERSITY OF GEORGIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DAVID GILBARG
Stanford University
Stanford, California

F. H. BROWNELL
University of Washington
Seattle 5, Washington

A. L. WHITEMAN
University of Southern California
Los Angeles 7, California

L. J. PAIGE
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH
T. M. CHERRY
D. DERRY

E. HEWITT
A. HORN
L. NACHBIN

M. OHTSUKA
H. L. ROYDEN
M. M. SCHIFFER

E. SPANIER
E. G. STRAUS
F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON
* * *

AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
HUGHES AIRCRAFT COMPANY
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 2120 Oxford Street, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

M. Altman, <i>An optimum cubically convergent iterative method of inverting a linear bounded operator in Hilbert space</i>	1107
Nesmith Cornett Ankeny, <i>Criterion for rth power residuacity</i>	1115
Julius Rubin Blum and David Lee Hanson, <i>On invariant probability measures I</i>	1125
Frank Featherstone Bonsall, <i>Positive operators compact in an auxiliary topology</i>	1131
Billy Joe Boyer, <i>Summability of derived conjugate series</i>	1139
Delmar L. Boyer, <i>A note on a problem of Fuchs</i>	1147
Hans-Joachim Bremermann, <i>The envelopes of holomorphy of tube domains in infinite dimensional Banach spaces</i>	1149
Andrew Michael Bruckner, <i>Minimal superadditive extensions of superadditive functions</i>	1155
Billy Finney Bryant, <i>On expansive homeomorphisms</i>	1163
Jean W. Butler, <i>On complete and independent sets of operations in finite algebras</i>	1169
Lucien Le Cam, <i>An approximation theorem for the Poisson binomial distribution</i>	1181
Paul Civin, <i>Involutions on locally compact rings</i>	1199
Earl A. Coddington, <i>Normal extensions of formally normal operators</i>	1203
Jacob Feldman, <i>Some classes of equivalent Gaussian processes on an interval</i>	1211
Shaul Foguel, <i>Weak and strong convergence for Markov processes</i>	1221
Martin Fox, <i>Some zero sum two-person games with moves in the unit interval</i>	1235
Robert Pertsch Gilbert, <i>Singularities of three-dimensional harmonic functions</i>	1243
Branko Grünbaum, <i>Partitions of mass-distributions and of convex bodies by hyperplanes</i>	1257
Sidney Morris Harmon, <i>Regular covering surfaces of Riemann surfaces</i>	1263
Edwin Hewitt and Herbert S. Zuckerman, <i>The multiplicative semigroup of integers modulo m</i>	1291
Paul Daniel Hill, <i>Relation of a direct limit group to associated vector groups</i>	1309
Calvin Virgil Holmes, <i>Commutator groups of monomial groups</i>	1313
James Fredrik Jakobsen and W. R. Utz, <i>The non-existence of expansive homeomorphisms on a closed 2-cell</i>	1319
John William Jewett, <i>Multiplication on classes of pseudo-analytic functions</i>	1323
Helmut Klingen, <i>Analytic automorphisms of bounded symmetric complex domains</i>	1327
Robert Jacob Koch, <i>Ordered semigroups in partially ordered semigroups</i>	1333
Marvin David Marcus and N. A. Khan, <i>On a commutator result of Taussky and Zassenhaus</i>	1337
John Glen Marica and Steve Jerome Bryant, <i>Unary algebras</i>	1347
Edward Peter Merkes and W. T. Scott, <i>On univalence of a continued fraction</i>	1361
Shu-Teh Chen Moy, <i>Asymptotic properties of derivatives of stationary measures</i>	1371
John William Neuberger, <i>Concerning boundary value problems</i>	1385
Edward C. Posner, <i>Integral closure of differential rings</i>	1393
Marian Reichaw-Reichbach, <i>Some theorems on mappings onto</i>	1397
Marvin Rosenblum and Harold Widom, <i>Two extremal problems</i>	1409
Morton Lincoln Slater and Herbert S. Wilf, <i>A class of linear differential-difference equations</i>	1419
Charles Robson Storey, Jr., <i>The structure of threads</i>	1429
J. François Treves, <i>An estimate for differential polynomials in $\partial/\partial z_1, \dots, \partial/\partial z_n$</i>	1447
J. D. Weston, <i>On the representation of operators by convolutions integrals</i>	1453
James Victor Whittaker, <i>Normal subgroups of some homeomorphism groups</i>	1469