

Pacific Journal of Mathematics

TWO EXTREMAL PROBLEMS

MARVIN ROSENBLUM AND HAROLD WIDOM

TWO EXTREMAL PROBLEMS

MARVIN ROSENBLUM AND HAROLD WIDOM

1. Introduction. Let \mathcal{S}_0 be the class of all complex trigonometric polynomials P of the form $P_0 + P_1 e^{i\phi} + P_2 e^{2i\phi} + \dots$. Let σ and μ be, respectively normalized Lebesgue measure and any finite non-negative Borel measure on the real interval $(-\pi, \pi]$. Suppose $\mu = \mu_A + \mu_S$, with $d\mu_A(\phi) = f(\phi)d\sigma(\phi)$, is the Lebesgue decomposition of μ into absolutely continuous and singular measures. In this note we shall be concerned with two generalizations of the problem Q_0 : Find

$$I_0(\mu) = \inf_{P \in \mathcal{S}_0} \left[\int |1 + e^{i\phi} P(e^{i\phi})|^2 d\mu(\phi) \right]^{\frac{1}{2}}.$$

Q_0 was solved by Szegö for the case $\mu = \mu_A$ and in general by M. G. Krein and Kolmogorov. They showed that $I_0(\mu) = \exp \frac{1}{2} \int \log f d\sigma$ if $\log f$ is integrable and $I_0(\mu) = 0$ otherwise. (See [3], pp. 44, 231.)

We shall consider:

Problem Q_1 : Suppose $\int |g|^2 d\mu < \infty$. Find

$$I_1(g, \mu) = \inf_{P \in \mathcal{S}_0} \left[\int |g + P|^2 d\mu \right]^{\frac{1}{2}},$$

and

Problem Q_2 : Suppose $\int |h| d\sigma < \infty$. Find

$$I_2(h, \mu) = \sup_{P \in \mathcal{S}_0} \left\{ \left| \int Ph d\sigma \right| / \left[\int |P|^2 d\mu \right]^{\frac{1}{2}} \right\}.$$

Clearly $I_1(e^{-i\phi}, \mu) = I_0(\mu)$. Also

$$[I_2(1, \mu)]^{-1} = \inf_{P \in \mathcal{S}_0} \left\{ \left[\int |P|^2 d\mu \right]^{\frac{1}{2}} / \left| \int Pd\sigma \right| \right\} = I_0(\mu),$$

so Q_0 is a particularization of both Q_1 and Q_2 . There are other special cases of Q_1 and Q_2 that can be found in the work of Szegö [5] and Grenander and Szegö [3]. Of particular interest are the following:

(i) Let $g(\phi) = e^{-i(k+1)\phi}$, where k is a positive integer. Then Q_1 is the problem of linear prediction k units ahead of time ([3], p. 184).

(ii) Let $h(\phi) = 1/(1 - \alpha e^{-i\phi})$, $|\alpha| < 1$. Then

$$I_2(h, \mu) = \sup_{P \in \mathcal{S}_0} \left\{ |P(\alpha)| / \left[\int |P|^2 d\mu \right]^{\frac{1}{2}} \right\}.$$

Received October 20, 1959, in revised form February 2, 1960. This work was done while both authors held National Science Foundation postdoctoral fellowships.

See [3], p. 48.

Throughout we shall indulge in the following notational conveniences: We shall write $I_1(g, f)$ and $I_2(h, f)$ for $I_1(g, \mu_A)$ and $I_2(h, \mu_A)$ respectively, and, in certain contexts, consider two functions identical that are equal everywhere except for a set of Lebesgue measure zero.

We have divided this note into six sections. First we indicate an interesting duality between $I_1(e^{-i\phi}g(\phi), f)$ and $I_2(g, 1/f)$ that relates the problems Q_1 and Q_2 under certain restrictive hypotheses. In section three we fashion the theory that will handle Q_1 and Q_2 . This is the solution of a Riemann-Hilbert problem (which we call problem Q_3), which is applied in §§ 4, 5 and 6 to Q_1 and Q_2 .

2. Duality of I_1 and I_2 . This will fall out of the following Banach space lemma:

Let \mathcal{P}_0 be a subspace of a Banach space \mathcal{L} and let \mathcal{P}_0^\perp be the annihilator of \mathcal{P}_0 in the dual space \mathcal{L}^* . If $g \in \mathcal{L}$, then

$$\inf \{ \|g + P\| : P \in \mathcal{P}_0 \} = \sup \{ |l(g)| : l \in \mathcal{P}_0^\perp, \|l\| \leq 1 \}.$$

For a proof see Bonsall [2].

THEOREM 1. Suppose f and $1/f$ are in $L^1(-\pi, \pi)$ and $\int |g|^2 f d\sigma < \infty$. Then

$$I_1(e^{-i\phi}g(\phi), f) = I_2(g, 1/f).$$

Sketch of proof. By the above lemma

$$I_1(e^{-i\phi}g(\phi), f) = \sup \left\{ \left| \int e^{-i\phi}g(\phi)h(\phi)f(\phi)d\sigma \right| / \left[\int |h|^2 f d\sigma \right]^{\frac{1}{2}} \right\},$$

where the supremum is taken over all h such that $\int e^{in\phi}h(\phi)f(\phi)d\sigma = 0$ for $n = 0, 1, 2, \dots$. Through the substitution $e^{-i\phi}hf = P$ it follows that

$$I_1(e^{-i\phi}g(\phi), f) = \sup \left\{ \left| \int P f d\sigma \right| / \left[\int |P|^2 \frac{1}{f} d\sigma \right]^{\frac{1}{2}} \right\},$$

where now the supremum is taken over all P such that $\int e^{in\phi}P(\phi)d\sigma = 0$ for $n = 1, 2, \dots$. It can be shown that it is sufficient merely to consider suprema for $P \in \mathcal{P}_0$, which proves the theorem.

The restrictive condition $1/f \in L^1(-\pi, \pi)$ seems essential to the formulation of the preceding duality relation, but at least this relation indicates that there exist close tie-ins between Q_1 and Q_2 . We shall solve a Riemann-Hilbert problem for the unit circle that, when applied to Q_1 and Q_2 , solves both.

3. **The Riemann-Hilbert problem Q_3 .** Let f be a non-negative function in $L^1 = L^1(-\pi, \pi)$, and suppose that \mathcal{P} is the closure of \mathcal{S}_0 in the Hilbert space $L^2(f)$ of functions square integrable with respect to the measure $f d\sigma$. Thus, for example, \mathcal{P} in $L^2(1) = L^2$ can be identified with the Hardy space H^2 . The problem Q_3 is:

Given $k \in L^1$, find functions $P \in \mathcal{P}$ and q satisfying

$$(1) \quad Pf = k + q, \quad \text{and}$$

$$(2) \quad \int qe^{-in\phi} d\sigma = 0, \quad n = 0, 1, \dots$$

(Note that since $\int |P|^2 f d\sigma < \infty$, we have $Pf \in L^1$ and so $q = Pf - k \in L^1$.)

We first list some prefatory material. We associate with any non-negative $f \in L^1$ such that $\log f \in L^1$ the analytic functions

$$(3) \quad \begin{aligned} F^+(z) &= \exp \frac{1}{2} \int \frac{e^{i\phi} + z}{e^{i\phi} - z} \log f(\phi) d\sigma(\phi), \quad |z| < 1, \\ F^-(z) &= \exp \frac{1}{2} \int \frac{z + e^{i\phi}}{z - e^{i\phi}} \log f(\phi) d\sigma(\phi), \quad |z| > 1. \end{aligned}$$

F^+ and F^- belong to H^2 and K^2 respectively, and $\overline{F^-(z)} = F^+(1/\bar{z})$ if $|z| > 1$. (A function $F(z)$ is said to belong to K^2 if $F(1/z)$ belongs to H^2 .) Since the boundary functions in H^2 and K^2 exist in mean square, we can define

$$(4) \quad \begin{aligned} f^+(\phi) &= \lim_{r \rightarrow 1^-} F^+(re^{i\phi}), \\ f^-(\phi) &= \lim_{r \rightarrow 1^+} F^-(re^{i\phi}). \end{aligned}$$

These functions satisfy

$$(5) \quad f(\phi) = f^-(\phi)f^+(\phi) = |f^+(\phi)|^2 = |f^-(\phi)|^2.$$

For any non-negative $f \in L^1$ and $\varepsilon > 0$ we define $F_\varepsilon^\pm(z)$, $f_\varepsilon^\pm(\phi)$ by (3) and (4) with f replaced by $f_\varepsilon = f + \varepsilon$. Here we need not assume that $\log f \in L^1$. Note that since $f + \varepsilon \geq \varepsilon > 0$, we have $1/F_\varepsilon^+ \in H^\infty$ and $1/F_\varepsilon^- \in K^\infty$. Moreover $|f_\varepsilon^+(\phi)|^2 = f(\phi) + \varepsilon$, so $|f_\varepsilon^-(\phi)| = |f_\varepsilon^+(\phi)| \geq [f(\phi)]^{1/2}$.

Next we define an operator $(\)_+$ as follows. Its domain D consists of all L^1 functions k with Fourier series $\sum_{-\infty}^\infty c_n e^{in\phi}$ such that $\sum_0^\infty |c_n|^2 < \infty$, and k_+ is the function with Fourier series $\sum_0^\infty c_n e^{in\phi}$. We define the operator $(\)_-$ by $k_- = k - k_+$. Notice that $k_+ \in H^2$ and $k_- \in K^1$ with $\int k_- d\sigma = 0$.

Our discussion of Q_3 proceeds in the following order. First we prove uniqueness. Then we solve Q_3 in certain special cases (these being sufficient, it will turn out, to handle Q_1), and finally find the solution in

the general case.

We are indebted to the referee for the proof of the next lemma.

LEMMA 2. Q_3 has at most one solution.

Proof. Suppose $Pf = q$ where $P \in \mathcal{S}$ and q satisfies (2). Then P is orthogonal, in $L^2(f)$, to all exponentials $e^{in\phi}$ ($n \geq 0$). Since P belongs to the closed manifold \mathcal{S} spanned by these exponentials we conclude $P = 0$.

One can formally solve Q_3 by means of the usual factorization methods (see [4], for example). Write $f = f^+f^-$, so $Pf = k + q$ implies

$$Pf^+ = \frac{k}{f^-} + \frac{q}{f^-}.$$

Applying $(\)_+$ to both sides we obtain $Pf^+ = (k/f^-)_+$, $P = (1/f^+)(k/f^-)_+$. The following theorem justifies this procedure in certain cases.

THEOREM 3. (i) Suppose $\log f \in L^1$ and $k/f^- \in D$. Then Q_3 has the solution

$$(6) \quad P = \frac{1}{f^+} \left(\frac{k}{f^-} \right)_+ \quad q = -f^- \left(\frac{k}{f^-} \right)_-.$$

(ii) Suppose $\log f \notin L^1$ and $k^2/f \in L^1$. Then Q_3 has the solution

$$P = \frac{k}{f} \quad q = 0.$$

Proof. (i) Let $\varepsilon > 0$. Since the function f^+ is outer, it follows from a theorem of Beurling [1] that there exists a $P_0 \in \mathcal{S}_0$ such that

$$\int \left| \left(\frac{k}{f^-} \right)_+ - P_0 f^+ \right|^2 d\sigma < \varepsilon.$$

Therefore by (5)

$$\int \left| \frac{1}{f^+} \left(\frac{k}{f^-} \right)_+ - P_0 \right|^2 f d\sigma < \varepsilon,$$

so P as defined in (6) belongs to \mathcal{S} . Furthermore, with q as defined in (6),

$$Pf - q = f^- \left[\left(\frac{k}{f^-} \right)_+ + \left(\frac{k}{f^-} \right)_- \right] = k.$$

It remains to show that $q \in K^1$. Certainly q belongs to $K^{1/2}$ since it is the product of the two K^1 functions $-f^-$ and $(k/f^-)_-$. But since also

$q = Pf - k$, it belongs to L^1 . Therefore ([6], p. 163) $q \in K^1$.

(ii) If $\log f \notin L^1$, the space \mathcal{P} is identical with $L^2(f)$ ([3], § 33) and so $k/f \in \mathcal{P}$.

We now give the complete solution of Q_3 .

THEOREM 4. (i) *The limit*

$$\lim_{\varepsilon \rightarrow 0+} \int |(k/f_\varepsilon^-)_+|^2 d\sigma$$

exists either finitely or infinitely.

(ii) *A necessary and sufficient condition that Q_3 have a solution P, q is that the limit be finite.*

(iii) *If the limit is finite then*

$$P = \lim (1/f_\varepsilon^+)(k/f_\varepsilon^-)_+$$

in the space $L^2(f)$, and

$$\int |P|^2 f d\sigma = \lim_{\varepsilon \rightarrow 0+} \int |(k/f_\varepsilon^-)_+|^2 d\sigma .$$

Proof. Assume first that Q_3 has a solution P, q and divide both sides of (1) by f_ε^- . Since $q/f_\varepsilon^- \in K^1$ and $\int q/f_\varepsilon^- d\sigma = 0$ we have $q/f_\varepsilon^- \in D$ and $(q/f_\varepsilon^-)_+ = 0$; also $Pf/f_\varepsilon^- \in L^2 \subset D$. Therefore we can apply $(\)_+$ to both sides, obtaining

$$(Pf/f_\varepsilon^-)_+ = (k/f_\varepsilon^-)_+ .$$

Consequently

$$(7) \quad \int |(k/f_\varepsilon^-)_+|^2 d\sigma \leq \int |Pf/f_\varepsilon^-|^2 d\sigma \leq \int |P|^2 f d\sigma ,$$

and so

$$(8) \quad \limsup_{\varepsilon \rightarrow 0+} \int |(k/f_\varepsilon^-)_+|^2 d\sigma < \infty .$$

Conversely suppose that $\{\varepsilon_n\}$ is a sequence of ε 's such that $\varepsilon_n \rightarrow 0+$ and

$$(9) \quad \int |(k/f_\varepsilon^-)_+|^2 d\sigma = O(1) \text{ for } \varepsilon = \varepsilon_n .$$

By Theorem 3(i) there corresponds to each $\varepsilon = \varepsilon_n$ a solution $P_\varepsilon, q_\varepsilon$ of $(f + \varepsilon)P_\varepsilon = k + q_\varepsilon$. We have

$$(10) \quad \int |P_\varepsilon|^2 f d\sigma \leq \int |P_\varepsilon|^2 f_\varepsilon d\sigma = \int |(k/f_\varepsilon^-)_+|^2 d\sigma = O(1) .$$

Thus there exists a subsequence of $\{\varepsilon_n\}$ such that $\{P_\varepsilon\}$ converges weakly

in $L^2(f)$ to an element $P \in \mathcal{P}$. It will follow that $P, Pf - k$ satisfies Q_3 if the L^1 function $q = Pf - k$ satisfies (2). We have for $n = 0, 1, 2, \dots$

$$\begin{aligned} \int q(\phi)e^{-in\phi}d\sigma &= \int \{P_\varepsilon(\phi)[f(\phi) + \varepsilon] - k(\phi)\}e^{-in\phi}d\sigma \\ &\quad + \int [P(\phi) - P_\varepsilon(\phi)]f(\phi)e^{-in\phi}d\sigma - \varepsilon \int P_\varepsilon(\phi)e^{-in\phi}d\sigma \\ &= J_1 + J_2 + J_3 . \end{aligned}$$

Theorem 3(i) implies that $J_1 = 0$. By the weak convergence of the P_ε we can make J_2 as small as desired by taking ε_n sufficiently small. Finally (10) implies that $\int |\varepsilon^{1/2}P_\varepsilon|^2 d\sigma = O(1)$, so by the Schwarz inequality $|J_3| \leq \varepsilon^{1/2} \int |\varepsilon^{1/2}P_\varepsilon| d\sigma = O(\varepsilon^{1/2})$ as $\varepsilon_n \rightarrow 0$. Thus P, q satisfy Q_3 , so (8), holds and (9) is true for *any* sequence $\{\varepsilon_m\}$ of ε 's that converge to $0+$. By what we have shown there corresponds to any such sequence $\{\varepsilon_m\}$ a subsequence such that P_ε converges weakly to the unique (Lemma 2) element P . Thus we can consider ε to be a real variable and conclude that P_ε converges weakly in $L^2(f)$ to $P \in \mathcal{P}$ as $\varepsilon \rightarrow 0+$ provided that

$$\liminf_{\varepsilon \rightarrow 0+} \int |k/f_\varepsilon^-|_+^2 d\sigma < \infty .$$

We next prove that in fact P_ε converges strongly to P in $L^2(f)$. It suffices to show that $\int |P_\varepsilon|^2 fd\sigma \rightarrow \int |P|^2 fd\sigma$. Weak convergence gives

$$\liminf_{\varepsilon \rightarrow 0+} \int |P_\varepsilon|^2 fd\sigma \geq \int |P|^2 fd\sigma .$$

On the other hand, as in (7),

$$\int |P_\varepsilon|^2 fd\sigma \leq \int |P_\varepsilon|^2 f_\varepsilon d\sigma = \int |(k/f_\varepsilon^-)|_+^2 d\sigma \leq \int |P|^2 fd\sigma .$$

so

$$\limsup_{\varepsilon \rightarrow 0+} \int |P_\varepsilon|^2 fd\sigma \leq \int |P|^2 fd\sigma .$$

Thus

$$\lim_{\varepsilon \rightarrow 0+} \int |P_\varepsilon|^2 fd\sigma$$

exists, and equals

$$\lim_{\varepsilon \rightarrow 0+} \int |(k/f_\varepsilon^-)|_+^2 d\sigma = \int |P|^2 fd\sigma .$$

Thus the proof is complete.

4. **Solution of Q_1 .** In Q_1 we wish to find

$$I_1(g, \mu) = \inf_{P \in \mathcal{S}'_0} \left[\int |g + P|^2 d\mu \right]^{\frac{1}{2}},$$

where g is a given function in $L^2(\mu)$. Since $I_1(g, \mu)$ represents the distance from g to the manifold \mathcal{S}_0 in $L^2(\mu)$, there exists a (unique) function P belonging to the closure \mathcal{S}' of \mathcal{S}_0 in $L^2(\mu)$ such that

$$I_1(g, \mu) = \left[\int |g + P|^2 d\mu \right]^{\frac{1}{2}}.$$

This function P is such that $g + P$ is orthogonal to \mathcal{S}_0 , so

$$\int [g(\phi) + P(\phi)]e^{-in\phi} d\mu(\phi) = 0 \quad n = 0, 1, 2, \dots .$$

It follows from a theorem of the brothers Riesz ([6], p. 158) that the measure ν given by

$$\nu(E) = \int_E [g(\phi) + P(\phi)] d\mu(\phi)$$

is absolutely continuous with respect to Lebesgue measure. Let F be a Borel set of Lebesgue measure zero such that $\mu_s((-\pi, \pi] - F) = 0$. Then $g + P$ vanishes on F almost everywhere with respect to μ_s , so

$$\int_F |g + P|^2 d\mu_s = 0$$

and

$$\int |g + P|^2 d\mu = \int_{\mathcal{E}} |g + P|^2 d\mu_A = \int |g + P|^2 f d\sigma .$$

Since $\mu \geq \mu_A$ it follows that $I_1(g, \mu) = I_1(g, f)$, and this common value is attained by the same extremizing function $P \in \mathcal{S}' \subset \mathcal{S}$.

Now,

$$\int [g(\phi) + P(\phi)]e^{-in\phi} f(\phi) d\sigma = 0 \quad n = 0, 1, \dots ,$$

so if we set $q = (g + P)f$ we have $Pf = -gf + q$, where $P \in \mathcal{S}$ and q satisfies (2). Since $(gf)^2/f = g^2f \in L^1$, we can apply Theorem 3 to this situation. The extremizing function

$$P = \begin{cases} -(1/f_+)(gf_+) & \text{if } \log f \in L^1 \\ -g & \text{if } \log f \notin L^1 , \end{cases}$$

and since

$$I_1(g, f) = \left[\int |g + P|^2 f d\sigma \right]^{\frac{1}{2}} = \left[\int |q|^2 f d\sigma \right]^{\frac{1}{2}}$$

we have

$$I_1(g, \mu) = I_1(g, f) = \begin{cases} \left[\int |(gf^+)_-|^2 d\sigma \right]^{\frac{1}{2}} & \text{if } \log f \in L^1 \\ 0 & \text{if } \log f \notin L^1. \end{cases}$$

5. **Solution of Q_2 .** Given $h \in L^1$, we will evaluate

$$I_2(h, \mu) = \sup_{P \in \mathcal{S}_0} \left\{ \left| \int Ph d\sigma \right| / \left[\int |P|^2 d\mu \right]^{\frac{1}{2}} \right\}.$$

Since $\mu \geq \mu_A$ it is clear that if $I_2(h, f)$ is finite so is $I_2(h, \mu)$. We shall show that, conversely, if $I_2(h, \mu)$ is finite then so is $I_2(h, f)$ and in fact $I_2(h, f) = I_2(h, \mu)$. So now suppose $I_2(h, \mu) < \infty$. Then the linear functional L on \mathcal{S}_0 given by

$$L(P) = \int Ph d\sigma$$

is bounded on $L^2(\mu)$. Therefore if \mathcal{S}' denotes the closure of \mathcal{S}_0 in $L^2(\mu)$, there is a uniquely determined $Q \in \mathcal{S}'$ such that $L(P) = \int P\bar{Q} d\mu$. Then we have

$$\int e^{-in\phi} [Q(\phi)d\mu(\phi) - \bar{h}(\phi)d\sigma(\phi)] = 0 \quad n = 0, 1, \dots.$$

We again apply the F. and M. Riesz theorem, and deduce that the measure ν given by

$$\nu(E) = \int_E Qd\mu - \int_E h d\sigma$$

is absolutely continuous with respect to Lebesgue measure. Letting F' be a Borel set of Lebesgue measure zero such that $\mu_s((-\pi, \pi] - F') = 0$, we see that Q vanishes on F' almost everywhere with respect to μ_s . Consequently

$$\int e^{-in\phi} [Q(\phi)f(\phi) - \bar{h}(\phi)]d\sigma(\phi) = 0 \quad n = 0, 1, \dots,$$

so $Qf = \bar{h} + q$, where $Q \in \mathcal{S}' \subset \mathcal{S}$ and q satisfies (2). Thus the linear functional

$$L(P) = \int Ph d\sigma = \int P\bar{Q}f d\sigma,$$

$P \in \mathcal{S}_0$, is bounded on $L^2(f)$, so $I_2(h, f)$ is finite and in fact equals $I_2(h, \mu)$. We deduce from Theorem 4 that

$$I_2(h, \mu) = I_2(h, f) = \lim_{\varepsilon \rightarrow 0+} \left[\int |(\bar{h}/f_\varepsilon^-)_+|^2 d\sigma \right]^{\frac{1}{2}},$$

and Q may be exhibited as an $L^2(f)$ limit in the mean.

6. Some formulae for $I_2(h, \mu)$. We can obtain a simpler formula for $I_2(h, \mu)$ if we assume that $h^2/f \in L^1$ and apply Theorem 3. Then

$$I_2(h, \mu) = \begin{cases} \left[\int |(\bar{h}/f^-)_+|^2 d\sigma \right]^{\frac{1}{2}} = \left[\int |(e^{-i\phi}h(\phi)/f^+(\phi))_-|^2 d\sigma(\phi) \right]^{\frac{1}{2}} & \text{if } \log f \in L^1, \\ \left[\int |h|^2/f d\sigma \right]^{\frac{1}{2}} & \text{if } \log f \notin L^1. \end{cases}$$

This, in conjunction with our solution of Q_1 , gives the duality discussed in Theorem 1. Note that the hypothesis $1/f \in L^1$ of Theorem 1 implies that $\log f \in L^1$.

Another simple formula for $I_2(h, \mu)$ is available if we know that the Fourier series $\sum_{-\infty}^{\infty} h_n e^{in\phi}$ of h is such that $h_{-n} = O(R_0^{-n})$ as $n \rightarrow +\infty$ for some $R_0 > 1$. Then the function $H(z) = \sum_0^{\infty} h_{-n} z^{-n}$ is analytic in $|z| > 1/R_0$. We have

$$\int |(\bar{h}/f_\varepsilon^-)_+|^2 d\sigma = \int |(e^{-i\phi}h(\phi)/f_\varepsilon^+(\phi))_-|^2 d\sigma,$$

which by the Parseval relation equals

$$\begin{aligned} \sum_{n=0}^{\infty} \left| \int e^{in\phi} h(\phi) f_\varepsilon^+(\phi) d\sigma \right|^2 &= \sum_{n=0}^{\infty} \left| \frac{1}{2\pi} \int_{|z|=1} z^{n+1} H(z) / F_\varepsilon^+(z) dz \right|^2 \\ &= \sum_{n=0}^{\infty} \left| \frac{1}{2\pi} \int_{|z|=R} z^{n+1} H(z) / F_\varepsilon^+(z) dz \right|^2, \end{aligned}$$

where $1/R_0 < R < 1$. Let us also assume that $\log f \in L^1$, so F^+ is well-defined and

$$H(Re^{i\phi})/F_\varepsilon^+(Re^{i\phi}) \longrightarrow H(Re^{i\phi})/F^+(Re^{i\phi})$$

in L^2 as $\varepsilon \rightarrow 0+$. It follows that

$$I_2(h, \mu)^2 = \sum_{n=0}^{\infty} \left| \frac{1}{2\pi} \int_{|z|=R} z^{n+1} H(z) / F^+(z) dz \right|^2.$$

Now, if we write

$$\frac{1}{F^+(z)} = \sum_{n=0}^{\infty} f_n z^n,$$

then

$$I_2(h, \mu)^2 = \sum_{n=0}^{\infty} \left| \sum_{m=0}^{\infty} h_{-n-m} f_m \right|^2.$$

Thus if H is the Hankel matrix $[h_{-n-m}]_{n,m=0}^{\infty}$, and Φ the column vector with components f_0, f_1, \dots , then

$$I_2(h, \mu) = \| H\Phi \|,$$

where the norm is that of l^2 .

For example, let α be such that $|\alpha| < 1$ and consider

$$\sup_{P \in \mathcal{F}} \left\{ |P(\alpha)| / \left(\int |P|^2 d\mu \right)^{\frac{1}{2}} \right\}.$$

Thus we wish to evaluate $I_2(1/(1 - \alpha e^{-i\phi}), \mu)$. Here $h_{-n} = \alpha^n$, $n = 0, 1, \dots$, so

$$I_2(h, \mu)^2 = \sum_{n=0}^{\infty} \left| \sum_{m=0}^{\infty} \alpha^{n+m} f_m \right|^2 = 1/[(1 - |\alpha|^2) F^+(\alpha)^2],$$

as in [2], p. 48.

BIBLIOGRAPHY

1. A. Beurling, *On two problems concerning linear transformations in Hilbert space*, Acta Math., **81** (1948), 239-255.
2. F. F. Bonsall, *Dual extremum problems in the theory of functions*, Jour. London Math. Soc., **31** (1956), 1-5-110.
3. U. Grenander and G. Szegö, *Toeplitz forms and their applications*, Berkeley and Los Angeles, 1958.
4. N. I. Muskhelishvili, *Singular integral equations*, Groningen, 1953.
5. Szegö, *Orthogonal polynomials*, A. M. S. colloquium publication, **23** (1939).
6. A. Zygmund, *Trigonometrical series*, New York, 1952.

INSTITUTE FOR ADVANCED STUDY
UNIVERSITY OF VIRGINIA
CORNELL UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DAVID GILBARG

Stanford University
Stanford, California

F. H. BROWNELL

University of Washington
Seattle 5, Washington

A. L. WHITEMAN

University of Southern California
Los Angeles 7, California

L. J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

T. M. CHERRY

D. DERRY

E. HEWITT

A. HORN

L. NACHBIN

M. OHTSUKA

H. L. ROYDEN

M. M. SCHIFFER

E. SPANIER

E. G. STRAUS

F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON
* * *

AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
HUGHES AIRCRAFT COMPANY
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 2120 Oxford Street, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 10, No. 4

December, 1960

M. Altman, <i>An optimum cubically convergent iterative method of inverting a linear bounded operator in Hilbert space</i>	1107
Nesmith Cornett Ankeny, <i>Criterion for rth power residuacity</i>	1115
Julius Rubin Blum and David Lee Hanson, <i>On invariant probability measures I</i>	1125
Frank Featherstone Bonsall, <i>Positive operators compact in an auxiliary topology</i>	1131
Billy Joe Boyer, <i>Summability of derived conjugate series</i>	1139
Delmar L. Boyer, <i>A note on a problem of Fuchs</i>	1147
Hans-Joachim Bremermann, <i>The envelopes of holomorphy of tube domains in infinite dimensional Banach spaces</i>	1149
Andrew Michael Bruckner, <i>Minimal superadditive extensions of superadditive functions</i>	1155
Billy Finney Bryant, <i>On expansive homeomorphisms</i>	1163
Jean W. Butler, <i>On complete and independent sets of operations in finite algebras</i>	1169
Lucien Le Cam, <i>An approximation theorem for the Poisson binomial distribution</i>	1181
Paul Civin, <i>Involutions on locally compact rings</i>	1199
Earl A. Coddington, <i>Normal extensions of formally normal operators</i>	1203
Jacob Feldman, <i>Some classes of equivalent Gaussian processes on an interval</i>	1211
Shaul Foguel, <i>Weak and strong convergence for Markov processes</i>	1221
Martin Fox, <i>Some zero sum two-person games with moves in the unit interval</i>	1235
Robert Pertsch Gilbert, <i>Singularities of three-dimensional harmonic functions</i>	1243
Branko Grünbaum, <i>Partitions of mass-distributions and of convex bodies by hyperplanes</i>	1257
Sidney Morris Harmon, <i>Regular covering surfaces of Riemann surfaces</i>	1263
Edwin Hewitt and Herbert S. Zuckerman, <i>The multiplicative semigroup of integers modulo m</i>	1291
Paul Daniel Hill, <i>Relation of a direct limit group to associated vector groups</i>	1309
Calvin Virgil Holmes, <i>Commutator groups of monomial groups</i>	1313
James Fredrik Jakobsen and W. R. Utz, <i>The non-existence of expansive homeomorphisms on a closed 2-cell</i>	1319
John William Jewett, <i>Multiplication on classes of pseudo-analytic functions</i>	1323
Helmut Klingens, <i>Analytic automorphisms of bounded symmetric complex domains</i>	1327
Robert Jacob Koch, <i>Ordered semigroups in partially ordered semigroups</i>	1333
Marvin David Marcus and N. A. Khan, <i>On a commutator result of Tausky and Zassenhaus</i>	1337
John Glen Marica and Steve Jerome Bryant, <i>Unary algebras</i>	1347
Edward Peter Merkes and W. T. Scott, <i>On univalence of a continued fraction</i>	1361
Shu-Teh Chen Moy, <i>Asymptotic properties of derivatives of stationary measures</i>	1371
John William Neuberger, <i>Concerning boundary value problems</i>	1385
Edward C. Posner, <i>Integral closure of differential rings</i>	1393
Marian Reichaw-Reichbach, <i>Some theorems on mappings onto</i>	1397
Marvin Rosenblum and Harold Widom, <i>Two extremal problems</i>	1409
Morton Lincoln Slater and Herbert S. Wilf, <i>A class of linear differential-difference equations</i>	1419
Charles Robson Storey, Jr., <i>The structure of threads</i>	1429
J. François Treves, <i>An estimate for differential polynomials in $\partial/\partial z_1, \dots, \partial/\partial z_n$</i>	1447
J. D. Weston, <i>On the representation of operators by convolutions integrals</i>	1453
James Victor Whittaker, <i>Normal subgroups of some homeomorphism groups</i>	1469