

Pacific Journal of Mathematics

ON THE ACTION OF A LOCALLY COMPACT GROUP ON E_n

FRANK J. HAHN

ON THE ACTION OF A LOCALLY COMPACT GROUP ON E_n

F. J. HAHN

It is known [2, p. 208] that if a locally compact group acts effectively and differentiably on E_n then it is a Lie group. The object of this note is to show that if the differentiability requirements are replaced by some weaker restrictions, given later on, the theorem is still true. Let G be a locally compact group acting on E_n and let the coordinate functions of the action be given by $f_i(g, x_1, \dots, x_n)$, $1 \leq i \leq n$. For economy we introduce the following notation

$$Q_{ij}(g, t, x) = \frac{f_i(g, x_1, \dots, x_j + t, \dots, x_n) - f_i(g, x_1, \dots, x_j, \dots, x_n)}{t}.$$

We denote by $\sigma(Q_{ij}(e, 0, x))$ the oscillation of $Q_{ij}(g, t, x)$ at the point $(e, 0, x)$.

Before proceeding there is one simple remark to be made on matrices. If $A = (a_{ij})$ is an $n \times n$ matrix such that $|a_{ij} - \delta_{ij}| < (1/n)$ then A is non-singular. If A were singular there would be a vector x such that $\sum_i x_i^2 = 1$ and $Ax = 0$. From the Schwarz inequality it follows that $x_i^2 = (\sum_j (a_{ij} - \delta_{ij})x_j)^2 < (1/n)$ and consequently $1 = \sum x_i^2 < 1$ which is impossible. If $|a_{ij} - \delta_{ij}| \leq (\alpha/n)$, where $0 < \alpha < 1$, then the determinant of A is bounded away from zero since the determinant is a continuous function and the set $\{a_{ij}: |a_{ij} - \delta_{ij}| \leq (\alpha/n)\}$ is compact in E_{n^2} .

THEOREM 1. *If T is a pointwise periodic homeomorphism of E_n then T is periodic.*

Proof. [2, p. 224.]

THEOREM 2. *If G is a compact, zero dimensional, monothetic group acting effectively on E_n and satisfying*

$$(*) \quad \sigma(Q_{ij}(e, 0, x)) < \frac{\varepsilon}{n}, \quad 0 < \varepsilon < 1, \quad \text{for each } x \text{ in } E_n;$$

then G is a finite cyclic group.

Proof. Since G is monothetic, let a be an element whose powers are dense in G . It is enough to show that there is a power of a which leaves E_n pointwise fixed since the action of G is effective.

If q is a positive integer we let

$$T_i^q(g, x) = x_i + f_i(g, x) + \cdots + f_i(g^{q-1}, x).$$

If $y = (y_i)$ and $x = (x_i)$ let

$$T_{ij}^q(g, x, y) = \frac{T_i^q(g, x_1, \dots, x_{j-1}, y_j, \dots, y_n) - T_i^q(g, x_1, \dots, x_j, y_{j+1}, \dots, y_n)}{y_j - x_j}$$

for $y_j \neq x_j$ and zero otherwise. If we let $y = f(g, x)$ then we obtain

$$\begin{aligned} f_i(g^q, x) - x_i &= T_i^q(g, y) - T_i^q(g, x) \\ &= \sum_{j=1}^n T_{ij}^q(g, x, y)(y_j - x_j) \\ &= q \cdot \sum_{i=1}^n \frac{1}{q} T_{ij}^q(g, x, y)(y_j - x_j). \end{aligned}$$

Because of the fact that $f_i(e, x) = x_i$ and because of (*) it follows that there is a compact neighborhood $U(x)$ of the identity of G such that if $g, \dots, g^q \in U(x)$ then $|(1/q)T_{ij}^q(g, x, y) - \delta_{ij}| \leq (\alpha/n)$, $0 < \varepsilon < \alpha < 1$. It follows that if T is the matrix with entries $(1/q)T_{ij}^q(g, x, y)$ then T is non-singular and its determinant is bounded away from zero uniformly in q , so the determinant of the inverse is bounded uniformly in q ; thus

$$(f(g, x) - x) = (y - x) = \left(\delta_{ij} \frac{1}{q} \right) \cdot T^{-1} \cdot (f(g^q, x) - x).$$

Since G is monothetic and zero dimensional there is a power of a such that if $g = a^p$ then all the powers of g lie in $U(x)$. Since $U(x)$ is compact it follows that the vectors $f(g^q, x) - x$ are bounded uniformly in q and thus $f(g, x) - x = f(a^p, x) - x = 0$. Hence a is pointwise periodic on E_n and it follows from Theorem 1 that it is periodic and consequently has a power leaving E_n pointwise fixed.

From this it follows quickly that if G is a locally compact group acting effectively on E_n and satisfying (*) then it is a Lie group. This follows from the fact that since G is effective it must be finite dimensional [1] and then if G is not a Lie group it must contain a compact, non-finite zero dimensional subgroup H [2, p. 237] which acts effectively. H has small subgroups which act effectively and it follows from Newman's theorem [3, 4] that H cannot have arbitrarily small elements of finite order. Thus H has an element a of infinite order such that the compact subgroup generated by a acts effectively on E_n and satisfies (*) but by Theorem 2 this is impossible.

BIBLIOGRAPHY

1. D. Montgomery, *Finite dimensionality of certain transformation groups*, Ill. J. Math. 1, No. 1 (1957), 28-35.

2. D. Montgomery and L. Zippin, *Topological Transformation Groups*, Interscience Publishers, New York (1955).
3. M. H. A. Newman, *A theorem on periodic transformations of spaces*, Quart. J. Math. **2** (1931), 1-8.
4. P. A. Smith, *Transformations of finite period, III, Newman's theorem*, Ann. of Math. **42** (1941), 446-458.

INSTITUTE FOR ADVANCED STUDY
PRINCETON, NEW JERSEY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RALPH S. PHILLIPS
Stanford University
Stanford, California

F. H. BROWNELL
University of Washington
Seattle 5, Washington

A. L. WHITEMAN
University of Southern California
Los Angeles 7, California

L. J. PAIGE
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH
T. M. CHERRY

D. DERRY
M. OHTSUKA

H. L. ROYDEN
E. SPANIER

E. G. STRAUS
F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
HUGHES AIRCRAFT COMPANY
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

Reprinted 1966 in the United States of America

Pacific Journal of Mathematics

Vol. 11, No. 1

November, 1961

A. A. Albert, <i>Generalized twisted fields</i>	1
Richard Arens, <i>Operational calculus of linear relations</i>	9
John Herbert Barrett, <i>Disconjugacy of a self-adjoint differential equation of the fourth order</i>	25
Paul Richard Beesack, <i>Hardy's inequality and its extensions</i>	39
Julius Rubin Blum and David Lee Hanson, <i>On invariant probability measures. II</i>	63
Robert Allen Bonic, <i>Symmetry in group algebras of discrete groups</i>	73
R. Creighton Buck, <i>Multiplication operators</i>	95
Jack Gary Ceder, <i>Some generalizations of metric spaces</i>	105
Meyer Dwass, <i>Random crossings of cumulative distribution functions</i>	127
Albert Edrei, Wolfgang H. J. Fuchs and Simon Hellerstein, <i>Radial distribution and deficiencies of the values of a meromorphic function</i>	135
William Cassidy Fox, <i>Harmonic functions with arbitrary local singularities</i>	153
Theodore Thomas Frankel, <i>Manifolds with positive curvature</i>	165
Avner Friedman, <i>A strong maximum principle for weakly subparabolic functions</i>	175
Watson Bryan Fulks and J. O. Sather, <i>Asymptotics. II. Laplace's method for multiple integrals</i>	185
Adriano Mario Garsia and Eugene Richard Rodemich, <i>An embedding of Riemann surfaces of genus one</i>	193
Irving Leonard Glicksberg, <i>Weak compactness and separate continuity</i>	205
Branko Grünbaum, <i>On a conjecture of H. Hadwiger</i>	215
Frank J. Hahn, <i>On the action of a locally compact group on E_n</i>	221
Magnus R. Hestenes, <i>Relative hermitian matrices</i>	225
G. K. Kalisch, <i>On similarity invariants of certain operators in L_p</i>	247
Yitzhak Katznelson and Walter Rudin, <i>The Stone-Weierstrass property in Banach algebras</i>	253
Samir A. Khabbaz, <i>The subgroups of a divisible group G which can be represented as intersections of divisible subgroups of G</i>	267
Marvin Isadore Knopp, <i>Construction of a class of modular functions and forms</i>	275
Charles Alan McCarthy, <i>Commuting Boolean algebras of projections</i>	295
T. M. MacRobert, <i>Transformations of series of E-functions</i>	309
Heinz Renggli, <i>An inequality for logarithmic capacities</i>	313
M. S. Robertson, <i>Applications of the subordination principle to univalent functions</i>	315
David Sachs, <i>Partition and modulated lattices</i>	325
Frank S. Scalora, <i>Abstract martingale convergence theorems</i>	347
Elbert A. Walker, <i>Torsion endomorphic images of mixed Abelian groups</i>	375
Morgan Ward, <i>The prime divisors of Fibonacci numbers</i>	379
Charles R. B. Wright, <i>On the nilpotency class of a group of exponent four</i>	387