

Pacific Journal of Mathematics

OPERATORS OF MEROMORPHIC TYPE WITH MULTIPLE POLES OF THE RESOLVENT

JOHN IRVING DERR AND ANGUS E. TAYLOR

OPERATORS OF MEROMORPHIC TYPE WITH MULTIPLE POLES OF THE RESOLVENT

JOHN DERR AND ANGUS E. TAYLOR

1. Introduction. Let X be a complex Banach space and let A be a bounded linear operator on X such that the spectrum $\sigma(A)$ is a denumerable set of points with $\lambda = 0$ as the only point of accumulation. Suppose, also, that each nonzero point of $\sigma(A)$ is a pole of the resolvent $R_\lambda(A)$. Then we shall call A an operator of meromorphic type.

Let $\lambda_1, \lambda_2, \dots$ be an enumeration of the distinct nonzero points of $\sigma(A)$, and let E_n be the residue of $R_\lambda(A)$ at λ_n . Then $E_n^2 = E_n$, $E_n \neq 0$, $E_n \neq I$, and $E_n E_m = 0$ if $m \neq n$. Moreover, E_n commutes with A . If the order of the pole at λ_n is q_n , then

$$(A - \lambda_n)^{q_n} E_n = 0, \quad (A - \lambda_n)^{q_n - 1} E_n \neq 0.$$

When $q_n > 1$, the singular part of the Laurent expansion of $R_\lambda(A)$ in the neighborhood of λ_n is

$$(1) \quad S_n(\lambda) = \frac{E_n}{\lambda - \lambda_n} + \sum_{k=2}^{q_n} \frac{F_n^{k-1}}{(\lambda - \lambda_n)^k},$$

where

$$(2) \quad F_n = (A - \lambda_n) E_n.$$

In this case

$$(3) \quad F_n^{q_n - 1} \neq 0, \quad F_n^{q_n} = 0.$$

If $q_n = 1$, we have

$$(4) \quad S_n(\lambda) = \frac{E_n}{\lambda - \lambda_n}, \quad F_n = 0.$$

Observe that $E_n F_n = F_n E_n = F_n$. For the relevant facts about the coefficients in the Laurent expansion of $R_\lambda(A)$ in the neighborhood of an isolated singularity, see Taylor [5]. (Numbers in square brackets refer to the works cited at the end of the paper.)

Throughout this paper we shall be concerned with series expansions of operator-valued functions of λ . All convergence questions are examined from the point of view of the uniform topology in the space of bounded linear operators on X . (We shall denote this space of operators by $[X]$.) Thus, if T_n and T are elements of $[X]$, $T_n \rightarrow T$ means $\|T_n - T\| \rightarrow 0$.

Because $S_n(\lambda)$ is regular when $|\lambda| > |\lambda_n|$, it has an expansion in powers of $1/\lambda$, convergent when $|\lambda| > |\lambda_n|$. If p is a positive integer, we denote by $P_n^{(p)}(\lambda)$ the sum of the terms of degree $\leq p$ in $1/\lambda$ in this expansion of $S_n(\lambda)$. An explicit formula will be given presently.

It turns out that a sequence $\{p_n\}$ can be chosen in such a way that the series

$$(5) \quad \sum_{n=1}^{\infty} [S_n(\lambda) - P_n^{(p_n)}(\lambda)]$$

converges when λ is not in $\sigma(A)$, and such that, moreover, the difference

$$(6) \quad R_\lambda(A) - \sum_{n=1}^{\infty} [S_n(\lambda) - P_n^{(p_n)}(\lambda)]$$

has removable singularities at the points $\lambda_1, \lambda_2, \dots$ and coincides in the resolvent set $\rho(A)$ with a function given by a series

$$(7) \quad \sum_{n=0}^{\infty} \frac{Q_n}{\lambda^n}$$

which converges for every nonzero λ . Here Q_0, Q_1, \dots are bounded linear operators. The sequence $\{p_n\}$ is not unique. It may be chosen along with the imposition of rather strong conditions on the mode of convergence of the series (5). In a paper by one of us (Taylor [6]) a study was made of the situation when all the poles are of the first order and a uniform convergence condition was imposed on (5). Subsequently, in a paper delivered at the Jerusalem Symposium on Linear Spaces, in July 1960, Taylor discussed the situation when an absolute and uniform convergence condition is placed on the series

$$(8) \quad \sum_n \|S_n(\lambda) - P_n^{(p_n)}(\lambda)\|.$$

Most of the results reported on by Taylor were for the case of simple poles. The theory for the multiple-pole case was begun by Derr in his doctoral dissertation [2]. Some contributions to the multiple-pole theory, especially where the condition of absolute convergence is imposed, were made by Taylor. In this paper we concentrate on the discussion of the multiple-pole case.

In Taylor [7] the following theorem was proved. We repeat the statement here for convenience.

THEOREM 1. *If A is of meromorphic type and if the foregoing notation is established, then there exists a sequence $\{p_n\}$ of positive integers such that for each $\delta > 0$, if m is so large that $|\lambda_k| < \delta$ when $k \geq m$, then the series*

$$(9) \quad \sum_{n=m}^{\infty} \|S_n(\lambda) - P_n^{(p_n)}(\lambda)\|$$

converges uniformly in the part of the plane for which $|\lambda| \geq \delta$. Moreover, there exists a sequence $\{Q_n\}$ of elements of $[X]$ such that the series (7) converges if $\lambda \neq 0$, and for each λ in $\rho(A)$ the resolvent of A has the expansion

$$(10) \quad R_\lambda(A) = \sum_{n=1}^{\infty} [S_n(\lambda) - P_n^{(p_n)}(\lambda)] + \sum_{n=0}^{\infty} \lambda^{-n} Q_n.$$

On the basis of this theorem we make the following definitions: A sequence $\{p_n\}$ of the sort specified in Theorem 1 will be called *an absolute index sequence for A relative to $\{\lambda_n\}$* . If $\{p_n\}$ is an absolute index sequence such that p_n has the same value p for every n , we shall say that A admits the absolute index p . We do not need to add the phrase “relative to $\{\lambda_n\}$ ” in this case, because the terms in the series (9) are nonnegative; on this account, if the series converges uniformly in the manner described, it remains thus uniformly convergent after an arbitrary rearrangement of the order of the terms. If A admits the absolute index p , but no smaller absolute index, we say that A has *minimal absolute index p* . The minimal absolute index depends only on A , not on any particular order of listing of the poles of $R_\lambda(A)$.

Now, it may happen that a sequence $\{p_n\}$ of positive integers has the property that for each $\delta > 0$, if m is chosen so large that $|\lambda_k| < \delta$ when $k \geq m$, then the series

$$(11) \quad \sum_{k=m}^{\infty} [S_n(\lambda) - P_n^{(p_n)}(\lambda)]$$

converges uniformly in the part of the plane for which $|\lambda| \geq \delta$. Observe carefully the difference between (9) and (11). The series (9) is numerical, whereas (11) is a series whose terms are members of $[X]$. With this modified condition on the sequence $\{p_n\}$, it will be true that the operator function $F(\lambda)$ defined by

$$(12) \quad F(\lambda) = \sum_{n=1}^{\infty} [S_n(\lambda) - P_n^{(p_n)}(\lambda)]$$

is analytic on $\rho(A)$, regular at $\lambda = \infty$, and has a pole at each of the points $\lambda_1, \lambda_2, \dots$, the singular part of the Laurent expansion in the neighborhood of λ_n being $S_n(\lambda)$. Therefore, under these conditions, $R_\lambda(A) - F(\lambda)$ will have removable singularities at the points $\lambda_1, \lambda_2, \dots$, and will be regular at ∞ . Accordingly, we shall have a representation of the form (10), where the series involving the Q_n 's converges when $\lambda \neq 0$.

A sequence $\{p_n\}$ for which we have uniform convergence as specified in connection with the series (11) will be called *a uniform index sequence for A relative to $\{\lambda_n\}$* . If $p_n = p$ for every n , we call p a *uniform index for A relative to $\{\lambda_n\}$* . We can also define a *minimal uniform*

index for A relative to $\{\lambda_n\}$, provided there exists at least one such uniform index. It is to be emphasized that the notion of a uniform index for A must be defined in relation to a specified way of enumerating the poles of $R_\lambda(A)$. This is because there is no guarantee that uniform convergence of the series (11) is preserved when the order of the term is rearranged. In fact, an example is known (Berkson [1]) in which A has the uniform index 1 relative to a certain enumeration $\{\lambda_n\}$, but the series (11) can be rearranged in such a way as to be divergent at all points of $\rho(A)$.

In Theorem 10 (§ 5) we give a necessary and sufficient condition for A to have minimal uniform index p relative to $\{\lambda_n\}$. The case $p = 1$ is handled in Theorem 5 (§ 3). The condition is stated entirely in terms of λ_n , E_n , and F_n . A corresponding necessary and sufficient condition for A to have minimal absolute index p is given in Theorem 13 (§ 6).

These results are of decisive importance for the theory of operators of meromorphic type. With the criteria of Theorems 10 and 13 it is much easier to test for uniform and absolute indices than would be possible by direct reliance on the definitions. The construction of examples is enormously facilitated.

Another important part of the paper is in § 4, where we discuss the decomposition $A = B + C$ for an operator A of meromorphic type and minimal uniform index 1 relative to $\{\lambda_n\}$. In this decomposition, B is of the same general character as A , with $\sigma(A) = \sigma(B)$, and B is *canonical* in a well-defined sense. Moreover, $BC = CB = 0$ and C is quasinilpotent. This decomposition is established in Theorem 8, and its uniqueness is described in Theorem 9.

2. Some preliminary results. In this section we assemble some results and formulas for later use.

LEMMA 1. *Suppose that $E, F, \in [X]$ and that*

$$(13) \quad E^2 = E, \quad EF = FE = F, \quad E \neq 0.$$

Suppose that there is a positive integer q such that

$$(14) \quad F^{q-1} \neq 0, \quad F^q = 0.$$

Then, if $\alpha \neq 0$ and $B = \alpha E + F$, the resolvent of B is

$$(15) \quad R_\lambda(B) = \frac{I - E}{\lambda} + \frac{E}{\lambda - \alpha} + \frac{F}{(\lambda - \alpha)^2} + \cdots + \frac{F^{q-1}}{(\lambda - \alpha)^q}$$

(the terms involving F occurring only if $q > 1$). The spectrum of B consists of 0 and α if $E \neq I$ and of α alone if $E = I$.

Proof. Let C denote the operator in the right member of formula (15), where λ is fixed, with $\lambda \neq 0, \lambda \neq \alpha$. It is an easy matter to verify that $C(\lambda - B) = (\lambda - B)C = I$, whence it follows that $\lambda \in \rho(B)$ and $R_\lambda(B) = C$ when λ is thus restricted. We leave the calculations to the reader. Now, α is an eigenvalue of B . For, if $q > 1$, choose x so that $y = F^{q-1}x \neq 0$; this is possible, by (14). Then $Ey = y$ because $EF = F$; hence $By = (\alpha E + F)F^{q-1}x = \alpha y$, because $F^q = 0$. If $q = 1$, choose x so that $y = Ex \neq 0$. We can do this, because $E \neq 0$. Then, since $F = 0$ in this case, $By = \alpha Ey = \alpha y$. Hence it is certain that $\alpha \in \sigma(B)$, regardless of the value of q .

If $I - E \neq 0$, and if $y = x - Ex \neq 0$, we see that $By = 0$, because $B = BE$. Hence $0 \in \sigma(B)$ in this case. On the other hand, if $E = I$, a direct calculation shows that $0 - B$ has the inverse

$$\frac{I}{-\alpha} + \frac{F}{(-\alpha)^2} + \dots + \frac{F^{q-1}}{(-\alpha)^q},$$

so that $0 \in \rho(B)$ in this case. This completes the proof of Lemma 1.

It now follows from general spectral theory that the spectral radius of $\alpha E + F$ is $|\alpha|$, and that $R_\lambda(B)$ is given by the series

$$(16) \quad R_\lambda(B) = \sum_{k=0}^{\infty} \frac{(\alpha E + F)^k}{\lambda^{k+1}}$$

when $|\lambda| > |\alpha|$. From (15) we then see that

$$(17) \quad \frac{E}{\lambda - \alpha} + \sum_{j=1}^{q-1} \frac{F^j}{(\lambda - \alpha)^{j+1}} = \frac{E}{\lambda} + \sum_{k=1}^{\infty} \frac{(\alpha E + F)^k}{\lambda^{k+1}},$$

provided that $|\lambda| > |\alpha|$.

The foregoing considerations can be applied to obtain a formula for $P_n^{(p)}(\lambda)$, which is, by definition, the sum of the terms of degree $\leq p$ in $1/\lambda$ in the expansion of $S_n(\lambda)$ in powers of $1/\lambda$. By comparing (1) and (17) we see that

$$(18) \quad S_n(\lambda) = \frac{E_n}{\lambda} + \sum_{k=1}^{\infty} \frac{(\lambda_n E_n + F_n)^k}{\lambda^{k+1}}$$

if $|\lambda| > |\lambda_n|$. Therefore

$$(19) \quad P_n^{(1)}(\lambda) = \frac{E_n}{\lambda},$$

$$(20) \quad P_n^{(p)}(\lambda) = \frac{E_n}{\lambda} + \sum_{k=2}^p \frac{(\lambda_n E_n + F_n)^{k-1}}{\lambda^k} \text{ if } p \geq 2.$$

The formula (18) could also be obtained from (1) by direct use of binomial

series. This was done, in the first instance, by Derr in his thesis (Derr [2]). The present method, using spectral theory, is rather elegant.

Now that we have the formula (18), it is easy to calculate the formulas for the operator coefficients Q_0, Q_1, \dots occurring in (10). For this calculation it is sufficient to assume that $\{p_n\}$ is a uniform index sequence for A relative to $\{\lambda_n\}$. (An absolute index sequence is also a uniform index sequence of course.) From general spectral theory it is known that

$$\frac{1}{2\pi i} \oint \lambda^j R_\lambda(A) d\lambda = \begin{cases} 0 & \text{if } j < 0 \\ I & \text{if } j = 0 \\ A^j & \text{if } j > 0, \end{cases}$$

where the integration is taken counter clockwise over a large circle enclosing $\sigma(A)$. From (10), (18), and (20) we find that

$$(21) \quad Q_0 = 0, \quad Q_1 = I,$$

and

$$(22) \quad Q_{j+1} = A^j - \sum_{p_n \leq j} (\lambda_n E_n + F_n)^j \quad \text{if } j \geq 1.$$

The series on the right in (22) is understood to mean

$$(23) \quad \sum_{n=1}^{\infty} \varepsilon_{n,j} (\lambda_n E_n + F_n)^j,$$

where $\varepsilon_{n,j} = 1$ if $p_n \leq j$ and $\varepsilon_{n,j} = 0$ if $j < p_n$.

It is part of the conclusion that the series in (23) is convergent in $[X]$. As a consequence, we have the following theorem:

THEOREM 2. *If A admits the uniform index p , the series*

$$(24) \quad \sum_{n=1}^{\infty} (\lambda_n E_n + F_n)^j$$

converges when $p \leq j$.

Concerning an absolute index sequence we have:

THEOREM 3. *If $\{p_n\}$ is an absolute index sequence for A , the series*

$$(25) \quad \sum_{n=1}^{\infty} \|\varepsilon_{n,j} (\lambda_n E_n + F_n)^j\|$$

converges for each $j \geq 1$.

Proof. We fix j and suppose $\delta > 0, \varepsilon > 0$. Choose N so large that $|\lambda_k| < \delta$ if $k \geq N$ and also so large that

$$(26) \quad \sum_{n=r}^s \|S_n(\lambda) - P_n^{(p_n)}(\lambda)\| < \varepsilon \delta^{-(j+1)}$$

if $N \leq r < s$ and $|\lambda| \geq \delta$. Since $|\lambda_n| < \delta$, $S_n(\lambda) - P_n^{(p_n)}(\lambda)$ can be calculated from (18) and (20):

$$S_n(\lambda) - P_n^{(p_n)}(\lambda) = \sum_{k=p_n}^{\infty} \frac{(\lambda_n E_n + F_n)^k}{\lambda^{k+1}}.$$

With integration in the counterclockwise sense over the circle on which $|\lambda| = \delta$ we have

$$\varepsilon_{n,j}(\lambda_n E_n + F_n)^j = \frac{1}{2\pi i} \oint \lambda^j [S_n(\lambda) - P_n^{(p_n)}(\lambda)] d\lambda.$$

Putting $\lambda = \delta e^{i\theta}$ and using (26), we see that

$$\sum_{n=r}^s \|\varepsilon_{n,j}(\lambda_n E_n + F_n)^j\| \leq \frac{\delta^{j+1}}{2\pi} \int_0^{2\pi} \left\{ \sum_{n=r}^s \|S_n(\lambda) - P_n^{(p_n)}(\lambda)\| \right\} d\theta < \varepsilon.$$

This proves Theorem 3.

COROLLARY. *If A admits the absolute index p, the series*

$$(27) \quad \sum_{n=1}^{\infty} \|(\lambda_n E_n + F_n)^j\|$$

converges if $j \geq p$.

3. Generalization of a theorem of Berkson. In this section we show how to construct an operator which has minimal uniform index 1. The argument is modelled after the proof of a theorem due to E. R. Berkson (see §1 of Berkson [1]). Our theorem allows for multiple poles of the resolvent, whereas in Berkson's work there are simple poles only. We also discuss conditions for the operator to have minimal *absolute* index 1.

THEOREM 4. *Let $\{\mu_n\}$ be a sequence of complex numbers with $\mu_n \neq 0$ and $\mu_n \rightarrow 0$. (We do not assume that $\mu_m \neq \mu_n$ if $m \neq n$.) Let $\{P_n\}$ and $\{R_n\}$ be sequences of elements in $[X]$ satisfying the conditions*

$$(28) \quad P_n \neq 0, P_n P_m = 0 \text{ if } m \neq n, P_n^2 = P_n,$$

$$(29) \quad P_n R_n = R_n P_n = R_n, P_n R_m = R_m P_n = 0 \text{ if } m \neq n,$$

$$(30) \quad R_m R_n = 0 \text{ if } m \neq n.$$

Finally, let $\{r_n\}$ be a sequence of positive integers, and suppose that

$$(31) \quad R_n^{r_n} = 0, R_n^{r_n-1} \neq 0.$$

Now suppose that

$$(32) \quad B = \sum_{n=1}^{\infty} (\mu_n P_n + R_n),$$

it being supposed that the series converges in $[X]$. Then $\sigma(B)$ consists of 0 and the distinct points among μ_1, μ_2, \dots . The resolvent of B is given by

$$(33) \quad R_\lambda(B) = \frac{I}{\lambda} + \sum_{n=1}^{\infty} \left[T_n(\lambda) - \frac{P_n}{\lambda} \right],$$

where

$$(34) \quad T_n(\lambda) = \begin{cases} \frac{P_n}{\lambda - \mu_n} & \text{if } r_n = 1 \\ \frac{P_n}{\lambda - \mu_n} + \frac{R_n}{(\lambda - \mu_n)^2} + \dots + \frac{R_n^{r_n-1}}{(\lambda - \mu_n)^{r_n}} & \text{if } r_n > 1. \end{cases}$$

If $\delta > 0$ and if m is such that $|\mu_n| < \delta$ when $n \geq m$, the series

$$(35) \quad \sum_{n=m}^{\infty} \left[T_n(\lambda) - \frac{P_n}{\lambda} \right]$$

converges uniformly when $|\lambda| \geq \delta$. (This implies, in particular, that $R_\lambda(B)$ has a pole at μ_n , so that B is of meromorphic type.)

Proof. We write

$$B_n = \sum_{k=1}^n (\mu_k P_k + R_k).$$

If λ is fixed, different from 0, μ_1, \dots, μ_n , we define

$$R_n(\lambda) = \frac{I}{\lambda} + \sum_{k=1}^n \left[T_k(\lambda) - \frac{P_k}{\lambda} \right].$$

The first step is to prove that $\sigma(B_n)$ is the set $\{0, \mu_1, \dots, \mu_n\}$ and that $R_n(\lambda)$ is the resolvent of B_n . Direct calculations [which we omit—they are based on (28)—(30)] show that

$$R_n(\lambda)(\lambda - B_n) = (\lambda - B_n)R_n(\lambda) = I$$

if λ is different from 0, μ_1, \dots, μ_n . Much as in the proof of Lemma 1 in § 2 we prove that μ_j is an eigenvalue of B_n if $1 \leq j \leq n$. To see that 0 is also an eigenvalue, let

$$P = I - (P_1 + \dots + P_n),$$

and observe that $PP_{n+1} = P_{n+1} \neq 0$, whence $P \neq 0$. Choose $y = Px$ so

that $y \neq 0$, and observe that $B_n y = 0$. This concludes the first step in the proof.

As the second step we apply a theorem of Newburgh [4]; since $BB_n = B_n B$ and $B_n \rightarrow B$, Newburgh's theorem enables us to conclude that $\sigma(B_n)$ converges to $\sigma(B)$ in the sense that the Hausdorff distance between the sets $\sigma(B)$ and $\sigma(B_n)$ approaches zero. In the present case this means that

$$\sigma(B) = \bigcup_n \sigma(B_n) = \{0, \mu_1, \mu_2, \dots\}.$$

At the third step we obtain the formula (33) for the resolvent of B . When $\lambda \in \rho(B)$ we have

$$\begin{aligned} R_n(\lambda)(\lambda - B) &= R_n(\lambda)(\lambda - B_n) - R_n(\lambda)(B - B_n) \\ &= I - R_n(\lambda) \sum_{k=n+1}^{\infty} (\mu_k P_k + R_k), \end{aligned}$$

and a short calculation, using (28)—(30), yields

$$R_n(\lambda)(\lambda - B) = I - \frac{I}{\lambda} \sum_{k=n+1}^{\infty} (\mu_k P_k + R_k).$$

If we multiply by $R_\lambda(B)$ and transpose, we obtain

$$(36) \quad R_\lambda(B) - R_n(\lambda) = \frac{I}{\lambda} R_\lambda(B) \sum_{k=n+1}^{\infty} (\mu_k P_k + R_k).$$

We now pass to the limit as $n \rightarrow \infty$. Because of the assumed convergence of the series (32), we obtain the validity of (33) when $\lambda \in \rho(B)$.

The last step is the one concerning uniform convergence. Now, $\|\lambda^{-1}R_\lambda(B)\|$ is bounded on any subset of $\rho(B)$ in which the distance of λ from $\sigma(B)$ has a positive lower bound. Consequently, we see from (36) that the convergence of $R_n(\lambda)$ to $R_\lambda(B)$ is uniform on any such subset of $\rho(B)$. This being established, suppose $\delta > 0$, and let m be such that $|\mu_n| < \delta$ if $n \geq m$. Consider the operator

$$\sum_{n=m}^{\infty} (\mu_n P_n + R_n).$$

We can apply the results proved thus far to it. Its spectrum is the set $\{0, \mu_m, \mu_{m+1}, \dots\}$, and its resolvent is the series

$$(37) \quad \frac{I}{\lambda} + \sum_{n=m}^{\infty} \left[T_n(\lambda) - \frac{P_n}{\lambda} \right].$$

Hence, when $|\lambda| \geq \delta$, the distance of λ from the spectrum is positive, and so the series (37) converges uniformly when $|\lambda| \geq \delta$. This completes the proof of Theorem 4. (To justify the parenthetical remark at the

end of the statement of the theorem it suffices to look at formulas (34) and observe that, for fixed n , the number of indices k such that $\mu_k = \mu_n$ is finite, because of the fact that $\mu_k \rightarrow 0$.)

If we add to the assumptions in Theorem 4 the assumption that $\mu_n \neq \mu_m$ if $n \neq m$, it is evident from the conclusions of the theorem that the residue of $R_\lambda(B)$ at μ_n is P_n . It then follows from the definition in § 1 that B is of minimal uniform index 1 relative to $\{\mu_n\}$.

At this point we can state a theorem giving an alternative criterion for an operator A of meromorphic type to be of minimal uniform index 1 relative to a given enumeration of its poles.

THEOREM 5. *Let A be an operator of meromorphic type, and let the meanings of λ_n, E_n, F_n be as in § 1. Then a necessary and sufficient condition that A have minimal uniform index 1 relative to $\{\lambda_n\}$ is that the series $\sum_{n=1}^{\infty} (\lambda_n E_n + F_n)$ converge in $[X]$.*

Proof. The necessity of the condition is stated in Theorem 2. The sufficiency of the condition follows from Theorem 4, by taking $\mu_n = \lambda_n$, $P_n = E_n$, $R_n = F_n$, $r_n = q_n$ in Theorem 4. Then, the assertion about the series (35) shows that the conditions for A to admit the uniform index 1 are satisfied.

We now consider the effect of strengthening the hypotheses of Theorem 4.

THEOREM 6. *To the assumptions of Theorem 4 we add the hypothesis that the series*

$$(38) \quad \sum_{n=1}^{\infty} \|\mu_n P_n + R_n\|$$

is convergent. Then we can conclude that, if $\delta > 0$ and if m is chosen so that $|\mu_k|$ when $k \geq m$, the series

$$(39) \quad \sum_{n=m}^{\infty} \left\| T_n(\lambda) - \frac{P_n}{\lambda} \right\|$$

converges uniformly when $|\lambda| \geq \delta$. Hence, in this case, if $\mu_n \neq \mu_m$ when $n \neq m$, the operator B defined by (32) has the minimal absolute index 1.

Proof. From Lemma 1 we see that, when λ is different from 0 and μ_n ,

$$[\lambda - (\mu_n P_n + R_n)]^{-1} = \frac{I}{\lambda} + \left[T_n(\lambda) - \frac{P_n}{\lambda} \right].$$

This leads to the identity

$$T_n(\lambda) - \frac{P_n}{\lambda} \frac{1}{\lambda} \left\{ \frac{I}{\lambda} + \left[T_n(\lambda) - \frac{P_n}{\lambda} \right] \right\} (\mu_n P_n + R_n).$$

Thus

$$(40) \quad \left\| \left| T_n(\lambda) - \frac{P_n}{\lambda} \right| \right\| \leq \frac{1}{|\lambda|} \left\{ \frac{1}{|\lambda|} + \left\| \left| T_n(\lambda) - \frac{P_n}{\lambda} \right| \right\| \right\} \|\mu_n P_n + R_n\|.$$

Now, as a consequence of the proven facts about the uniform convergence of the series (35), we can assert that the factor

$$\frac{1}{|\lambda|} \left\{ \frac{1}{|\lambda|} + \left\| \left| T_n(\lambda) - \frac{P_n}{\lambda} \right| \right\| \right\}$$

is bounded uniformly with respect to n and λ if $|\lambda| \geq \delta$ and n is sufficiently large. The asserted uniform convergence of (39) now follows from (40) and the assumed convergence of (38).

Just as we got Theorem 5 from Theorem 4 and Theorem 2, so we get the following theorem from Theorem 6 and the corollary of Theorem 3.

THEOREM 7. *With A, λ_n, E_n, F_n as in § 1, A has minimal absolute index 1 if and only if the series $\sum_{n=1}^{\infty} \|\lambda_n E_n + F_n\|$ converges.*

With the assumptions of Theorem 6 it is not hard to see that B has minimal absolute index 1, even if the points μ_1, μ_2, \dots are not all distinct. Since $\mu_n \rightarrow 0$, there is only a finite set of n 's for which $\mu_n = \mu_1$. We let $\lambda_1 = \mu_1, E_1 =$ the sum of the P_n 's for which $\mu_n = \lambda_1, F_1 =$ the sum of the corresponding R_n 's. Then let λ_2 be the first μ_n different from λ_1 , and continue in the obvious way. It turns out that the pole at λ_1 is of order q_1 , where q_1 is the largest of the r_n 's corresponding to μ_n 's with $\mu_n = \lambda_1$. Likewise for the pole at λ_2 , and so on. Because of the convergence of (38) we see that

$$B = \sum_{n=1}^{\infty} (\lambda_n E_n + F_n)$$

with

$$\sum_{n=1}^{\infty} \|\lambda_n E_n + F_n\| < \infty.$$

Theorem 6 now applies, with λ_n, E_n, F_n, q_n in place of μ_n, P_n, R_n, r_n . (We leave to the reader the verification that the required conditions on E_n and F_n , corresponding to (28)—(31), are satisfied.)

4. Operators of minimal uniform index 1. We return now to the notations of §§ 1 and 2. Suppose that A is of meromorphic type, with minimal uniform index 1 relative to $\{\lambda_n\}$. By (10), (19), (21) and (22) we

see that

$$(41) \quad R_\lambda(A) = \frac{I}{\lambda} + \sum_{n=1}^{\infty} \left[S_n(\lambda) - \frac{E_n}{\lambda} \right] \\ + \sum_{j=1}^{\infty} \frac{1}{\lambda^{j+1}} \left[A^j - \sum_{n=1}^{\infty} (\lambda_n E_n + F_n)^j \right].$$

Let us define an operator B by the formula

$$(42) \quad B = \sum_{n=1}^{\infty} (\lambda_n E_n + F_n).$$

We know from Theorem 4 that $\sigma(B) = \sigma(A)$, that

$$(43) \quad R_\lambda(B) = \frac{I}{\lambda} + \sum_{n=1}^{\infty} \left[S_n(\lambda) - \frac{E_n}{\lambda} \right],$$

and that B is also an operator of meromorphic type with minimal uniform index 1 relative $\{\lambda_n\}$. We also see, from Theorem 7, that if A has minimal absolute index 1, the same is true of B .

When B is related to A by (42) under the circumstances here described, and when it turns out that $B = A$, we shall say that A is *canonical relative to* $\{\lambda_n\}$. The reader is reminded that E_n is the residue of $R_\lambda(A)$ at λ_n and that $F_n = (A - \lambda_n)E_n$.

Since $F_n E_n = F_n$, it is clear from (42) that $(B - \lambda_n)E_n = F_n$. From (43) we see that the residue of $R_\lambda(B)$ at λ_n is E_n . Hence the B given by (42) is canonical relative to $\{\lambda_n\}$, even if A is not. We may refer to B as *the canonical part of* A relative to $\{\lambda_n\}$. This naturally prompts us to investigate the nature of the operator $A - B$, which we denote by C .

THEOREM 8. *If A has minimal uniform index 1 relative to $\{\lambda_n\}$, and if B is the canonical part of A relative to $\{\lambda_n\}$, the operator $C = A - B$ is quasinilpotent [$\sigma(C) = (0)$], and $BC = CB = 0$.*

Proof. From $\lambda_n E_n + F_n = A E_n$ we see that

$$B = \sum_{n=1}^{\infty} A E_n \quad \text{and} \quad B E_m = A E_m.$$

Then (because $AB = BA$)

$$AB = \sum_{n=1}^{\infty} A^2 E_n, \quad B^2 = \sum_{n=1}^{\infty} A B E_n = AB,$$

whence $BC = CB = AB - B^2 = 0$. It now follows easily by induction that $A^n = B^n + C^n$. It also follows from (42) by induction that

$$B^j = \sum_{n=1}^{\infty} (\lambda_n E_n + F_n)^j, \quad j = 1, 2, \dots$$

On referring back to (41) and (43) we now see that

$$(44) \quad R_\lambda(A) = R_\lambda(B) + \sum_{j=1}^{\infty} \frac{C^j}{\lambda^{j+1}}.$$

The series here is convergent if $\lambda \neq 0$. This implies that C is quasinilpotent, i.e. that $\sigma(C)$ consists of 0 alone, or that $\|C^n\|^{1/n} \rightarrow 0$. This completes the proof.

The foregoing considerations raise the question as to whether the decomposition $A = B + C$ is unique, in the following sense. Is it conceivable that we could write $A = B_1 + C_1$, where C_1 is quasinilpotent, $B_1 C_1 = C_1 B_1 = 0$, and B_1 is an operator of meromorphic type which is of minimal uniform index 1 and canonical, relative to some enumeration of the poles of $R_\lambda(B_1)$, but such that B_1 is *not* the canonical part of A relative to an enumeration of the poles of $R_\lambda(A)$? The answer to this question is negative, as we now show.

THEOREM 9. *Let B_1 be of meromorphic type, of minimal uniform index 1 and canonical relative to an enumeration $\{\mu_n\}$ of the distinct poles of $R_\lambda(B_1)$. Let P_n be the residue of $R_\lambda(B_1)$ at μ_n , and let $R_n = (B_1 - \mu_n)P_n$. Let C_1 be a quasinilpotent operator such that $B_1 C_1 = C_1 B_1 = 0$. Let $A_1 = B_1 + C_1$. Then A_1 is of meromorphic type with $\sigma(A_1) = \sigma(B_1)$; A_1 has minimal uniform index 1 relative to $\{\mu_n\}$, and B_1 is the canonical part of A_1 relative to $\{\mu_n\}$.*

Proof. From $B_1 C_1 = C_1 B_1 = 0$ we deduce that $A_1 B_1 = B_1^2 = B_1 A_1$. From general spectral theory we know that μ_n is an eigenvalue of B_1 , so there exists x_n for which $B_1 x_n = \mu_n x_n \neq 0$. Then $A_1 B_1 x_n = B_1^2 x_n = B_1(\mu_n x_n) = \mu_n B_1 x_n$, so that μ_n is also an eigenvalue of A_1 . Since the spectrum is closed, it follows that $\sigma(B_1) \subset \sigma(A_1)$. Next, we show that $\rho(B_1) \subset \rho(A_1)$ and that

$$(45) \quad R_\lambda(A_1) = R_\lambda(B_1) + \sum_{n=1}^{\infty} \frac{C_1^n}{\lambda^{n+1}}$$

if $\lambda \in \rho(B_1)$. It will then follow that $\sigma(A_1) = \sigma(B_1)$.

To begin with, if $|\lambda| > \|B_1\|$ we know that

$$R_\lambda(B_1) = \sum_{n=0}^{\infty} \frac{B_1^n}{\lambda^{n+1}}.$$

It is then an easy calculation to show that

$$(46) \quad (\lambda - A_1) \left[R_\lambda(B_1) + \sum_{n=1}^{\infty} \frac{C_1^n}{\lambda^{n+1}} \right] = I$$

if $|\lambda| > \|B_1\|$. Because $\rho(B_1)$ is a connected set, and the left member of (46) is an analytic function on $\rho(B_1)$, it follows that (46) is valid for each λ in $\rho(B_1)$. The same argument applies with the order of the factors reversed. Hence we conclude that $\rho(B_1) \subset \rho(A_1)$ and that (45) holds.

From the form of (45) and the assumption on B_1 it follows that μ_n is a pole of $R_\lambda(A_1)$, the residue there being the same as the residue of $R_\lambda(B_1)$, namely P_n . In fact, we can also see that when we calculate the Laurent series for each of $R_\lambda(A_1)$ and $R_\lambda(B_1)$ in powers of $\lambda - \mu_n$, we get the same terms for both in the case of negative powers of $\lambda - \mu_n$. In particular, the coefficient of $(\lambda - \mu_n)^{-2}$ in one case is $(A_1 - \mu_n)P_n$, and in the other case it is $(B_1 - \mu_n)P_n = R_n$. Therefore $R_n = (A_1 - \mu_n)P_n$. It is now clear that A_1 is of minimal uniform index 1 relative to $\{\mu_n\}$ and that B_1 is the canonical part of A_1 .

5. Operators of minimal uniform index p . Returning once more to the notation of § 1, let us assume that A is of meromorphic type and that it has minimal uniform index p relative to $\{\lambda_n\}$. We are now interested in the case when $p > 1$. By (10), (21) and (22) we see that

$$(47) \quad R_\lambda(A) = \frac{I}{\lambda} + \frac{A}{\lambda^2} + \cdots + \frac{A^{p-1}}{\lambda^p} + \sum_{n=1}^{\infty} [S_n(\lambda) - P_n^{(p)}(\lambda)] \\ + \sum_{j=p}^{\infty} \frac{1}{\lambda^{j+1}} \left[A^j - \sum_{n=1}^{\infty} (\lambda_n E_n + F_n)^j \right].$$

This formula takes an especially simple form if

$$(48) \quad A^p = \sum_{n=1}^{\infty} (\lambda_n E_n + F_n)^p,$$

because

$$\lambda_n E_n + F_n = A E_n, \quad (\lambda_n E_n + F_n)^j = A^j E_n,$$

so that

$$A^p = \sum_{n=1}^{\infty} A^p E_n,$$

whence it follows that

$$A^j = \sum_{n=1}^{\infty} A^j E_n \quad \text{if } j \geq p,$$

and so (47) becomes, in this special case,

$$(49) \quad R_\lambda(A) = \frac{I}{\lambda} + \frac{A}{\lambda^2} + \cdots + \frac{A^{p-1}}{\lambda^p} + \sum_{n=1}^{\infty} [S_n(\lambda) - P_n^{(p)}(\lambda)].$$

When A is of minimal uniform index p relative to $\{\lambda_n\}$, and when

(48) holds, we say that A is canonical of order p relative to $\{\lambda_n\}$. In § 7 we discuss an unsolved problem involving the concept of a canonical operator of order p , for the case in which A itself is not canonical.

We now present a generalization of Theorem 5.

THEOREM 10. *Let A be of meromorphic type, and let λ_n, E_n, F_n be as in § 1. Suppose that p is a positive integer, and suppose that the series $\sum_{n=1}^{\infty} (\lambda_n E_n + F_n)^p$ is convergent in $[X]$. Then A admits the uniform index p relative to $\{\lambda_n\}$. Hence (see Theorem 2), in order that A have minimal uniform index p relative to $\{\lambda_n\}$, it is necessary and sufficient that $\sum_{n=1}^{\infty} (\lambda_n E_n + F_n)^p$ converge in $[X]$ and that $\sum_{n=1}^{\infty} (\lambda_n E_n + F_n)^{p-1}$ not converge in $[X]$.*

Proof. Our argument is valid if $p \geq 1$. Hence this theorem includes Theorem 5; the proof is different from that of Theorem 5, however, even when $p = 1$. The argument hinges on two identities. Let $H_n = \lambda_n E_n + F_n$. Then

$$(50) \quad S_n(\lambda) - P_n^{(p)}(\lambda) = \lambda^{-p} R_\lambda(H_n) H_n^p$$

if $\lambda \neq 0$ and $\lambda \neq \lambda_n$. The other identity is

$$(51) \quad R_\lambda(H_n) H_n^p = R_\lambda(A) H_n^p;$$

it is valid if $\lambda \in \rho(A)$. To prove (50) we may assume $|\lambda| > |\lambda_n|$. When (50) has been proved under this condition the general assertion about the validity of (50) follows by the principle of analytic continuation, because the expressions on each side of the equality in (50) are analytic in λ except at 0 and λ_n . (We see by Lemma 1 in § 2 that $\sigma(H_n) = \{0, \lambda_n\}$.) Now, from (18) and (20) we see that

$$S_n(\lambda) - P_n^{(p)}(\lambda) = \sum_{k=p}^{\infty} \frac{H_n^k}{\lambda^{k+1}} \quad \text{if } |\lambda| > |\lambda_n|.$$

From the Neumann series for $R_\lambda(H_n)$ we see that

$$\lambda^{-p} R_\lambda(H_n) H_n^p = \frac{H_n^p}{\lambda^p} \sum_{j=0}^{\infty} \frac{H_n^j}{\lambda^{j+1}} = \sum_{k=p}^{\infty} \frac{H_n^k}{\lambda^{k+1}}.$$

Thus (50) is proved.

In proving (51), because of the principle of analytic continuation, we may restrict ourselves to values of λ so large that $\|A - \lambda_n\| < |\lambda - \lambda_n|$. With this restriction, noting that $\lambda - A = (\lambda - \lambda_n) - (A - \lambda_n)$, we can write

$$(52) \quad R_\lambda(A) = \sum_{k=0}^{\infty} \frac{(A - \lambda_n)^k}{(\lambda - \lambda_n)^{k+1}}.$$

Now from $F_n = (A - \lambda_n)E_n$, $H_n = \lambda_n E_n + F_n$, and $E_n^2 = E_n$, we deduce that $H_n F_n = (A - \lambda_n)H_n$. It then follows readily that $H_n^p F_n^k = (A - \lambda_n)^k H_n^p$ for all positive integers p and k . Since $F_n^k = 0$ if $k \geq q_n$, it follows from (52) that

$$(53) \quad R_\lambda(A)H_n^p = \sum_{k=0}^{q_n-1} \frac{H_n^p F_n^k}{(\lambda - \lambda_n)^{k+1}}.$$

On the other hand, we know from (15) that

$$R_\lambda(H_n) = \frac{I - E_n}{\lambda} + \frac{E_n}{\lambda - \lambda_n} + \frac{F_n}{(\lambda - \lambda_n)^2} + \cdots + \frac{F_n^{q_n-1}}{(\lambda - \lambda_n)^{q_n}};$$

because $H_n^p E_n = H_n^p$, we see from this and (53) that (51) holds.

We now turn to the proof of Theorem 10. We have to show that the convergence of the series $\sum_{n=1}^{\infty} H_n^p$ implies that the series

$$\sum_{n=m}^{\infty} [S_n(\lambda) - P_n^{(p)}(\lambda)]$$

converges uniformly when $|\lambda| \geq \delta$, provided merely that $\delta > 0$ and that m is chosen in such a way that $|\lambda_k| < \delta$ when $k \geq m$. When δ and m are specified, there may perhaps be a finite set of λ_i 's for which $|\lambda_i| \geq \delta$. We can diminish δ slightly, if necessary, so as to have $|\lambda_i| > \delta$ for all these λ_i 's and still have $|\lambda_k| < \delta$ if $k \geq m$. If we give the proof with this modified δ , that is certainly sufficient. Choose a positive number α so small that, if $|\lambda_i| > \delta$, the closed circular disk of radius α and center λ_i lies entirely outside the circle on which $|\lambda| = \delta$, and so that any two such disks do not touch or overlap. Let $E(\delta, \alpha)$ be the set of λ 's such that $|\lambda| \geq \delta$ and $|\lambda - \lambda_i| = \alpha$ if $|\lambda_i| > \delta$. There is some constant M such that $\|\lambda^{-p} R_\lambda(A)\| \leq M$ if $\lambda \in E(\delta, \alpha)$. Now, suppose that $\varepsilon > 0$. By hypothesis we can choose an integer N so large that

$$M \left\| \sum_{n=j}^k H_n^p \right\| < \varepsilon$$

if $N \leq j \leq k$. By (50) and (51) we can write

$$\sum_{n=j}^k [S_n(\lambda) - P_n^{(p)}(\lambda)] = \lambda^{-p} R_\lambda(A) \sum_{n=j}^k H_n^p,$$

provided that $\lambda \in \rho(A)$. Therefore certainly

$$(54) \quad \left\| \sum_{n=j}^k [S_n(\lambda) - P_n^{(p)}(\lambda)] \right\| < \varepsilon$$

if $N \leq j \leq k$, provided that $\lambda \in E(\delta, \alpha)$. In fact, however, (54) holds whenever $|\lambda| \geq \delta$; that is, it holds even if $|\lambda| \geq \delta$, $|\lambda_i| > \delta$, and $|\lambda - \lambda_i| < \alpha$. For, the function

$$\sum_{n=j}^k [S_n(\lambda) - P_n^{(p)}(\lambda)]$$

is holomorphic in λ when $|\lambda - \lambda_i| \leq \alpha$, and its norm is less than ε when $|\lambda - \lambda_i| = \alpha$; therefore the norm is also less than ε when $|\lambda - \lambda_i| < \alpha$, by the maximum modulus theorem, which is valid for analytic functions with values in a complex Banach space (see Taylor [5], Theorem 4. 42-A, or Hille-Phillips [3], p. 100). This completes the proof of Theorem 10.

In the next theorem we present information about the operator A^p under the assumption that A has minimal uniform index p .

THEOREM 11. *Suppose that A is of meromorphic type and of minimal uniform index p relative to $\{\lambda_n\}$, with $p > 1$. (We adopt the notation of § 1 as regards λ_n, E_n, F_n .) Define operators L, M by the formulas*

$$(55) \quad L = \sum_{n=1}^{\infty} (\lambda_n E_n + F_n)^p, \quad M = A^p - L.$$

Let

$$(57) \quad G_n = (\lambda_n E_n + F_n)^p - \lambda_n^p E_n,$$

$$(57) \quad V_n(\lambda) = \begin{cases} \frac{E_n}{\lambda - \lambda_n^p} \text{ if } q_n = 1 \\ \frac{E_n}{\lambda - \lambda_n^p} + \frac{G_n}{(\lambda - \lambda_n^p)^2} + \dots + \frac{G_n^{q_n-1}}{(\lambda - \lambda_n^p)^{q_n}}. \end{cases}$$

Then L is an operator of meromorphic type with $\sigma(L)$ consisting of $\lambda_1^p, \lambda_2^p, \dots$ and 0. The resolvent of L is

$$(58) \quad R_\lambda(L) = \frac{I}{\lambda} + \sum_{n=1}^{\infty} \left[V_n(\lambda) - \frac{E_n}{\lambda} \right].$$

If $\delta > 0$ and if m is such that $|\lambda_n^p| < \delta$ when $n \geq m$, then the series

$$\sum_{n=m}^{\infty} \left[V_n(\lambda) - \frac{E_n}{\lambda} \right]$$

converges uniformly when $|\lambda| \geq \delta$. The operator M is quasinilpotent, and $LM = ML = 0$. Finally, if all the points $\lambda_1^p, \lambda_2^p, \dots$ are distinct, A^p has minimal uniform index 1 relative to $\{\lambda_n^p\}$, and L is the canonical part of A^p relative to $\{\lambda_n^p\}$.

Proof. By expanding $(\lambda_n E_n + F_n)^p$, we see from (56) that G_n can be expressed in the form $G_n = F_n J_n = J_n F_n$, where $J_n \in [X]$. It is then easy to see that $E_n G_n = G_n E_n = G_n$ and $G_n^{q_n} = 0$. We shall show that $G_n^{q_n-1} \neq 0$ if $q_n > 1$. In fact, from (56) we can write $G_n = p \lambda_n^{p-1} F_n + F_n^2 K_n$, where $K_n \in [X]$ and $F_n K_n = K_n F_n$. Thus $G_n^{q_n-1} = 0$ would imply

$$F_n^{q_n-1} [p \lambda_n^{p-1} I + F_n K_n]^{q_n-1} = 0.$$

Since $F_n^{q_n} = 0$ and $p\lambda_n^{p-1} \neq 0$, this would imply $F_n^{q_n-1} = 0$, contrary to fact.

We can now apply Theorem 4, taking $\mu_n = \lambda_n^p$, $P_n = E_n$, $R_n = G_n$, $r_n = q_n$. From (55) and (56) we see that

$$L = \sum_{n=1}^{\infty} (\lambda_n^p E_n + G_n);$$

therefore we obtain at once our assertions about $\sigma(L)$ and $R_\lambda(L)$. The convergence behavior of the series (58) insures that $R_\lambda(L)$ has a pole at λ_n^p .

Now consider the operator M defined in (55). From (2) we see that $AE_n = \lambda_n E_n + F_n$, so we can write

$$L = \sum_{n=1}^{\infty} A^n E_n,$$

from which it follows that $LA^p = A^p L = L^2$. Then $LM = L(A^p - L) = 0$, and likewise $ML = 0$. It follows that $A^{np} = L^n + M^n$, $n = 1, 2, \dots$. To prove that M is quasinilpotent we shall show that the series

$$\sum_{n=1}^{\infty} \frac{M^n}{\lambda^{n+1}}$$

converges if $\lambda \neq 0$. Now, from (55) it follows readily that

$$L^j = \sum_{n=1}^{\infty} (\lambda_n E_n + F_n)^{j_p} \quad \text{if } j \geq 1.$$

Hence, what we wish to prove is that the series

$$(59) \quad \sum_{j=1}^{\infty} \frac{A^{j_p} - \sum_{n=1}^{\infty} (\lambda_n E_n + F_n)^{j_p}}{\lambda^{j+1}}$$

is convergent if $\lambda \neq 0$. We know from (47) that the series

$$\sum_{k=p}^{\infty} \frac{A^k - \sum_{n=1}^{\infty} (\lambda_n E_n + F_n)^k}{\lambda^{j+1}}$$

converges, and from this we can deduce the convergence of the series in (59). If $\lambda \neq 0$, choose a circle of radius ε centered at 0, where $0 < \varepsilon, \varepsilon^p < |\lambda|$. Then, integration around this circle in the counterclockwise sense yields

$$\frac{1}{2\pi i} \oint \frac{dt}{(\lambda - t^p)t^{k+1}} = \begin{cases} \frac{1}{\lambda^{k+1}} & \text{if } k = jp, j = 1, 2, \dots \\ 0 & \text{if } k \text{ is not a multiple of } p. \end{cases}$$

Therefore, if

$$D_k = A^k - \sum_{n=1}^{\infty} (\lambda_n E_n + F_n)^k,$$

we have

$$\frac{1}{2\pi i} \oint \frac{1}{\lambda - t^p} \left(\sum_{k=p}^{\infty} \frac{D_k}{t^{k+1}} \right) dt = \sum_{j=1}^{\infty} \frac{D_{jp}}{\lambda^{j+1}},$$

the last series on the right being convergent. By comparison with (59), we see that we have finished the proof that M is quasinilpotent.

It is clear from the situation that L is of minimal uniform index 1 and that L is canonical relative to $\{\lambda_n^p\}$ when the points $\lambda_1^p, \lambda_2^p, \dots$ are all distinct. The final assertion of Theorem 11 is then a consequence of Theorem 9.

6. Operators of minimal absolute index p . Suppose that A is of meromorphic type and of minimal absolute index p . We follow the notation of § 1. It is then certainly true that A admits the uniform index p relative to $\{\lambda_n\}$, although p is not necessarily the minimal uniform index. Let L, M , and G_n be defined by (55) and (56). By the corollary of Theorem 3 we know that the series

$$\sum_{n=1}^{\infty} \|\lambda_n^p E_n + G_n\|$$

is convergent. From the remarks at the end of § 3 we see that L has minimal absolute index 1, regardless of whether or not the point $\lambda_1^p, \lambda_2^p, \dots$ are all distinct. Under our present assumptions the operator M defined in (55) is quasinilpotent, for the reasoning about this in the proof of Theorem 11 is all applicable. It is obvious that L is of minimal uniform index 1 and canonical relative to any enumeration of the distinct poles of $R_\lambda(L)$. It then follows as in Theorem 9 and its proof that $\sigma(A^p) = \sigma(L)$ and that

$$(60) \quad R_\lambda(A^p) = R_\lambda(L) + \sum_{n=1}^{\infty} \frac{M_n}{\lambda^{n+1}} \quad \text{if } \lambda \in \sigma(A^p).$$

From (60) and the definitions in § 1 we can then see that A^p is of meromorphic type and minimal absolute index 1. We state all this formally:

THEOREM 12. *If A is of meromorphic type and minimal absolute index p , A^p is of meromorphic type and minimal absolute index 1.*

The following theorem is analogous to Theorem 10.

THEOREM 13. *Let A be of meromorphic type, and let the notation of § 1 be adopted. Then A admits the absolute index p if and only if the series*

$$(61) \quad \sum_{n=1}^{\infty} \|(\lambda_n E_n + F_n)^p\|$$

is convergent. Hence, A has minimal absolute index p (a positive integer) if and only if

$$\sum_{n=1}^{\infty} \|(\lambda_n E_n + F_n)^p\| < \infty \text{ and } \sum_{n=1}^{\infty} \|(\lambda_n E_n + F_n)^{p-1}\| = \infty .$$

Proof. In view of the corollary of Theorem 3, we have only to prove that A admits the absolute index p when the series (61) is convergent.

We begin by calling attention once more to (50) and (51), and to the notation $H_n = \lambda_n E_n + F_n$. We are supposing that

$$\sum_{n=1}^{\infty} \|H_n^p\| < \infty .$$

Suppose $\delta > 0$, and let m be such that $|\lambda_k| < \delta$ if $k \geq m$. We assert the existence of a positive number C that

$$(62) \quad \|R_{\lambda}(H_n)H_n^p\| \leq C \|H_n^p\|$$

if $|\lambda| \geq \delta$ and $n \geq m$. Once this is proved, the fact that

$$\sum_{n=m}^{\infty} \|S_n(\lambda) - P_n^{(p)}(\lambda)\|$$

converges uniformly when $|\lambda| \geq \delta$ will follow from (50) and (62), and our proof will be accomplished. There is only a finite set of k 's for which $|\lambda_k| \geq \delta$. Choose $\varepsilon > 0$ so small that $\lambda \in \rho(A)$ if $0 < |\lambda - \lambda_k| \leq \varepsilon$ for these k 's. We know that $\|R_{\lambda}(A)\|$ is bounded by some positive constant C_0 when λ is at a distance greater than or equal to ε from $\sigma(A)$. Consider the situation if $0 < |\lambda - \lambda_k| \leq \varepsilon$ and $|\lambda_k| \geq \delta$. We know there is a function of λ with values in $[X]$, call it f_k , such that f_k is analytic when $|\lambda - \lambda_k| \leq \varepsilon$ and

$$R_{\lambda}(A) = f_k(\lambda) + S_k(\lambda)$$

if $0 < |\lambda - \lambda_k| \leq \varepsilon$. Now $H_n^p E_k = H_n^p F_k = 0$ if $n \neq k$. Therefore, in view of the definition of $S_k(\lambda)$ in section 1, $R_{\lambda}(A)H_n^p = f_k(\lambda)H_n^p$, and we see from (51) that $\|R_{\lambda}(H_n)H_n^p\| = \|f_k(\lambda)H_n^p\|$ will have a bound $C_k \|H_n^p\|$ if $|\lambda - \lambda_k| \leq \varepsilon$. If we choose C at least as large as C_0 and the largest of the finite set of C_k 's, (62) will hold. This ends the proof.

7. An open problem. Suppose that A is of meromorphic type and of minimal uniform index p relative to $\{\lambda_n\}$, where $p > 1$. Suppose further that A is not canonical of order p relative to $\{\lambda_n\}$ (see § 5). Is there in this case some uniquely determined operator B which deserves

to be called the canonical part of A relative to $\{\lambda_n\}$? For the case $p = 1$ we know that this question has an affirmative answer. The details are given in § 4; the required B is given by (42). For the case $p > 1$, however, we have been unable to answer the question. We are even unable to say with certainty what properties should characterize the required B . We would certainly want B to have the same spectrum as A , and for each λ_n we would want the Laurent expansion of $R_\lambda(B)$ in powers of $\lambda - \lambda_n$ to consist of $S_n(\lambda)$ plus a series of nonnegative powers of $\lambda - \lambda_n$. We would also require that

$$(63) \quad B^p = \sum_{n=1}^{\infty} (\lambda_n E_n + F_n)^p .$$

The requirements thus far stated would be sufficient to imply that B is of minimal uniform index p and canonical of order p relative to $\{\lambda_n\}$, with

$$(64) \quad R_\lambda(B) = \frac{I}{\lambda} + \frac{B}{\lambda^2} + \dots + \frac{B^{p-1}}{\lambda^p} + \sum_{n=1}^{\infty} [S_n(\lambda) - P_n^{(p)}(\lambda)]$$

when $\lambda \in \rho(B)$. It is not clear, however that there is at most one B fulfilling all these conditions. Nor is there, so far as we can see, any method of proving the existence of a B with these properties.

One possible starting point might be to define

$$B_n = \sum_{k=1}^n (\lambda_k E_k + F_k) .$$

It may be proved that $\sigma(B_n) = \{0, \lambda_1, \dots, \lambda_n\}$ and that

$$R_\lambda(B_n) = \frac{I}{\lambda} + \frac{B_n}{\lambda^2} + \dots + \frac{B_n^{p-1}}{\lambda^p} + \sum_{k=1}^n [S_k(\lambda) - P_n^{(p)}(\lambda)] .$$

But, if $p > 1$, B_n has no limit in the sense of convergence in $[X]$. It is true that

$$B_n^p \rightarrow \sum_{k=1}^{\infty} (\lambda_k E_k + F_k)^p .$$

What appears to be needed is some way of assigning a limit B to the sequence $\{B_n\}$ in some generalized sense, and then proving that this limit B is the unique operator with the required properties. In particular cases it may (and does) happen that we can define B by $Bx = \lim_{n \rightarrow \infty} B_n x$ for each x . But we have no general theory.

8. Examples. In this concluding section we show how to construct an operator A with minimal uniform index 1 or 2 such that A either admits no absolute index at all, or admits an arbitrarily prescribed minimal absolute index (greater than or equal to the minimal uniform

index, of course).

We also show how to construct an operator A with arbitrarily preassigned minimal uniform index k .

In both examples we consider operators A acting in the space l^1 . If $x = (\xi_1, \xi_2, \dots)$ and $y = (\eta_1, \eta_2, \dots)$ are points of l^1 with $y = Ax$, we take the defining equations to be


$$\eta_i = \sum_{j=1}^{\infty} \alpha_{ij} \xi_j, \quad i = 1, 2, \dots$$

Thus A is represented by an infinite matrix. For operators acting in l^1 the norm is

$$\|A\| = \sup_j \sum_{i=1}^{\infty} |\alpha_{ij}|.$$

EXAMPLE 1

For our first example we take the matrix representation of A to have the form



where the only nonzero matrix entries are in the square blocks $B_1, C_1, B_2, C_2, B_3, \dots$ down the main diagonal. The block B_n is to be a two-by-two block

$$B_n : \begin{bmatrix} \lambda_n & 0 \\ \mu_n & \lambda_n \end{bmatrix},$$

and the block C_n is to have r_n rows and columns, with the only nonzero elements 1's in the subdiagonal:

$$C_n : \begin{bmatrix} 0 & \dots & 0 \\ 1 & 0 & \dots & 0 \\ \vdots & & & \\ 0 & \dots & 1 & 0 \end{bmatrix}.$$

We require $0 \leq r_n$, $\sup r_n = r < \infty$. If $r_n = 0$ the block C_n is to be

nonexistent, and if $r_n = 1$, C_n is to be a block with the single entry 0. We assume $\{\lambda_n\}$ is a sequence of distinct nonzero numbers such that $\lambda_n \rightarrow 0$; $\{\mu_n\}$ can be any bounded sequence.

It is not difficult to compute $R_\lambda(A)$. Each point λ_n is a pole of the second order and all other nonzero points are in $\rho(A)$. The block corresponding to B_n in the matrix representing $R_\lambda(A)$ is

$$\begin{array}{|c|c|} \hline \frac{1}{\lambda - \lambda_n} & 0 \\ \hline \frac{\mu_n}{(\lambda - \lambda_n)^2} & \frac{1}{\lambda - \lambda_n} \\ \hline \end{array}$$

The block corresponding to C_n is

$$\begin{array}{|cccc|} \hline \frac{1}{\lambda} & 0 & \dots & 0 \\ \hline \frac{1}{\lambda^2} & \frac{1}{\lambda} & & 0 \\ \hline \vdots & & & 0 \\ \hline \frac{1}{\lambda^{r_n}} & \dots & & \frac{1}{\lambda} \\ \hline \end{array}$$

The matrices representing E_n and F_n , respectively, have the blocks

$$\begin{array}{|c|c|} \hline 1 & 0 \\ \hline 0 & 1 \\ \hline \end{array} \quad \begin{array}{|c|c|} \hline 0 & 0 \\ \hline \mu_n & 0 \\ \hline \end{array}$$

in place of B_n , and all other entries are 0.

It is now easy to see that, if $l < m$,

$$\left\| \sum_{n=l}^m (\lambda_n E_n + F_n)^p \right\| = \sup_{l \leq n \leq m} [|\lambda_n|^p + p|\lambda_n^{p-1}\mu_n|].$$

Consequently, A admits the uniform index 2 relative to $\{\lambda_n\}$. It admits the uniform index 1 if and only if $\mu_n \rightarrow 0$.

We can also see that

$$\|(\lambda_n E_n + F_n)^p\| = |\lambda_n|^p + p|\lambda_n^{p-1}\mu_n|.$$

Therefore A admits the absolute index p if and only if

$$\sum_{n=1}^{\infty} [|\lambda_n|^p + p|\lambda_n^{p-1}\mu_n|] < \infty .$$

From this discussion we see that we can arrange for A to have any prescribed minimal absolute index (greater than or equal to the minimal uniform index relative to $\{\lambda_n\}$). Or, we can also arrange for A not to admit any absolute index at all. For instance, if $\lambda_n = [\log(n + 1)]^{-1}$, and $\mu_n = n^{-1}$, the minimal uniform index is 1 and there is no absolute index. If $k > 1$, $\lambda_n = n^{-k/(k-1)^2}$, and $\mu_n = 1$, the minimal uniform index is 2 and the minimal absolute index is k .

Let us return to the general case of this example. Suppose we have the situation where the minimal uniform index is 2. The operator A is not canonical, but we can express it in the form $A = B + C$, where B satisfies (63) and (64) with $p = 2$, and C is nilpotent (of order r), with $BC = CB = 0$. We can define B as the operator whose matrix representation is like the matrix representation of A except that the blocks C_1, C_2, \dots have all been replaced by blocks of zeros. It can also be seen that

$$Bx = \sum_{n=1}^{\infty} (\lambda_n E_n + F_n)x$$


for each x , so that

$$B = \sum_{n=1}^{\infty} (\lambda_n E_n + F_n) ,$$

with convergence in the strong (instead of the uniform) operator topology.

EXAMPLE 2

For this example we again use the matrix representation of A and a scheme of blocks:



The form of the block A_n is to be

$$A_n : \begin{bmatrix} \lambda_{2n-1} & 0 \\ \mu_n & \lambda_{2n} \end{bmatrix} .$$

We assume that $\{\lambda_n\}$ is a sequence of distinct nonzero numbers such

that $\lambda_n \rightarrow 0$. For $\{\mu_n\}$ we take any bounded sequence. This time all the poles are simple; the block corresponding to A_n in the matrix representation of $R_\lambda(A)$ is

$$\begin{array}{cc} \frac{1}{\lambda - \lambda_{2n-1}} & 0 \\ \frac{\mu_n}{(\lambda - \lambda_{2n-1})(\lambda - \lambda_{2n})} & \frac{1}{\lambda - \lambda_{2n}} \end{array}$$

The matrix representing E_{2n-1} has just one nonzero block; it is in the position occupied by A_n , and it is

$$\begin{array}{cc} 1 & 0 \\ \alpha_n & 0 \end{array},$$

where

$$(65) \quad \alpha_n = \frac{\mu_n}{\lambda_{2n-1} - \lambda_{2n}}.$$

The matrix representing E_{2n} is of the same type, but its sole nonzero block is

$$\begin{array}{cc} 0 & 0 \\ -\alpha_n & 1 \end{array}.$$

From these facts we can readily see that A admits the absolute index p if and only if

$$\sum_{n=1}^{\infty} |\lambda_{2n-1}|^p (1 + |\alpha_n|) < \infty$$

and

$$\sum_{n=1}^{\infty} |\lambda_{2n}|^p \beta_n < \infty,$$

where $\beta_n = \max(1, |\alpha_n|)$. Equivalently, the condition is that each of the following four series must converge:

$$(66) \quad \sum_{n=1}^{\infty} |\lambda_{2n-1}|^p, \quad \sum_{n=1}^{\infty} |\lambda_{2n}|^p, \quad \sum_{n=1}^{\infty} |\lambda_{2n-1}|^p |\alpha_n|, \quad \sum_{n=1}^{\infty} |\lambda_{2n}|^p |\alpha_n|.$$

In investigating the question of a uniform index for A relative to $\{\lambda_n\}$, it is convenient to begin by calculating A^p . The matrix representation of A^p has the same form as that of A , with the block A_n replaced by

$$\begin{array}{cc} \lambda_{2n-1}^p & 0 \\ \mu_n \frac{\lambda_{2n-1}^p - \lambda_{2n}^p}{\lambda_{2n-1} - \lambda_{2n}} & \lambda_{2n}^p \end{array}$$

This block is the same as the sole nonzero block in the matrix representation of

$$\lambda_{2n-1}^p E_{2n-1} + \lambda_{2n}^p E_{2n}.$$

It is then easy to see that the matrix representation of

$$A^p = \sum_{k=1}^{2n} \lambda_k^p E_k$$

is derived from the matrix representation of A^p simply by putting blocks of zeros in place of the first n blocks down the diagonal. From this it may be proved that A admits the uniform index p relative to $\{\lambda_n\}$ if and only if

$$A^p = \sum_{n=1}^{\infty} \lambda_n^p E_n,$$

and that this occurs if and only if

$$(67) \quad \|\lambda_{2n-1}^p E_{2n-1} + \lambda_{2n}^p E_{2n}\| \rightarrow 0$$

and

$$(68) \quad \|\lambda_{2n-1}^p E_{2n-1}\| \rightarrow 0$$

as $n \rightarrow \infty$.

Condition (67) is equivalent to

$$\max \{|\lambda_{2n-1}^p| + |\lambda_{2n-1}^p - \lambda_{2n}^p| |\alpha_n|, |\lambda_{2n}^p|\} \rightarrow 0,$$

and condition (68) is equivalent to

$$|\lambda_{2n-1}^p| (1 + |\alpha_n|) \rightarrow 0.$$

Since we assumed at the outset that $\lambda_n \rightarrow 0$, it is now easy to see that A admits the uniform index p relative to $\{\lambda_n\}$ if and only if

$$(69) \quad \frac{|\lambda_{2n-1}^p| |\mu_n|}{|\lambda_{2n-1} - \lambda_{2n}|} \rightarrow 0 \quad \text{and} \quad \frac{|\lambda_{2n}^p| |\mu_n|}{|\lambda_{2n-1} - \lambda_{2n}|} \rightarrow 0$$

If we take $\lambda_n = n^{-1}$, it turns out that A admits the uniform index p relative to $\{\lambda_n\}$ if and only if $\mu_n n^{2-p} \rightarrow 0$, and A admits the absolute index p if and only if

$$\sum_{n=1}^{\infty} \frac{|\mu_n|}{n^{p-2}} < \infty .$$

In particular, since $\{\mu_n\}$ is bounded, A certainly admits the uniform index 3 and the absolute index 4, but these indices are not necessarily minimal.

If we take $\lambda_n = n^{-1/k}$ (with k a positive integer) a simple calculation with the binomial series shows that each of the expressions

$$\frac{|\lambda_{2n-1}|^p}{|\lambda_{2n-1} - \lambda_{2n}|} \text{ and } \frac{|\lambda_{2n}|^p}{|\lambda_{2n-1} - \lambda_{2n}|}$$

is asymptotically equivalent to

$$k \left(\frac{1}{2n-1} \right)^{(p-1-k)/k} .$$

Consequently, A admits the uniform index p relative to $\{\lambda_n\}$ if and only if

$$n^{(k+1-p)/k} \mu_n \rightarrow 0 .$$

With $\mu_n = n^{-2/k}$, we can see that A has minimal uniform index k relative to $\{\lambda_n\}$. These same choices of λ_n and μ_n give A the minimal absolute index $2k$.

REFERENCES

1. Earl Berkson, *Sequel to a paper of A. E. Taylor*, Pacific J. Math., **10** (1960), 767-776.
2. John Derr, *Multiple poles of the resolvent for operators of finite Mittag-Leffler type*, Doctoral dissertation, University of California, Los Angeles, 1962.
3. E. Hille and R. S. Phillips, *Functional analysis and semi-groups* (revised edition), American Mathematical Society Colloquium Publications, vol. 31, New York, 1957.
4. J. D. Newburgh, *The variation of spectra*, Duke Math. J., **18** (1951), 165-176.
5. Angus E. Taylor, *Introduction to functional analysis*, John Wiley, New York, 1958.
6. ———, *Mittag-Leffler expansions and spectral theory*, Pacific J. Math., **10** (1960), 1049-1056.
7. ———, *Spectral theory and Mittag-Leffler type expansions of the resolvent*, Proceedings of the International Symposium on Linear Spaces, The Hebrew University, Jerusalem, 1960.

THE RAND CORPORATION
 THE UNIVERSITY OF CALIFORNIA, LOS ANGELES

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RALPH S. PHILLIPS

Stanford University
Stanford, California

M. G. ARSOVE

University of Washington
Seattle 5, Washington

A. L. WHITEMAN

University of Southern California
Los Angeles 7, California

LOWELL J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

T. M. CHERRY

D. DERRY

M. OHTSUKA

H. L. ROYDEN

E. SPANIER

E. G. STRAUS

F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Jonathan L. Alperin, <i>Groups with finitely many automorphisms</i>	1
Martin Arthur Arkowitz, <i>The generalized Whitehead product</i>	7
John D. Baum, <i>Instability and asymptoticity in topological dynamics</i>	25
William Aaron Beyer, <i>Hausdorff dimension of level sets of some Rademacher series</i>	35
Frank Herbert Brownell, III, <i>A note on Cook's wave-matrix theorem</i>	47
Gulbank D. Chakerian, <i>An inequality for closed space curves</i>	53
Inge Futtrup Christensen, <i>Some further extensions of a theorem of Marcinkiewicz</i>	59
Charles Vernon Coffman, <i>Linear differential equations on cones in Banach spaces</i>	69
Eckford Cohen, <i>Arithmetical notes. III. Certain equally distributed sets of integers</i>	77
John Irving Derr and Angus E. Taylor, <i>Operators of meromorphic type with multiple poles of the resolvent</i>	85
Jacob Feldman, <i>On measurability of stochastic processes in products space</i>	113
Robert S. Freeman, <i>Closed extensions of the Laplace operator determined by a general class of boundary conditions, for unbounded regions</i>	121
Robert E. Fullerton, <i>Geometric structure of absolute basis systems in a linear topological space</i>	137
Dieter Gaijer, <i>On conformal mapping of nearly circular regions</i>	149
Andrew Mattei Gleason and Hassler Whitney, <i>The extension of linear functionals defined on H^∞</i>	163
Seymour Goldberg, <i>Closed linear operators and associated continuous linear operators</i>	183
Basil Gordon, Aviezri Siegmund Fraenkel and Ernst Gabor Straus, <i>On the determination of sets by the sets of sums of a certain order</i>	187
Branko Grünbaum, <i>The dimension of intersections of convex sets</i>	197
Paul Daniel Hill, <i>On the number of pure subgroups</i>	203
Robert Peter Holten, <i>Generalized Goursat problem</i>	207
Alfred Horn, <i>Eigenvalues of sums of Hermitian matrices</i>	225
Henry C. Howard, <i>Oscillation and nonoscillation criteria for $y''(x) + f(y(x))p(x) = 0$</i>	243
Taqdir Husain, <i>S-spaces and the open mapping theorem</i>	253
Richard Eugene Isaac, <i>Markov processes and unique stationary probability measures</i>	273
John Rolfe Isbell, <i>Supercomplete spaces</i>	287
John Rolfe Isbell, <i>On finite-dimensional uniform spaces. II</i>	291
N. Jacobson, <i>A note on automorphisms of Lie algebras</i>	303
Antoni A. Kosinski, <i>A theorem on families of acyclic sets and its applications</i>	317
Marvin David Marcus and H. Minc, <i>The invariance of symmetric functions of singular values</i>	327
Ralph David McWilliams, <i>A note on weak sequential convergence</i>	333
John W. Milnor, <i>On axiomatic homology theory</i>	337
Victor Julius Mizel and Malempati Madhusudana Rao, <i>Nonsymmetric projections in Hilbert space</i>	343
Calvin Cooper Moore, <i>On the Frobenius reciprocity theorem for locally compact groups</i>	359
Donald J. Newman, <i>The Gibbs phenomenon for Hausdorff means</i>	367
Jack Segal, <i>Convergence of inverse systems</i>	371
Józef Siciak, <i>On function families with boundary</i>	375
Hyman Joseph Zimmerberg, <i>Two-point boundary conditions linear in a parameter</i>	385