

Pacific Journal of Mathematics

GROUP MEMBERSHIP IN SEMIGROUPS

HUBERT COLLINGS KENNEDY

GROUP MEMBERSHIP IN SEMIGROUPS

HUBERT C. KENNEDY

An element a of a semigroup S is said to be a *group member* in S if there is a subgroup G of S which contains a . This notion was introduced in 1909 by Arthur Ranum in discussing group membership in certain sets of matrices. In 1927 he extended the notion to group membership in finite dimensional algebras. Using somewhat different methods, the subject was revived separately by two sets of authors, W. E. Barnes and H. Schneider in 1957, and H. K. Farahat and L. Mirsky in 1956 and 1958. They discussed the notion of group membership in polynomials in an element algebraic over a field and in rings of various types. With regard to rings, we say that an element a is a group member in a ring R if a is a group member of the semigroup formed by elements of R and its multiplicative operation. Barnes and Schneider posed the following question: If S is a subring of a ring R and an element a of S is a group member in R , under what conditions is it a group member in S [1, p. 168]? Farahat and Mirsky echo this question, in a note added in proof to their 1958 article, and state that one of their theorems gives a "partial answer" to this question [3, p. 244]. We propose to return to another of Ranum's notions, that of "associated group member," and develop the theory for semigroups. We shall see that this will lead to a more complete solution of the question proposed, and one which applies equally well to semigroups in general.

It is known [3, p. 232] that if a is a group member in a semigroup S , then there is a maximal subgroup $M(a)$ of S , containing a , whose identity is the identity of every subgroup containing a . Further, the inverse of a , relative to this identity, is the same in every subgroup containing a . Distinct maximal subgroups are disjoint. We say that n is the *group index* of a in S if a^n is a group member in S and n is the smallest positive integer for which this is true. (Thus the group index of a group member is 1.) The following theorem is perhaps not so well known.

THEOREM 1. *If S is a semigroup and n is the group index of an element a in S , then a^t is a group member in S if and only if $t \geq n$. Furthermore, if $t \geq n$, then $M(a^t) = M(a^n)$.*

Proof. Let a^t be a group member in S . By the definition of group index, t cannot be less than n . Hence $t \geq n$.

Conversely, let $t \geq n$. If $t = n$, then $a^t = a^n$ and so a^t is a group

member in S since a^n is a group member in S . Now suppose $t > n$. Denote the identity of $M(a^n)$ by e , and for simplicity of notation, let b be the inverse of a relative to e . Then e is a two-sided identity for a^t , since

$$a^t e = (a^{t-n} a^n) e = a^{t-n} (a^n e) = a^{t-n} a^n = a^t,$$

and

$$e a^t = e (a^n a^{t-n}) = (e a^n) a^{t-n} = a^n a^{t-n} = a^t.$$

We next show that a^t is invertible relative to e . Let p be an integer such that $pn > 2t$. Then $a^{pn-t} b^p$ is a right inverse for a^t , since

$$a^t (a^{pn-t} b^p) = (a^t a^{pn-t}) b^p = a^{pn} b^p = (a^n b)^p = e^p = e.$$

We have also that $b^p a^{pn-t}$ is a left inverse for a^t , since

$$(b^p a^{pn-t}) a^t = b^p (a^{pn-t} a^t) = b^p a^{pn} = (b a^n)^p = e^p = e.$$

But e is a two-sided identity for $a^{pn-t} b^p$, because

$$(a^{pn-t} b^p) e = (a^{pn-t} b^{p-1}) (b e) = (a^{pn-t} b^{p-1}) b = a^{pn-t} b^p,$$

and

$$e (a^{pn-t} b^p) = (e a^t) (a^{pn-2t} b^p) = a^t (a^{pn-2t} b^p) = a^{pn-t} b^p.$$

Similarly, we may show that e is a two-sided identity for $b^p a^{pn-t}$. The uniqueness of the inverse of a^t follows from

$$\begin{aligned} a^{pn-t} b^p &= e (a^{pn-t} b^p) = e^p (a^{pn-t} b^p) = (b a^n)^p (a^{pn-t} b^p) = (b^p a^{pn}) (a^{pn-t} b^p) \\ &= (b^p a^{pn-t}) (a^{pn} b^p) = (b^p a^{pn-t}) (a^n b)^p = (b^p a^{pn-t}) e^p \\ &= (b^p a^{pn-t}) e = b^p a^{pn-t}. \end{aligned}$$

Thus a^t is invertible relative to e . Hence a^t is an element of $M(e)$, and so a^t is a group member in S . It follows that $M(a^t) = M(e) = M(a^n)$.

THEOREM 2. *If S is a semigroup, a has group index n in S , and e is the identity of $M(a^n)$, then eae is a group member in S and $(eae)^n = a^n$.*

Proof. Clearly e is a two-sided identity for eae . Let b be the inverse of a^n relative to e . We shall show that $a^{n-1}b$ is a two-sided inverse for eae relative to e , and that e is a two-sided identity for $a^{n-1}b$. We have that $a^{n-1}b$ is a right inverse for eae relative to e , for

$$\begin{aligned} (eae)(a^{n-1}b) &= (eae)[a^{n-1}(eb)] = (eae)[a^{n-1}(a^n b)b] \\ &= (eae)[a^n(a^{n-1}b)b] = (ea)(ea^n)(a^{n-1}b)b \end{aligned}$$

$$\begin{aligned} &= (ea)a^n(a^{n-1}b)b = (ea^n)a(a^{n-1}b)b \\ &= a^n(a^nb)b = a^n(eb) = a^nb = e . \end{aligned}$$

Similarly, we may show that ba^{n-1} is a left inverse for eae relative to e . That e is a two-sided identity for $a^{n-1}b$ follows from

$$(a^{n-1}b)e = a^{n-1}(be) = a^{n-1}b ,$$

and

$$\begin{aligned} e(a^{n-1}b) &= e[a^{n-1}(eb)] = e[a^{n-1}(a^nb)b] = e[a^n(a^{n-1}b)b] \\ &= (ea^n)(a^{n-1}b)b = a^n(a^{n-1}b)b = a^{n-1}(a^nb)b \\ &= a^{n-1}(eb) = a^{n-1}b . \end{aligned}$$

Similarly, we may show that e is a two-sided identity for ba^{n-1} . The uniqueness of the inverse of eae now follows from

$$ba^{n-1} = (ba^{n-1})e = (ba^{n-1})(a^nb) = (ba^n)(a^{n-1}b) = e(a^{n-1}b) = a^{n-1}b .$$

We have thus shown that eae is a group member in S . We now show that $(eae)^n = a^n$. We note first that $(eae)^n = ea^n e$, for

$$\begin{aligned} (eae)^2 &= (eae)(eae) = eaeae = eaea(a^nb) = ea(ea^n)(ab) \\ &= eaa^na b = ea^2(a^nb) = ea^2e , \end{aligned}$$

and by induction, $(eae)^n = ea^n e$. But e is a two-sided identity for a^n , so that $(eae)^n = ea^n e = a^n$.

The element eae of the above theorem is said to be the *associated group member* of a in S . (Note that a is a group member in S if and only if a is identical with its associated group member.)

THEOREM 3. *Let S be a semigroup, let T be a sub-semigroup of S , and let a belong to T . If a has group index n in S and group index t in T , then $n = t$.*

Proof. Clearly $n \leq t$. Now let e be the identity of the group in T to which a^t belongs. But a^t belongs to the maximal group $M(a^n)$ in S , so that e is the identity of any group in S to which a^n belongs. Then $ea^n e = a^n$. But $ea^n e$ is the associated group member in T of a^n . Hence, by Theorem 2, a^n is a group member in T . Thus $t = n$.

We are now in a position to answer the question proposed in the first paragraph.

THEOREM 4. *Let S be a semigroup and let T be a sub-semigroup of S . If an element a in T is a group member in S , then it is a group member in T if and only if some power of a is a group member in T .*

Proof. Suppose n is the group index of a in T . Since a is a group member in S , its group index in S is 1. By Theorem 3, $n = 1$. Hence, a is a group member in T .

The following theorem points out the wide application of this solution. (Proofs may be found in the places cited in the footnotes.)

THEOREM 5. *Every element of each of the following has finite group index:*

- (1) *finite semigroup* [4, pp. 12, 13]
- (2) *division ring or a direct sum of division rings* [4, pp. 19, 20]
- (3) *semi-divided ring* [3, p. 234]
- (4) *finite dimensional algebra* [6, p. 292].

REFERENCES

1. W. E. Barnes and H. Schneider, *The group membership of a polynomial in an element algebraic over a field*, Arkiv für Mathematik, **8** (1957), 166-168.
2. H. K. Farahat and L. Mirsky, *A condition for diagonality of matrices*, Amer. Math. Monthly, **63** (1956), 410-412.
3. ———, *Group membership in rings of various types*, Mathematische Zeitschrift, **70** (1958), 231-244.
4. H. C. Kennedy, *Group Membership in Rings*. Ann Arbor: University Microfilms, 1961.
5. Arthur Ranum, *The Group Membership of Singular Matrices*, Amer. J. Math., **31** (1909), 13-41.
6. ———, *The groups belonging to a linear associative algebra*, Amer. J. Math., **49** (1927), 285-308.

PROVIDENCE COLLEGE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RALPH S. PHILLIPS

Stanford University
Stanford, California

M. G. ARSOVE

University of Washington
Seattle 5, Washington

A. L. WHITEMAN

University of Southern California
Los Angeles 7, California

LOWELL J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

T. M. CHERRY

D. DERRY

M. OHTSUKA

H. L. ROYDEN

E. SPANIER

E. G. STRAUS

F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insa'susha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 12, No. 2

February, 1962

William George Bade and Robert S. Freeman, <i>Closed extensions of the Laplace operator determined by a general class of boundary conditions</i>	395
William Browder and Edwin Spanier, <i>H-spaces and duality</i>	411
Stewart S. Cairns, <i>On permutations induced by linear value functions</i>	415
Frank Sydney Cater, <i>On Hilbert space operators and operator roots of polynomials</i>	429
Stephen Urban Chase, <i>Torsion-free modules over $K[x, y]$</i>	437
Heron S. Collins, <i>Remarks on affine semigroups</i>	449
Peter Crawley, <i>Direct decompositions with finite dimensional factors</i>	457
Richard Brian Darst, <i>A continuity property for vector valued measurable functions</i>	469
R. P. Dilworth, <i>Abstract commutative ideal theory</i>	481
P. H. Doyle, III and John Gilbert Hocking, <i>Continuously invertible spaces</i>	499
Shaul Foguel, <i>Markov processes with stationary measure</i>	505
Andrew Mattei Gleason, <i>The abstract theorem of Cauchy-Weil</i>	511
Allan Brasted Gray, Jr., <i>Normal subgroups of monomial groups</i>	527
Melvin Henriksen and John Rolfe Isbell, <i>Lattice-ordered rings and function rings</i>	533
Amnon Jakimovski, <i>Tauberian constants for the $[J, f(x)]$ transformations</i>	567
Hubert Collings Kennedy, <i>Group membership in semigroups</i>	577
Eleanor Killam, <i>The spectrum and the radical in locally m-convex algebras</i>	581
Arthur H. Kruse, <i>Completion of mathematical systems</i>	589
Magnus Lindberg, <i>On two Tauberian remainder theorems</i>	607
Lionello A. Lombardi, <i>A general solution of Tonelli's problem of the calculus of variations</i>	617
Marvin David Marcus and Morris Newman, <i>The sum of the elements of the powers of a matrix</i>	627
Michael Bahir Maschler, <i>Derivatives of the harmonic measures in multiply-connected domains</i>	637
Deane Montgomery and Hans Samelson, <i>On the action of $SO(3)$ on S^n</i>	649
J. Barros-Neto, <i>Analytic composition kernels on Lie groups</i>	661
Mario Petrich, <i>Semicharacters of the Cartesian product of two semigroups</i>	679
John Sydney Pym, <i>Idempotent measures on semigroups</i>	685
K. Rogers and Ernst Gabor Straus, <i>A special class of matrices</i>	699
U. Shukla, <i>On the projective cover of a module and related results</i>	709
Don Harrell Tucker, <i>An existence theorem for a Goursat problem</i>	719
George Gustave Weill, <i>Reproducing kernels and orthogonal kernels for analytic differentials on Riemann surfaces</i>	729
George Gustave Weill, <i>Capacity differentials on open Riemann surfaces</i>	769
G. K. White, <i>Iterations of generalized Euler functions</i>	777
Adil Mohamed Yaqub, <i>On certain finite rings and ring-logics</i>	785