Pacific Journal of Mathematics

LIMITS AND BOUNDS FOR DIVIDED DIFFERENCES ON A JORDAN CURVE IN THE COMPLEX DOMAIN

J. H. CURTISS

Vol. 12, No. 4

April 1962

LIMITS AND BOUNDS FOR DIVIDED DIFFERENCES ON A JORDAN CURVE IN THE COMPLEX DOMAIN

J. H. CURTISS

1. Introduction. Let $S_{n+1} = \{z_1, z_2, \dots, z_{n+1}\}$ be a set of n + 1 complex numbers and let f be a function on a set containing S_{n+1} to the complex numbers. The divided difference $d_n = d_n(f | z_1, z_2, \dots, z_{n+1})$ of order n formed for the function f in the points S_{n+1} is defined in a recursive manner as follows:

$$egin{aligned} &d_1 = d_1(f \,|\, z_1, z_2) = rac{f(z_1) - f(z_2)}{z_1 - z_2} \ &d_2 = d_2(f \,|\, z_1, z_2, z_3) = rac{d_1(f \,|\, z_1, z_2) - d_1(f \,|\, z_3, z_2)}{z_1 - z_3} \ &dots \ &d_n = d_n(f \,|\, z_1, z_2, \, \cdots, \, z_{n+1}) \ &= rac{d_{n-1}(f \,|\, z_1, z_2, \, \cdots, \, z_n) - d_{n-1}(f \,|\, z_{n+1}, \, z_2, \, \cdots, \, z_n)}{z_1 - z_{n+1}} \ \end{aligned}$$

The definition requires further discussion when the points in S_{n+1} are not all distinct. We shall suppose that they are distinct unless provision is explicitly made for coincidences.

It can be proved by induction [7, p. 15] that if

$$\omega_{n+1}(z) = (z-z_1)(z-z_2)\cdots(z-z_{n+1})$$
 ,

then

(1.1)
$$d_n = \sum_{k=1}^{n+1} \frac{f(z_k)}{\omega'_{n+1}(z_k)}$$

where the prime denotes differentiation of $\omega_{n+1}(z)$ with respect to z. This formula shows that d_n is a symmetric function of z_1, z_2, \dots, z_{n+1} .

The divided differences of a function given on the real line play a prominent role in the mathematics of computation. Their counterparts in the complex plane have appeared in various classical studies of approximation by complex polynomials. The formal algebra of complex

Received January 10, 1962. This research was supported by the United States Air Force through the Air Force Office of Scientific Research of the Air Research and Development Command, under Contract No. AF 49 (638)-862.

¹ We use the words "points" and "numbers" interchangeably in referring to the arguments in divided differences. This follows the practice in interpolation theory. It is consistent within this terminology to speak of "coincident points" z_k .

divided differences is of course much the same as for the real case, but the analytical properties of complex divided differences, such as asymptotic behavior and representability by integrals, are in some cases quite different. It would appear that these analytical properties have not received much attention in the literature, although some of them seem interesting.

A primary motivation for the present paper was the need to establish that under certain smoothness hypotheses on a function f given on a Jordan curve C, the divided difference of f of a fixed order formed in points on C is uniformly bounded in modulus for all choices of the points. This property was required in a study of complex interpolation in random points [1]. The existence of the bound is proved in §2 below for the case in which C is the unit circle. The extension to more general Jordan curves appears in §3. In §4, the asymptotic behavior of successive divided differences of order n formed in n+1 points on a Jordan curve, $n = 1, 2, \cdots$, which in their totality become everywhere dense in a certain way on the curve, is investigated. It is found that the behavior to be expected in cases important in the theory of complex interpolation is that the *n*th divided difference multiplied by the (n+1)th power of the transfinite diameter, or capacity, of the curve C in question approaches the limit $\int_{a} f dz/(2\pi i)$. Section 4 is essentially self-contained and can be read separately.

As was mentioned above, the impetus for this study came from a particular application. It is hoped that the results may turn out to be useful in other directions. However, the general spirit in which this paper is written is that of interest in the subject for itself alone, and the possible applications will not be considered further.

2. An upper bound for the modulus of a divided difference formed on the unit circle. If the numbers z_k are all real numbers, and if f is continuous on a closed interval \overline{I} of the real line containing S_{n+1} and possesses an *n*th derivative $f^{(n)}$ at each point of the corresponding open interval *I*, then by elementary calculus [7, p. 24] it can be shown that there exists a number x_0 in *I* such that $d_n = f^{(n)}(x_0)/n!$. Thus if $|f^{(n)}|$ is uniformly bounded everywhere on *I*, so also is $|d_n|$ for all choices of S_{n+1} on \overline{I} . Again, with real points S_{n+1} , if $f^{(n-1)}$ is absolutely continuous [8, pp. 364 ff.] on \overline{I} , then the iterated integral on the right side of the following formula (in which we define z_{n+2} as meaning z_1),

$$(2.1) \qquad d_n(f \mid z_1, z_2, \cdots, z_{n+1}) \\ = \int_0^1 \int_0^{y_1} \cdots \int_0^{y_{n-1}} f^{(n)} \Big[z_2 + \sum_{j=1}^n y_j (z_{j+2} - z_{j+1}) \Big] dy_1 \, dy_2 \cdots \, dy_n ,$$

has meaning for all choices of S_{n+1} in which the points z_j are distinct,

and indeed can be used to extend the definition of d_n to cases involving confluent points. It is easily shown by induction that the formula is true [7, pp. 17–18]. Thus it follows with $f^{(n-1)}$ absolutely continuous and $|f^{(n)}|$, where it exists, uniformly bounded on *I*, that $|d_n|$ is also uniformly bounded for all choices of S_{n+1} on \overline{I} such that completion of the definition of d_n through (2.1) is possible. If *M* is the least upper bound of $|f^{(n)}|$ on *I*, then $|d_n| \leq M/n!$.

The formula (2.1) is no longer generally valid when the numbers S_{n+1} are not all real, and the derivation of a bound for $|d_n|$ in terms of a given bound for $|f^{(n)}|$ is not so readily accomplished. In the remainder of the section we shall consider this problem in the case in which S_{n+1} lies on the unit circle in the complex plane.

In the development, we shall use a complex-variable type of interpretation of the derivatives of a function g given on the circle C: |z|=1. The symbol $g^{(1)}(z_1)$ will mean

$$\lim_{z o z_1} d_{\scriptscriptstyle 1}(g \,|\, z,\, z_1) = \lim_{z o z_1} rac{g(z) - g(z_1)}{z - z_1} \;, \; \; |\, z \,| = 1, |\, z_1 \,| = 1 \;,$$

provided of course that the limit exists. Higher derivatives $g^{(k)}$ are to be defined recursively. The circle C can be parametrized in a one-toone manner by the equation $z = e^{i\theta}$, with $\alpha \leq \theta < \alpha + 2\pi$, where α is chosen arbitrarily. If this is done, then

$$g^{\scriptscriptstyle (1)}(z) = rac{dg}{d heta} \cdot rac{d heta}{de^{i heta}} = rac{dg}{d heta} \cdot rac{1}{ie^{i heta}} \, .$$

The chief result is this:

THEOREM 2.1 Let the function f be given on C: |z| = 1 together with its first n-1 derivatives $f^{(1)}, f^{(2)}, \dots, f^{(n-1)}$. Let the points S_{n+1} lie on C and be distinct. Then if $f^{(n-1)}$ satisfies the Lipschitz condition:

$$| f^{(n-1)}(z) - f^{(n-1)}(t) | \leq \lambda | z - t |, \lambda > 0$$
 ,

for all z and t on C, it follows that

$$(2.2) | d_n(f | z_1, z_2, \cdots, z_{n+1}) | \leq \frac{\frac{\pi}{2} \left(\frac{\sqrt{\pi}}{2}\right)^{n-1} \lambda_n}{\Gamma\left(\frac{n+1}{2}\right)}$$

uniformly for all such S_{n+1} , where λ_n is the least upper bound of

$$|f^{(n)}(z)|, |z| = 1$$
.

The symbol Γ in (2.2) refers to the Gamma Function.

The hypothesis on $f^{(n-1)}$ has the implication that $f^{(n-1)}$ be absolutely continuous on |z| = 1, and that it therefore be the indefinite integral of a derivative $f^{(n)}$ existing everywhere on |z| = 1 with the possible exception of a set of Lebesgue measure zero. Moreover the implication is that $|f^{(n)}|$, on the set where it exists, is bounded and its least upper bound λ_n does not exceed λ .

Our proof involves integral representations, and it is important to be explicit about the integral calculus to be used.

Consider two points $e^{i\alpha_2}$ and $e^{i\alpha_1}$ on the unit circle. The complex line integral of a function g given on the unit circle extended over either one of the two arcs of the circle joining these points, directed from $e^{i\alpha_2}$ to $e^{i\alpha_1}$, is to be defined as a Lebesgue integral with respect to the parameter θ in the parametrization $z = e^{i\theta}$. That is, if A is the chosen directed arc, then

(2.3)
$$\int_{A} g(z) dz = \int_{\alpha_2}^{\alpha_1} g(e^{i\theta}) i e^{i\theta} d\theta .$$

If g is continuous in a neighborhood of $e^{i\alpha_1}$, then

(2.4)
$$\frac{d}{de^{i\alpha_1}}\int_A g(z)dz = \frac{g(e^{i\alpha_1})ie^{i\alpha_1}}{ie^{i\alpha_1}} = g(e^{i\alpha_1}) \ .$$

The notation for the integral on the right side of (2.3) is ambiguous in that it does not indicate which one of the two possible directed arcs A is being integrated over. However in the sequel we shall be dealing only with complex line integrals on |z| = 1 which are independent of the path of integration. Such an integral extended over either arc directed from z_2 to z_1 , $|z_1| = |z_2| = 1$, will be denoted by

$$\int_{z_2}^{z_1} g(z) dz \; .$$

If the two arcs joining z_2 to z_1 are are of equal length, then $\alpha_2 = \alpha_1 \pm \pi$, so the variation of θ in (2.3) is over a closed interval of length π of which one endpoint is α_2 . If the two paths are not of equal length, then the shorter one corresponds under $z = e^{i\theta}$ to an interval of values of θ of which one endpoint is α_2 and the other one, say α' , is such that $z_1 = e^{i\alpha_1} = e^{i\alpha'}$ and $|\alpha_2 - \alpha'| < \pi$. (For example, if α_1 and α_2 are restricted to the interval $[0, 2\pi]$ and if $\alpha_2 > \alpha_1, \alpha_2 - \alpha_1 > \pi$, then we take $\alpha' = 2\pi + \alpha_1$.)

We shall now drop the parentheses around superscripts indicating derivatives of functions, but it is to be understood that superscripts can also be exponents when the context requires, as in $(z - z_2)^k$. In the case of divided differences, a derivative superscript will always indicate a partial derivative with respect to the first argument when the

notation in the first paragraph of the Introduction is being used. That is

$$d^k_{{}_m} = d^k_{{}_m}(f \,|\, z_1, z_2, \, \cdots, \, z_{m+1}) \; rac{\partial^{(k)} d_{{}_m}(f \,|\, z_1, \, z_2, \, \cdots, \, z_{m+1})}{\partial z_1^{(k)}} \;.$$

To prove Theorem 2.1 we need two lemmas, of which the first is as follows.

LEMMA 2.1 Let the function f given on C: |z| = 1 be such that its (n-1)st derivative exists everywhere on C and is absolutely continuous. Then

$$(2.5) d_1^{h-1}(f \mid z_1, z_2) = \frac{\int_{z_2}^{z_1} (t-z_2)^{h-1} f^h(t) dt}{(z_1-z_2)^h} , \\ |z_1| = 1, \quad |z_2| = 1, \quad z_1 \neq z_2, h = 1, 2, \cdots, n .$$

The integral is independent of the path of integration on C.

The absolute continuity of f^{n-1} implies the absolute continuity of f, f^2, \dots, f^{n-2} , and so implies that each of these functions including f^{n-1} is the indefinite integral of its derivative.

In the case h = 1, with $z_1 = e^{i\alpha_1}, z_2 = e^{i\alpha_2}$,

$$rac{\int_{z_2}^{z_1} f'(t)\,dt}{z_1-z_2} = rac{\int_{x_2}^{x_1} f'(e^{i heta})ie^{i heta}d heta}{z_1-z_2} = rac{f(z_1)-f(z_2)}{z_1-z_2} = rac{\int_{x_2}^{x_1\pm 2\pi} f'(e^{i heta})ie^{i heta}d heta}{z_1-z_2} \ .$$

The second and fourth members of the equation show that whether $\alpha_1 > \alpha_2$ or $\alpha_1 < \alpha_2$, the integral is independent of the path. Thus the Lemma is true for h = 1.

Suppose now that (2.5) gives a valid representation of d_1^{k-1} , $1 \leq k < h$, with the integral independent of the path and $z_1 \neq z_2$. Then using (2.4), we have after a brief computation

$$(2.6) d_1^k = \frac{\partial d_1^{k-1}}{\partial z_1} \frac{(z_1 - z_2)^k f^k(z_1) - k \int_{z_2}^{z_1} (t - z_2)^{k-1} f^k(t) dt}{(z_1 - z_2)^{k+1}} \,.$$

Because of the absolute continuity of f^k , integration by parts is valid

in the integral in (2.6), with f^k to be differentiated and $(t - z_2)^{k-1}$ to be integrated with respect to t. We thereby immediately obtain (2.5) with h = k + 1, and the integral is again independent of the path. The premises of the induction are true for k = 1, so this establishes the Lemma.

LEMMA 2.2 Let the function g be given at all points on |z| = 1with the possible exception of a set of Lebesgue measure zero; let $|g(z)| \leq M$ where defined on |z| = 1, and let g be such that

$$I_{k}(z_{\scriptscriptstyle 1},\, z_{\scriptscriptstyle 2}) = \int_{z_{\scriptscriptstyle 2}}^{z_{\scriptscriptstyle 1}} (t - z_{\scriptscriptstyle 2})^{k} g(t)\, dt, \;\;\; |\, z_{\scriptscriptstyle 1}\, | = |\, z_{\scriptscriptstyle 2}\, | = 1, \;\;\; k \ge 0 \;,$$

is independent of the path of integration. Then

(2.7)
$$\left| \frac{I_k(z_1, z_2)}{(z_1 - z_2)^{k+1}} \right| \leq \frac{\sqrt{\pi}}{2} \cdot \frac{\Gamma\left(\frac{k+1}{2}\right)}{\Gamma\left(\frac{k+2}{2}\right)} M, \ k = 0, 1, 2, \cdots,$$

for all z_1 and z_2 , $z_1 \neq z_2$, on |z| = 1.

When k = 0, the right side of (2.7) reduces to $\pi M/2$.

For the proof, we make the shorter arc joining $z_2 = e^{i\alpha_2}$ and $z_1 = e^{i\alpha_1}$ (or either of the two arcs if they are equal in length) correspond under $z = e^{i\theta}$ to a θ -interval $[\alpha_2, \alpha']$ or $[\alpha', \alpha_2]$, where α' is such that $z_1 = e^{i\alpha'}$ and $|\alpha_2 - \alpha'| \leq \pi$. Thus

$$I_k(z_{\scriptscriptstyle 1},\,z_{\scriptscriptstyle 2})=\int_{x_{\scriptscriptstyle 2}}^{x'}\!(e^{i heta}-\,e^{ilpha_{\scriptscriptstyle 2}})^kg(e^{i heta})ie^{i heta}d heta$$
 .

For the case k = 0, we use the inequality $|\sin \theta| \ge |2\theta/\pi|, -\pi/2 \le \theta \le \pi/2$, which is merely an expression of the fact that $\sin \theta$ is convex on $[0, \pi/2]$. We also use the identity $e^{i\alpha} - e^{i\beta} = 2ie^{i(\alpha+\beta)/2} \sin [(\alpha - \beta)/2]$. Then since $|\alpha' - \alpha_2|/2 \le \pi/2$, it follows that $|z_1 - z_2| = 2 \sin [(\alpha' - \alpha_2)/2]| \ge (2/\pi) |2(\alpha' - \alpha_2)/2|$. Now $|I_0(z_1, z_2)| \le M |\alpha' - \alpha_2|$, so the inequality (2.7) follows at once for k = 0.

For $k \ge 1$ we have (recalling the restriction on $|\alpha' - \alpha_2|$),

(2.8)
$$\frac{|I_k(z_1, z_2)|}{|z_1 - z_2|^{k+1}} \leq \frac{M2^k \left| \int_{\alpha_2}^{\alpha'} \left| \sin^k \left(\frac{\theta - \alpha_2}{2} \right) \right| d\theta \right|}{2^{k+1} \left| \sin^{k+1} \frac{\alpha' - \alpha_2}{2} \right|}$$
$$= \frac{M}{2} \frac{\left| \int_{\alpha_2}^{\alpha_1} \sin^k \left| \frac{\theta - \alpha_2}{2} \right| d\theta \right|}{\sin^{k+1} \left| \frac{\alpha' - \alpha_2}{2} \right|}.$$

We make the substitution $\beta = |\theta - \alpha_2|/2$ in the integral and let $\gamma = |\alpha' - \alpha_2|/2 \leq \pi/2$. By examination of the various cases corresponding to k even or odd and $\alpha' < \alpha_2, \alpha' > \alpha_2$ we find that the righthand member of (2.8) is always equal to

Inspection of its derivative shows that $S(\gamma)$ increases steadily with γ on the interval $0 \leq \gamma \leq \pi/2$. The value of $S(\pi/2)$ is given by the well known formula

$$\int_{_{0}}^{^{\pi/2}} \sin^{k} heta d heta = rac{\sqrt{\pi}}{2} rac{\Gamma\left(rac{k+1}{2}
ight)M}{\Gamma\left(rac{k+2}{2}
ight)}$$

Thus

$$rac{\mid I_k(z_1,\,z_2)\mid}{\mid z_1-z_2\mid^{k+1}} \leq MS\!\left(rac{\pi}{2}
ight) = rac{\sqrt{2}}{2}rac{arGamma\left(rac{k+1}{2}
ight)}{arGamma\left(rac{k+2}{2}
ight)}\,M$$
 ,

as was to be proved.

Now let g in Lemma 2.2 be f^n and M be λ_n , where f is the function appearing in the Theorem. The two lemmas establish that

(2.9)
$$d_1^{n-1}(f | z_1, z_2) = \frac{\lambda_n \sqrt{\pi} \Gamma\left(\frac{n}{2}\right)}{2\Gamma\left(\frac{n+1}{2}\right)}, \quad z_1 \neq z_2.$$

As a function of z_1 , d_1^{n-1} is continuous for $z_1 \neq z_2$ and uniformly bounded in modulus. Consider next $d_1^{n-2}(f z_1, z_2)$ as a function of z_1 . This function has a continuous derivative for $z_1 \neq z_2$, which is moreover uniformly bounded in modulus. Therefore this function is absolutely continuous in z_1 for z_1 on any closed arc of the unit circle not containing z_2 . But the uniform boundedness of $|d_1^{n-1}|$ implies that $d_1^{n-2}(f | z_1, z_2)$ is of uniformly bounded variation in z_1 on the entire unit circle with the point z_2 deleted. By a well-known theorem [8, p. 372, Ex. 6] it follows that as z_1 approaches z_2 from either side, d_1^{n-2} approaches a limit; and if the limit is the same for approach from either side, then when the definition of d_1^{n-2} is completed by this limit at $z_1 = z_2$ the function d_1^{n-2} will be an absolutely continuous function for all z_1 on $|z_1| = 1$. To investigate the limit, we write (2.5) in the form

$$d_1^{n-2} = \left[rac{\int_{lpha_2}^{lpha_1} (e^{i heta} - e^{ilpha_2})^{n-2} f^{n-1}(e^{i heta}) i e^{i heta} d heta}{\left(\sin rac{lpha_1 - lpha_2}{2}
ight)^{n-1}}
ight] \cdot rac{1}{(2i e^{i(lpha_1 + lpha_2)/2})^{n-1}} \; .$$

The limit as $z_1 \rightarrow z_2$, $\alpha_1 \rightarrow \alpha_2$, of the expression in square brackets can be evaluated by l'Hospital's rule used with (2.4) and with the fact that f^{n-1} is continuous. We find that for z_1 approaching z_2 on either side, there is the unique limit

$$\lim_{z_1 \to z_2} \, d_1^{n-2}(f \,|\, z_1, \, z_2) = \frac{f^{n-1}(z_2)}{n-1} \; .$$

Thus with proper completion of the definition of d_1^{n-2} at $z_1 = z_2$, this function is an absolutely continuous function of z_1 on $|z_1| = 1$. Similarly we can complete the definition of $d_1^h(f | z_1, z_2)$ for $h = n - 3, n - 4, \cdots$, 1, 0, so that the resulting function is in each case an absolutely continuous function of z_1 on $|z_1| = 1$. We assume henceforth without change in notation that for each relevant value of h, the proper extension of the definition of d_1^h at $z_1 = z_2$ has been made.

What this establishes is that the completed first order divided difference $d_1(f | z_1, z_2)$, as a function of z_1 , together with its first n-2partial derivatives with respect to z_1 , have the same smoothness and integrability properties as does f and its first n-1 derivatives. That is to say, $d_1, d_{11}^2, \dots, d_1^{n-2}$ are absolutely continuous functions of z_1 and moreover the derivative of d_1^{n-2} , where it exists, is uniformly bounded in modulus.

The absolute continuity of the derivatives permits the inductive argument which we used to establish (2.5) to be used again to prove that

$$egin{aligned} &d_2^{h-1}(f\,|\,z_1,z_2,z_3)=rac{\partial^{h-1}}{\partial z_1^{h-1}}d_1(d_1(\,f\,|\,z,z_2)\,|\,z_1,z_3)\ &=rac{\int_{z_3}^{z_1}\!\!\!(t-z_3)^{h-1}d_1^h(f\,|\,t,z_2)\,dt}{(z_1-z_3)^h}\ &|\,z_1\,|\,=1,\;|\,z_2\,|\,=1,\;z_1
eq z_3,\;h=1,\,2,\,\cdots,\,n-1 \;. \end{aligned}$$

By (2.9) and Lemma 2.2, d_2^{n-2} as a function of z_1 is uniformly bounded in modulus. (It is not important at the moment to know how the bound depends on z_2 and z_3 .) The definitions of d_2^{n-3} , d_2^{n-4} , \cdots , d_2 can now be completed by continuity at $z_1 = z_3$ so that in each case the resulting function of z_1 is absolutely continuous on $|z_1| = 1$. Again we assume without change of notation that the proper extensions have been made.

Proceeding in this way, we establish the chain of equations

in which $d_n^0 = d_n$, $d_0^n = f^n$. Theorem 2.1 now can be proved by back substitution into (2.10), beginning with (2.9) and using Lemma 2.2 at each stage. Thus to start with, at least for $z_1 \neq z_2$ and $z_1 \neq z_3$,

$$|\,d_{\scriptscriptstyle 2}^{n-2}\,| \leq rac{\sqrt{\pi}}{2} rac{\Gamma\left(rac{n-1}{2}
ight)}{\Gamma\left(rac{n}{2}
ight)} \!\left[\!rac{\lambda_n\sqrt{\pi}\,\Gamma\left(rac{n}{2}
ight)}{2\Gamma\!\left(rac{n+1}{2}
ight)}
ight].$$

Similarly,

$$egin{aligned} &|d_3^{n-3}| \leq \left(rac{\sqrt{\pi}}{2} \Gammaigg(rac{n-2}{2}igg) {2\Gammaigg(rac{n-2}{2}igg)} {2\Gammaigg(rac{n}{2}igg)}
ight) & \left[rac{\sqrt{\pi}}{2} \Gammaigg(rac{n-1}{2}igg) {2\Gammaigg(rac{n}{2}igg)}
ight)
ight] \ &= \left(rac{\sqrt{\pi}}{2}
ight)^3 rac{\Gammaigg(rac{n-2}{2}igg)} {\Gammaigg(rac{n-2}{2}igg)} \lambda_n \ , \end{aligned}$$

and so forth. We finally find that

$$|\,d_n^{\scriptscriptstyle 0}\,| \leqq rac{\pi}{2} \Big(rac{\sqrt{\pi}}{2}\Big)^{n-1} rac{\lambda_n}{arGamma\left(rac{n+1}{2}
ight)}\,,$$

as stated in the conclusion of the Theorem.

It is clear from the proof that under the hypotheses of Theorem 2.1 on f, it is possible to extend the definition of d_n by continuity so as to admit point sets S_{n+1} in which coincidences occur, and then (2.2) will still be valid. However we shall not study this question in detail under the hypotheses of Theorem 2.1 on f, which were chosen as being natural to achieve boundedness of $|d_n|$ in the case of distinct points. (The boundedness of $d_1 = |f(z_1) - f(z_2)| / |z_1 - z_2|$ is equivalent to a Lipschitz condition on f.)

The method of proof with only slight modifications can be used to establish the following result:

THEOREM 2.2 Under the hypothesis of Theorem 2.1 concerning f,

and with the added hypothesis that $f^{(n)}$ is continuous on some open arc of |z| = 1 containing the point z_1 , the following equation completes the definition of d_n by continuity for the case in which all the points S_{n+1} coincide at z_1 :

(2.11)
$$d_n(f | z_1, z_1, \cdots, z_1) = \frac{f^{(n)}(z_1)}{n!}.$$

If $f^{(n)}$ is everywhere continuous on |z| = 1, then (2.2) is valid after proper completion of the definition of d_n for all choices of S_{n+1} without restrictions as to coincidences.

We conclude this section with two comments. In the first place it is clear that by repeated back-substitution into (2.10), a single formula for d_n in terms of $f^{(n)}$ involving repeated integration can be written out. It would be somewhat similar in appearance to a variant of (2.1) which appears in [5, p. 18, ex. 7].

In the second place, it may be that for $n \ge 2$ the bound in (2.2) can be improved. For n = 1 it is the best bound possible, as can be seen from this trivial example: Let f be real and let its graph over a period in the $(\theta, f(e^{i\theta}))$ -plane be a line segment joining (0, 0) to (π, π) and another line segment joining (π, π) to $(2\pi, 0)$. For this function the maximum of d_1 is $\pi/2$, the least upper bound of |f'| is one, and the right side of (2.2) is $(\pi/2) \cdot 1$, which is as small as it can be. However the general bound was derived through Lemma 2.2 in which the two points z_1 and z_2 were placed at opposite ends of a diameter to obtain the numerical appraisal. Such wide-apart spacing is of course not possible for the case of three or more points on the unit circle. The bound given by (2.1) in the real case under the hypothesis of Theorem 2.1 is $\lambda_n/n!$, which is much smaller than that in (2.2).

3. Boundedness of the modulus of a divided difference formed on a general Jordan curve. A generalization of Theorem 2.1 to the case in which the unit circle is replaced by a more general Jordan curve is not hard to derive. In doing so, for simplicity we shall not try to keep track of the structure of the upper bound, and shall suppress various details in the proof.

A Jordan curve is homeomorphic to a circle. It can be represented by a parametric equation $z = \phi(\theta)$, where ϕ is continuous in the real variable θ with period 2π , and where for each given point z on the curve, any two solutions of $z = \phi(\theta)$ differ by an integral multiple of 2π . Our considerations here will be restricted to Jordan curves such that the first derivative $d\phi/b\theta = \psi(\theta)$ exists for all θ and is continuous, and $\psi(\theta) \neq 0$ for all θ . Such a Jordan curve will be said to be "admissible". (Presumably in what follows the definition of admissibility can be slightly relaxed.)

LEMMA 3.1. If $z = \phi(\theta)$ is a parametric equation of an admissible Jordan curve, then there exist numbers m and M, 0 < m < M, such that

$$m \leq \left|rac{\phi(heta_1) - \phi(heta_2)}{e^{i heta_1} - e^{i heta_2}}
ight| \leq M$$

for all θ_1 and θ_2 .

The divided difference appearing in the above inequality is to be interpreted as meaning $\psi(\theta_2)/ie^{i\theta_2}$ when $\theta_1 = \theta_2$.

The existence of an upper bound M follows from Lemma 2.2 with k = 0 and $g(e^{i\theta}) = \psi(\theta)/ie^{i\theta}$. The existence of the lower bound can be established by an elementary indirect argument which we omit².

As in the unit circle case, it is convenient to interpret the derivatives of a function on a Jordan curve to the complex numbers as limits of complex-variable difference quotients. Specifically for any function g given on a Jordan curve C, the symbol $g'(z_1)$ means $\lim_{z\to z_1} d_1(g \mid z, z_1)$, z and z_1 on C, and higher derivatives are to be defined recursively. If C is admissible, then

$$g'(z) = rac{d(g(\phi(heta)))}{d heta} \cdot rac{1}{\psi(heta)} \cdot$$

Integrals are to be defined as in (2.3) with $ie^{i\theta}$ in that formula replaced by $\psi(\theta)$. With this replacement and with $e^{i\alpha_1}$ replaced by $\phi(\alpha_1)$, (2.4) is is valid. An integral over C with limits of integration z_1 and z_2 which is independent of the path will be written as

$$\int_{z_2}^{z_1} g(z) dz .$$

The notation implies of course that the arc over which the integration takes place is directed from z_2 to z_1 . The derivatives of divided differences are always partials with respect to the first apparent argument.

The generalization of Theorem 2.1 is as follows:

THEOREM 3.1. Let the function f be given on an admissible Jordan curve C, together with f^1, f^2, \dots, f^{n-1} . Let f^{n-1} satisfy the Lipschitz condition

$$|f^{n-1}(z) - f^{n-1}(t)| \leq \lambda |z-t|, \lambda > 0$$
 ,

for all z and t on C. Let the points $S_{n+1} = \{z_1, z_2, \dots, z_{n+1}\}$ lie on C and be distinct. Then there exists a constant M depending only on n, λ , and C, and independent of S_{n+1} , such that $|d_n(f | z_1, z_2, \dots, z_{n+1})| \leq M$.

² Various related but deeper results may be found in [6, Section 2.5].

The proof starts with a generalization of Lemma 2.1.

LEMMA 3.2. Let the function f, given on an admissible Jordan curve C, be such that its (n-1)th derivative exists everywhere on C and is absolutely continuous as a function of θ , $z = \phi(\theta)$. Then

$$d_1^{h-1}(f \,|\, z_1, z_2) = rac{\int_{z_2}^{z_1} (t-z_2)^{h-1} f^h(t) \, dt}{(z_1-z_2)^h}$$

,

for z_1 and z_2 on C, $z_1 \neq z_2$, $h = 1, 2, \dots, n$, and the integral is independent of the path of integration on C.

The argument used to establish Lemma 2.1 carries over to Lemma 3.2 with only minor changes, and will not be restated here.

LEMMA 3.3. Let the function G be given on the admissible Jordan curve C with the possible exception of a set of Lebesgue measure zero; let |G| be bounded on C and be such that

$$J_k(z_1, z_2) = \int_{z_2}^{z_1} (t - z_2)^k G(t) dt$$

is independent of the path of integration on C for all z_1 and z_2 on C. Then for each $k, k = 0, 1, \dots$, there exists a constant M_k , depending only on G and C, and such that

$$\left|rac{{J}_k(z_1,\,z_2)}{(z_1-z_2)^{k+1}}
ight| \leq M_k$$

for all z_1 and z_2 on C, $z_1 \neq z_2$.

The Lebesgue measure in the theorem means measure on the θ -line after the transformation $\phi^{-1}: z \to \theta$.

To prove this lemma, we let $z_1 = \phi(\alpha_1), z_2 = \phi(\alpha_2), t = \phi(\theta)$, and write

(3.1)
$$\frac{J_k(z_1, z_2)}{(z_1 - z_2)^{k+1}} = \left[\frac{e^{i\alpha_1} - e^{i\alpha_2}}{\phi(\alpha_1) - \phi(\alpha_2)}\right]^{k+1} \frac{\int_{\alpha_2}^{\alpha_1} (e^{i\theta} - e^{i\alpha_2})^k g(e^{i\theta}) i e^{i\theta} d\theta}{(e^{i\alpha_1} - e^{i\alpha_2})^{k+1}},$$

where

$$(3.2) g(e^{i\theta})ie^{i\theta} = G(\phi(\theta)) \left[\frac{\phi(\theta) - \phi(\alpha_2)}{e^{i\theta} - e^{i\alpha_2}} \right]^k \psi(\theta) \; .$$

By Lemma 3.1, the quantities in the square brackets in (3.1) and (3.2)

are both uniformly bounded in modulus for all θ , α_1 , and α_2 , $\theta \neq \alpha_2$, $\alpha_1 \neq \alpha_2$. For any fixed α_2 , $g(e^{i\theta})$ as given by (3.2) is integrable, and its modulus is uniformly bounded for all θ and α_2 . The integral in (3.1), considered as an integral over an arc of the unit circle, is independent of the path of integration. Thus the hypotheses of Lemma 2.2 are satisfied by gas given by (3.2). The truth of Lemma 3.3 now follows immediately.

Theorem 3.1 can now be proved by the use of Lemmas 3.2 and 3.3 in the same way that Theorem 2.1 was proved. The hypotheses on fand C imply that $f^{n-1}(e^{i\theta})$ is an absolutely continuous function of $e^{i\theta}$ and of θ , and that its derivative with respect to $\phi(\theta)$, where it exists, is uniformly bounded in modulus. The same is true for its derivative with respect to θ . (These facts follow from the existence of numbers λ_1 and λ_2 such that with $z = e^{i\theta}$, $t = e^{i\omega}$,

$$|f^{n-1}(z)-f^{n-1}(t)|\leq \lambda \,|\,\phi(heta)-\phi(lpha)\,|\leq \lambda_1\,|\,e^{i heta}-e^{ilpha}\,|\leq \lambda_2\,|\, heta-lpha\,|$$
 .

(Here we used Lemma 3.1 in passing from the second member to the third member of the chain.) The functions f, f^1, \dots, f^{n-2} are also absolutely continuous and have uniformly bounded derivatives.

We can now re-establish the recursion formulas (2.10), which look exactly the same as before and so will not be repeated here. The integrals in (2.10) are of course now complex line integrals over C. Thereafter by back-substitution, using Lemma 3.3 at each stage, we establish the existence of the bound for $|d_n|$.

The analogous generalization of Theorem 2.2 is also valid. The proper definition of d_n for confluent points is again given by (2.11). It is worth noting that what gives simplicity to our results and minimizes the restrictions on C is the complex-variable type of definition which we are using for derivatives of functions given on C.

4. Some asymptotic properties of divided differences formed on a Jordan curve. In this section we shall be considering an infinite sequence of divided differences

$$d_1(f \mid z_{11}, z_{12}), d_2(f z_{21}, z_{22}, z_{23}), \cdots, d_n(f \mid z_{n1}, z_{n2}, \cdots, z_n, z_{u+1}), \cdots,$$

formed for a function f given on a Jordan curve C in the z-plane. Do there exist sequences of point sets $S_{n+1} = \{z_{n1}, z_{n2}, \dots, z_n, {}_{n+1}\}, n=1, 2, \dots$ such that $\lim_{n\to\infty} d_n$ exists for all functions f belonging to an interestingly wide class; and if so, what is this limit?

Let D be the region interior to the curve C, let K be the unlimited region exterior to C, and let \overline{D} be $D \cup C$. There exists an analytic function

(4.1)
$$z = \chi(w) = cw + c_0 + \frac{c_1}{w} + \frac{c_2}{w^2} + \cdots, c > 0$$
,

univalent for |w| > 1, which maps |w| > 1 conformally onto K so that the points at infinity in the z-plane and w-plane correspond. According to the Osgood-Taylor-Carathéodory Theorem, $\chi(w)$ can be extended in a continuous and one-to-one manner onto |w| = 1, and $\chi(e^{i\theta}) = \phi(\theta)$ then gives a parametric equation for C of the type considered above in § 3. The number c is called the transfinite diameter (Robin's constant, capacity) of C.

If a function f is analytic on C, it is also analytic in a region (perhaps multiply connected) which contains C in its interior. Let $w = Re^{i\theta}$ in (4.1). There is a largest value of R, say $\rho > 1$, such that f is analytic at every point of the intersection of $C \cup K$ with the region interior to the Jordan curve C_R : $z = \chi(Re^{i\theta}), 0 \le \theta \le 2\pi$. (See [9, p. 79].) A curve such as C_R is called a level curve of the map given by (4.1).

With $z_{nk} = \chi(e^{i\theta_{nk}}), 0 \leq \theta_{nk} < 2\pi$, let $N_n(\theta)$ be the number of elements of the set $\{\theta_{n1}, \theta_{n2}, \dots, \theta_{n,n+1}\}$ falling into the closed interval $[0, \theta]$. The numbers $\theta_{nk}, k = 1, \dots, n+1, n = 1, 2, \dots$ are said to be equidistributed on $[0, 2\pi]$ if $\lim_{n\to\infty} N_n(\theta)/(n+1) = \theta/2\pi$; and when this happens, the corresponding sequence of point sets $S_{n+1}, n = 1, 2, \dots$, is said to be equidistributed on C.

Our first result is as follows:

THEOREM 4.1. Let f be analytic on \overline{D} and let the sequence $\{S_{n+1}\}$ be equidistributed on C. Let ρ be the largest value of |w| in the map (4.1) such that f is analytic interior to the level curve $C_{|w|}$. Then for any R, $1 < R < \rho$, there exists a constant M depending on f, R, and C, but not on n, such that

$$|\, d_{\,_n}(f \,|\, z_{\,_n1},\, \cdots, z_{\,_n},\,_{n+1})\,| \leq rac{M}{(cR)^{n+1}} \;.$$

Thus $\lim_{n\to\infty} c^{n+1}d_n = 0$.

To prove this, we use the formula [5, p. 11]

(4.2)
$$d_n = rac{1}{2\pi i} \int_{\sigma_R} rac{f(t)}{\omega_{n+1}(t)} \, dt, \qquad 0 < R <
ho \ ,$$

where $\omega_{n+1}(z) = (z - z_{n1})(z - z_{n2}) \cdots (z - z_{n,n+1})$. This formula can be used to complete the definition of d_n by continuity for the case of confluent points z_{nk} . We then refer to a classical result of L. Fejér [4], [9, pp. 167 ff]: If $\{S_{n+1}\}$ is equidistributed on a Jordan curve C, then

$$\lim_{n o\infty}\mid \omega_{n+1}(z)\mid^{_{1/(n+1)}}=c\mid w\mid$$
 , $z=\chi(w)$,

uniformly for z on any closed subset of K. This implies that if z lies on $C_{\mathbb{R}}$ and R_1 is such that $1 < R_1 < R$, then for all n sufficiently large,

$$rac{c^{n+1}}{\mid \omega_{n+1}(z)\mid} \leq rac{1}{R_1^{n+1}} \ .$$

Letting M_R be the maximum of |f(z)| on C_R , and L_R be the length of C_R , we appraise (4.2) as follows:

(4.3)
$$c^{n+1} \mid d_n \leq rac{M_R L_R}{2\pi R_1^{n+1}}$$

and the theorem follows from this.

THEOREM 4.2. Let C be rectifiable, let f be analytic on C, and let S_{n+1} be the transform under (4.1) of n+1 distinct points equally spaced on |w| = 1. Then

(4.4)
$$\lim_{n\to\infty} c^{n+1} d_n = \frac{1}{2\pi i} \int_{\sigma} f(t) dt .$$

This is consistent with Theorem 4.1, because the sequence $\{S_{n+1}\}$ in Theorem 4.2 is equidistributed and the integral in (4.4) would be zero if f were analytic on \overline{D} .

To prove the theorem we use a generalization of (4.2),

(4.5)
$$d_n = \frac{1}{2\pi i} \left(\int_{\sigma_R} + \int_{\sigma'} \right) \frac{f(t)}{\omega_{n+1}(t)} dt$$

which is easily established by the calculus of residues. Here C_R , R > 1, is a level curve of (4.1) and C' is a suitably chosen rectifiable curve lying in D. The curves C_R and C' are chosen so that f is analytic on the closed annular region bounded by C_R and C'. Integration on C' is in the opposite sense to that on C_R .

The appraisal given by (4.3) is valid for the first integral in (4.5), and it shows that this integral vanishes in the limit. The following result of the author [2] is available for the second integral: From the hypotheses of Theorem 4.2 on C and S_{n+1} it follows that

$$\lim_{n\to\infty}\omega_{n+1}(z)/c^{n+1}=-1$$

uniformly for z on any closed subset of D. This implies that

$$egin{aligned} &\lim_{n o\infty}&rac{1}{2\pi i}\int_{\sigma'}rac{c^{n+1}f(t)}{\omega_{n+1}(t)}\,dt\ &=rac{1}{2\pi i}\int_{\sigma'}&\lim_{n o\infty}rac{c^{n+1}f(t)}{\omega_{n+1}(t)}\,dt\ &=&-rac{1}{2\pi i}\int_{\sigma'}f(t)\,dt=rac{1}{2\pi i}\int_{\sigma}f(t)\,dt\;, \end{aligned}$$

which completes the proof.

Generalizations of the above theorems to the case in which D is replaced by a finite number of mutually exterior Jordan regions can be developed by the methods to be found in Walsh's book [9, Chap. VII].

The results from which the above two theorems are derived were originally established in studying the convergence of sequences of polynomials found by interpolation to the function f on C. Let $L_{n+1}(z) = L_{n+1}(f; z | S_{n+1})$ be the (unique) polynomial in z of degree at most n which is determined by the condition that it shall coincide with f(z) at each of the points S_{n+1} , assumed to be distinct. Then from the standard formula

$$L_{n+1}(lpha) = \omega_{n+1}(lpha) \sum_{k=1}^{n+1} rac{f(z_{nk})}{\omega_{n+1}'(z_{nk})(lpha-z_{nk})}$$

it is seen by comparison with (1.1) that

$$(4.6) L_{n+1}(g; \alpha \mid S_{n+1}) = \omega_{n+1}(\alpha) d_n(f \mid z_{n1}, \cdots, z_{n,n+1})$$

where $f(z) = g(z)/(\alpha - z)$. The following result of the author [2], [3] is relevant: Let the curve C be such that $\chi'(w)$ is nonvanishing and of bounded variation for |w| = 1. Let g be bounded and integrable in the sense of Riemann on C. Let the points S_{n+1} be the transforms under (4.1) of distinct points equally spaced on the unit circle. Then

$$\lim_{n\to\infty} L_{n+1}(g;\alpha\mid S_n) = \frac{1}{2\pi i} \int_\sigma \frac{g(t)}{t-\alpha} dt$$

uniformly for α on any closed subset of D.

We now write (4.6) in the form

(4.7)
$$c^{n+1}d_n = \left[-\frac{c^{n+1}}{\omega_{n+1}(\alpha)}\right] \left[-L_{n+1}(g; \alpha \mid S_{n+1})\right].$$

If α is a fixed point of D and f is bounded and Riemann integrable on C, then so is g and conversely. We recall that $-\omega_{n+1}(\alpha)/c^{n+1}$ tends to unity at each point of D as n becomes infinite. It follows from these facts that the limiting value of (4.7) is

$$\lim_{n\to\infty} c^{n+1}d_n = \frac{-1}{2\pi i}\int_{\sigma} \frac{g(t)}{t-\alpha} dt = \frac{1}{2\pi i}\int_{\sigma} f(t) dt .$$

We summarize formally:

THEOREM 4.3. If the points S_{n+1} are transforms under (4.1) of distinct points equally spaced on the unit circle, and if C is such that χ' is nonvanishing and of bounded variation for |w| = 1, and if f is

bounded and integrable in the sense of Riemann on C, then

$$\lim_{n\to\infty}c^{n+1}d_n=\frac{1}{2\pi i}\int_{\sigma}f(t)\,dt\;.$$

References

1. J. H. Curtiss, "Interpolation in equidistributed points on the unit circle", to appear in the Pacific Journal of Mathematics.

2. _____, Riemann sums and the fundamental polynomials of Lagrange interpolation" Duke Math. 8 (1941), 634-646.

3. ____, Interpolation in regularly distributed points, Trans. Amer. Math. Soc., 38 (1935), 458-473.

4. L. Fejér, "Interpolation und konforme Abbildung, Gottinger Nachrichten, (1918), 319-331.

5. Milne-Thomson, L. M., The Calculus of Finite Differences, MacMillan and Co., London, 1933.

6. W. E. Sewell, Degree of Approximation by Polynomials in the Complex Domain, Princeton University Press, Princeton, 1942.

7. J. F. Steffensen, Interpolation, Williams and Wilkins, Baltimore, 1927.

8. E. C. Titchmarsh, Theory of Functions, Oxford University Press, Oxford, 1932.

9. J. L. Walsh, Interpolation and Approximation by Rational Functions in the Complex Domain, Second edition, Amer. Math. Soc., Providence, 1956.

UNIVERSITY OF MIAMI

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RALPH S. PHILLIPS Stanford University Stanford, California

M. G. ARSOVE University of Washington Seattle 5, Washington A. L. WHITEMAN University of Southern California Los Angeles 7, California

LOWELL J. PAIGE University of California Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH	D. DERRY	H. L. ROYDEN	E. G. STRAUS
T. M. CHERRY	M. OHTSUKA	E. SPANIER	F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA CALIFORNIA INSTITUTE OF TECHNOLOGY UNIVERSITY OF CALIFORNIA MONTANA STATE UNIVERSITY UNIVERSITY OF NEVADA NEW MEXICO STATE UNIVERSITY OREGON STATE UNIVERSITY UNIVERSITY OF OREGON OSAKA UNIVERSITY UNIVERSITY OF SOUTHERN CALIFORNIA STANFORD UNIVERSITY UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON

AMERICAN MATHEMATICAL SOCIETY CALIFORNIA RESEARCH CORPORATION SPACE TECHNOLOGY LABORATORIES NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics Vol. 12, No. 4 April, 1962

Tsuyoshi Andô, On fundamental properties of a Banach space with a cone	1163
Sterling K. Berberian, A note on hyponormal operators	1171
Errett Albert Bishop, Analytic functions with values in a Frechet space	1177
(Sherman) Elwood Bohn, Equicontinuity of solutions of a quasi-linear	
equation	1193
Andrew Michael Bruckner and E. Ostrow, Some function classes related to the	
class of convex functions	1203
J. H. Curtiss, <i>Limits and bounds for divided differences on a Jordan curve in the complex domain</i>	1217
P. H. Doyle, III and John Gilbert Hocking, <i>Dimensional invertibility</i>	1235
David G. Feingold and Richard Steven Varga, <i>Block diagonally dominant matrices</i>	
and generalizations of the Gerschgorin circle theorem	1241
Leonard Dubois Fountain and Lloyd Kenneth Jackson, A generalized solution of the	
boundary value problem for $y'' = f(x, y, y')$	1251
Robert William Gilmer, Jr., Rings in which semi-primary ideals are primary	1273
Ruth Goodman, <i>K</i> -polar polynomials	1277
Israel Halperin and Maria Wonenburger, On the additivity of lattice	
completeness	1289
Robert Winship Heath, Arc-wise connectedness in semi-metric spaces	1301
Isidore Heller and Alan Jerome Hoffman, On unimodular matrices	1321
Robert G. Heyneman, <i>Duality in general ergodic theory</i>	1329
Charles Ray Hobby, Abelian subgroups of p-groups	1343
Kenneth Myron Hoffman and Hugo Rossi, <i>The minimum boundary for an analytic polyhedron</i>	1347
Adam Koranyi. The Bergman kernel function for tubes over convex cones.	1355
Pesi Rustom Masani and Jack Max Robertson. <i>The time-domain analysis of a</i>	
continuous parameter weakly stationary stochastic process	1361
William Schumacher Massey. <i>Non-existence of almost-complex structures on</i>	
quaternionic projective spaces	1379
Deane Montgomery and Chung-Tao Yang, A theorem on the action of SO(3)	1385
Ronald John Nunke, A note on Abelian group extensions	1401
Carl Mark Pearcy. A complete set of unitary invariants for operators generating	
finite W*-algebras of type I	1405
Edward C. Posner, Integral closure of rings of solutions of linear differential	
equations	1417
Duane Sather, Asymptotics. III. Stationary phase for two parameters with an application to Bessel functions	1423
I Śladkowska. Bounds of analytic functions of two complex variables in domains	1123
with the Beroman-Shilov boundary.	1435
Joseph Gail Stampfli Hyponormal operators	1453
George Gustave Weill Some extremal properties of linear combinations of kernels	1.55
on Riemann surfaces	1459
Edward Takashi Kobayashi, Errata: "A remark on the Niienhuis tensor"	1467