

Pacific Journal of Mathematics

ON THE GENERATION OF DISCONTINUOUS GROUPS

JOSEPH LEHNER

ON THE GENERATION OF DISCONTINUOUS GROUPS

JOSEPH LEHNER

In a paper in this Journal (v. 11, p. 675) M. I. Knopp remarked that $G(j)$, the principal congruence subgroup of level $j \geq 2$ of the modular group, can be generated exclusively by parabolic transformations if and only if it is of genus zero. The following natural generalization is easily proved :

Let Γ be a horocyclic¹ group of genus g . Then Γ possesses a system of generators consisting entirely of parabolic and elliptic elements if and only if $g = 0$.

Knopp's result is a special case, since $G(j)$ has no elliptic substitutions.

For the proof we appeal to the classical result that Γ has a canonical fundamental region whose sides are conjugated by elliptic and parabolic substitutions and $2g$ hyperbolic substitutions $A_1, B_1, \dots, A_g, B_g$ (cf. [1], p. 182 ff). These substitutions generate Γ . If $g = 0$, the hyperbolic ones are absent and the conclusion follows.

Conversely, let Γ be generated by elliptic and parabolic transformations T_1, \dots, T_s . Let the domain of existence of Γ be, for example, the upper half-plane H . Denote by H^+ the union of H and the parabolic cusps of Γ . If $g > 0$ there exists an abelian integral of the first kind, that is, a function F regular in H^+ such that

$$(*) \quad F(Lz) = F(z) + C(L)$$

for all $L \in \Gamma$. Each T_i has a fixed point lying in H^+ . Letting z tend to this fixed point in $(*)$, we see that $C(T_i) = 0$, $i = 1, \dots, s$. Since

$$C(L_1 L_2) = C(L_1) + C(L_2),$$

and the T_i generate Γ , we have

$$C(L) = 0$$

for all $L \in \Gamma$. The abelian integral F has zero periods and is therefore an automorphic function on Γ . Since it is regular in the closed fundamental region, it is a constant. Differentiating, we conclude that there are no abelian differentials of the first kind except 0,

Received June 15, 1962.

¹ A discontinuous group Γ is called horocyclic (*Grenzkreisgruppe*) if there is a fixed disk (or half-plane) preserved by each element of Γ and every boundary point of the disk is a limit point of Γ .

whence Γ is of genus 0. This completes the proof.

That a group of genus 0 cannot always be generated entirely by parabolic elements is shown by the following example, supplied by Morris Newman. Let H be the group generated by $G = G(3)$ and T , where $T\tau = -1/\tau$. Since T is of period 2 and commutes with G , we have

$$H = G + TG.$$

Now G is of genus 0, as is known. Let $f(\tau)$ be a univalent function on G with a simple pole at $\tau_0 \neq i$. Then $f(\tau) + f(-1/\tau)$ is univalent on H , which is therefore of genus 0. A parabolic element P of H cannot lie in TG , for P has trace ± 2 whereas $TG \equiv T \pmod{3}$ has trace divisible by 3. Hence P is in G , and therefore every product of parabolic elements of H is also in G . It follows that H cannot be generated by parabolic elements alone.

Instead of $G(3)$ we could also have used $G(4)$ or $G(5)$.

REFERENCE

1. R. Fricke-F. Klein, *Vorlesungen über die Theorie der automorphen Funktionen*, vol. 1, Teubner, Leipzig, 1897.

NATIONAL BUREAU OF STANDARDS AND MICHIGAN STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RALPH S. PHILLIPS
Stanford University
Stanford, California

M. G. ARSOVE
University of Washington
Seattle 5, Washington

J. DUGUNDJI
University of Southern California
Los Angeles 7, California

LOWELL J. PAIGE
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH
T. M. CHERRY

D. DERRY
M. OHTSUKA

H. L. ROYDEN
E. SPANIER

E. G. STRAUS
F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Pacific Journal of Mathematics

Vol. 13, No. 1

March, 1963

Frantz Woodrow Ashley, Jr., <i>A cone of super-(L) functions</i>	1
Earl Robert Berkson, <i>Some metrics on the subspaces of a Banach space</i>	7
Felix Earl Browder and Walter Strauss, <i>Scattering for non-linear wave equations</i>	23
Edmond Darrell Cashwell and C. J. Everett, <i>Formal power series</i>	45
Frank Sydney Cater, <i>Continuous linear functionals on certain topological vector spaces</i>	65
John Douglas Dixon, <i>General group extensions</i>	73
Robert Pertsch Gilbert, <i>On harmonic functions of four variables with rational p_4-associates</i>	79
Irving Leonard Glicksberg, <i>On convex hulls of translates</i>	97
Simon Hellerstein, <i>On a class of meromorphic functions with deficient zeros and poles</i>	115
Donald William Kahn, <i>Secondary cohomology operations which extend the triple product</i>	125
G. K. Leaf, <i>A spectral theory for a class of linear operators</i>	141
R. Sherman Lehman, <i>Algebraic properties of the composition of solutions of partial differential equations</i>	157
Joseph Lehner, <i>On the generation of discontinuous groups</i>	169
S. P. Lloyd, <i>On certain projections in spaces of continuous functions</i>	171
Fumi-Yuki Maeda, <i>Generalized spectral operators on locally convex spaces</i>	177
Donald Vern Meyer, <i>E^3 modulo a 3-cell</i>	193
William H. Mills, <i>An application of linear programming to permutation groups</i>	197
Richard Scott Pierce, <i>Centers of purity in abelian groups</i>	215
Christian Pommerenke, <i>On meromorphic starlike functions</i>	221
Zalman Rubinstein, <i>Analytic methods in the study of zeros of polynomials</i>	237
B. N. Sahney, <i>On the Nörlund summability of Fourier series</i>	251
Tôru Saitô, <i>Regular elements in an ordered semigroup</i>	263
Lee Meyers Sonneborn, <i>Level sets on spheres</i>	297
Charles Andrew Swanson, <i>Asymptotic estimates for limit point problems</i>	305
Lucien Waelbroeck, <i>On the analytic spectrum of Arens</i>	317
Alvin (Murray) White, <i>Singularities of a harmonic function of three variables given by its series development</i>	321
Kôichi Yamamoto, <i>Decomposition fields of difference sets</i>	337
Chung-Tao Yang, <i>On the action of $SO(3)$ on a cohomology manifold</i>	353