

# Pacific Journal of Mathematics

**ON THE ACTION OF  $SO(3)$  ON A COHOMOLOGY MANIFOLD**

CHUNG-TAO YANG

# ON THE ACTION OF $SO(3)$ ON A COHOMOLOGY MANIFOLD

C. T. YANG

**1. Introduction.** Let the rotation group  $SO(3)$  of the euclidean 3-space act on a locally compact Hausdorff space  $X$  such that the highest dimension of the orbits is 3. Then the following results can be found in Montgomery-Samelson [2].

**THEOREM 1.** *If  $X$  is an integral cohomology  $n$ -manifold and an integral cohomology  $n$ -sphere, then the principal isotropy group is either trivial or contains a dihedral subgroup of order 4.*

**THEOREM 2.** *If  $X$  is the  $n$ -sphere, if the action of  $SO(3)$  on  $X$  is differentiable and if the union of all the singular orbits is of dimension  $< n - 2$ , then the principal isotropy group is trivial.*

The purpose of the present paper is to generalize these theorems.

Basic notations, concepts and theorems which are often used in the study of topological transformation groups will not be given in this paper. Any reader who is not familiar with them may consult [1] or the references given in [1].

Throughout the paper,  $Z$  denotes the ring of integer,  $R$  denotes the field of rational numbers,  $p$  denotes a prime and  $Z_p$  denotes the field of integers mod  $p$ .  $G$  denotes the rotation group  $SO(3)$  of the euclidean 3-space,  $T$  denotes a circle group in  $G$  and  $N$  denotes the normalizer of  $T$ . Notice that  $N/T$  is of order 2 and that every element of  $N - T$  is of order 2. Also notice that the cyclic subgroup of  $T$  of order  $p$ , which we also denote by  $Z_p$ , has  $N$  as its normalizer in  $G$ . As in [1],  $H_c^k(M; L)$  denotes the  $k$ th Alexander-Wallace-Spanier cohomology group of  $M$  with compact support and with coefficients in  $L$ . When  $M$  is compact, it is also written  $H^k(M; L)$ .

## 2. On the action of $N$ .

**LEMMA 1.** *Let  $N$  be the normalizer of a circle group  $T$  in  $SO(3)$  and let  $N$  act on an orientable connected rational cohomology  $m$ -manifold  $Y$  such that*

- (i) *all the orbits are 1-dimensional and*
- (ii)  *$N/T$  acts freely on  $Y/T$ .*

---

Received June 6, 1962. The research is supported in part by the National Science Foundation and in part by the U. S. Army Research Office.

Then  $Y/N$  is a connected rational cohomology  $(m - 1)$ -manifold which is orientable or nonorientable according as the elements of  $N - T$  reverse or preserve the orientation of  $Y$ .

*Proof.* Since all the orbits are 1-dimensional, the sheaf

$$\mathcal{H}^1(Tx; R) = \bigcup H^1(Tx; R)$$

on  $Y/T$  is constant so that  $Y/T$  is an orientable connected rational  $(m - 1)$ -manifold. Since  $N/T$  acts freely on  $Y/T$ ,  $Y/N = (Y/T)/(N/T)$  is a connected rational cohomology  $(m - 1)$ -manifold. Moreover,  $Y/N$  is orientable or nonorientable according as  $N - T$  preserves the orientation of  $Y/T$ .

Let  $h \in N - T$  and let  $y \in Y$ . Then there is a connected slice  $K$  of the transformation group  $(N, Y)$  at  $y$  and a connected neighborhood  $Q$  of the identity in  $T$  such that the map  $Q \times K \rightarrow QK$ , given by  $(g, x) \rightarrow gx$ , is a homeomorphism onto and such that  $hQh^{-1} = Q$ . Clearly  $hK$  is a slice of  $(N, Y)$  at  $hy$  and the map  $Q \times hK \rightarrow QhK (=hQK)$ , given  $(g, x) \rightarrow gx$ , is a homeomorphism onto. Let  $Y, Q, K$  and  $hK$  be oriented such that the maps  $Q \times K \rightarrow Y$  and  $Q \times hK \rightarrow Y$ , given by  $(g, x) \rightarrow gx$ , are orientation-preserving. Since the map  $Q \rightarrow Q$ , given by  $t \rightarrow hth^{-1}$ , reverses the orientation of  $Q$ , it follows that  $h$  reverses the orientation of  $Y$  if and only if  $h: K \rightarrow hK$  is orientation-preserving.

Let  $\pi: Y \rightarrow Y/T$  be the canonical projection. Since the sheaf  $\mathcal{H}^1(Tx; R)$  on  $Y/T$  is constant,  $Y/T$  may be oriented such that the maps  $\pi|_K$  and  $\pi|_{hK}$  are both orientation-preserving. Hence  $h$  reverses the orientation of  $Y$  if and only if  $N - T = hT$  preserves the orientation of  $Y/T$ . This completes the proof of Lemma 1.

**LEMMA 2.** *Let  $X$  be a connected mod 2 cohomology  $n$ -manifold with  $H_c^k(X; Z_2) = 0$  for  $k = n - 1, n - 2$ , and let  $G = \text{SO}(3)$  act on  $X$ . If the principal isotropy group is a finite group of even order, then it contains a dihedral subgroup of order 4. Moreover, the stationary point set of every cyclic subgroup of  $G$  of order 2 is a connected mod 2 cohomology  $(n - 2)$ -manifold and that of every dihedral subgroup of  $G$  of order 4 is a mod 2 cohomology  $(n - 3)$ -manifold.*

*Proof.* Let  $Z_2$  be the cyclic subgroup of  $T$  of order 2. Since the principal isotropy group is a finite group of even order, the stationary point set  $F(Z_2)$  of  $Z_2$  intersects every principal orbit at a 1-dimensional set. Therefore  $F(Z_2)$  is of mod 2 cohomology dimension  $n - 2$  everywhere and hence it is a mod 2 cohomology  $(n - 2)$ -manifold [1; p. 76].

By hypothesis,

$$H_c^n(X; Z_2) = Z_2,$$

$$H_c^k(X; Z_2) = 0 \text{ for } k > n \text{ and for } k = n - 1, n - 2.$$

Using Smith sequence [1; p. 41], one can easily see that

$$H_c^{n-2}(F(Z_2); Z_2) = Z_2.$$

Hence  $F(Z_2)$  is connected.

Let  $D_2$  be a dihedral subgroup of  $N$  of order 4. We assert that the stationary point set  $F(D_2)$  of  $D_2$  is not empty. Assume that the assertion is false; then we have a fibre map

$$\lambda: X \rightarrow G/N$$

given by  $\lambda(gF(Z_2)) = gN, g \in G$ . Therefore we have a spectral sequence  $\{E_r\}$  whose  $E_2$ -term is given by

$$E_2^{s,t} = H^s(G/N; \mathcal{H}_c^t(gF(Z_2); Z_2))$$

and whose  $E_\infty$ -term is associated with  $H_c^*(X; Z_2)$ . Clearly

$$E_2^{1,n-2} = Z_2,$$

$$E_2^{s,t} = 0 \text{ for } s < 0 \text{ and for } s > 2.$$

It follows that

$$E_\infty^{1,n-2} = E_2^{1,n-2} = Z_2.$$

This is impossible as we know that  $H_c^{n-1}(X; Z_2) = 0$ .

By Borel theorem [1; p. 182],  $F(D_2)$  is a mod 2 cohomology  $(n-3)$ -manifold. Since  $F(D_2)$  intersects every orbit at a finite set, it must intersect principal orbits. Hence the principal isotropy group contains a dihedral subgroup of order 4.

**LEMMA 3.** *Let  $N$  be the normalizer of a circle group  $T$  in  $SO(3)$  and let  $Y$  be a connected mod 2 cohomology  $m$ -manifold with  $H_c^{m-1}(Y; Z_2) = 0$ . If  $N$  acts on  $Y$  such that the principal orbits are 1-dimensional and such that for some  $h \in N - T$ , the fixed point set  $F(h)$  of  $h$  is a mod 2 cohomology  $(m-1)$ -manifold, then the stationary point set  $F(N)$  of  $N$  is a mod 2 cohomology  $(m-2)$ -manifold.*

*Proof.* Since the principal orbits are 1-dimensional,  $m \geq 1$  and the stationary point set of  $T$  is of mod 2 cohomology dimension  $\leq m-2$ . Hence for  $m=1$ ,  $F(N) = \phi$  so that our conclusion is trivial.

Let  $m \geq 2$ . Let  $D_2$  be a dihedral subgroup of  $N$  of order 4 and let  $D_2 - T = \{h_1, h_2\}$ . Since all the elements of  $N - T$  are conjugate

to one another in  $N$ , it follows from our hypothesis that the fixed point set  $F(h_i)$  of  $h_i$  is a mod 2 cohomology  $(m-1)$ -manifold,  $i=1, 2$ . By hypothesis,  $H_c^m(Y; Z_2) = Z_2$  and  $H_c^k(Y; Z_2) = 0$  for  $k > m$  and for  $k = m-1$ . We infer that  $H_c^{m-1}(F(h_1); Z_2) = Z_2$  so that  $Y - F(h_1)$  has exactly two components each of which is mapped into the other under  $h_1$ . Similarly  $H_c^{m-1}(F(h_2); Z_2) = Z_2$  so that  $F(h_2)$  is connected. Since  $h_1 h_2 = h_2 h_1$ ,  $h_1 F(h_2) = F(h_2)$ . Hence  $F(h_1) \cap F(h_2) \neq \phi$ .

By Borel theorem,  $F(D_2) = F(h_1) \cap F(h_2)$  is a mod 2 cohomology  $(m-2)$ -manifold. Moreover, in the vicinity of  $F(D_2)$ ,  $F(D_2)$  coincides with the fixed point set of  $h_1 h_2$  which is independent of the choice of  $D_2$  in  $N$ . Hence  $F(N) = F(D_2)$  is a mod 2 cohomology  $(m-2)$ -manifold.

### 3. Main theorems.

**THEOREM 1.** *Let  $X$  be an orientable connected integral cohomology  $n$ -manifold with  $H_c^k(X; Z) = 0$  for  $k = n-1, n-2$  and let  $G = \text{SO}(3)$  act on  $X$ . If the principal isotropy group is finite, then it is either trivial or contains a dihedral subgroup of order 4.*

*Proof.* Suppose that the principal isotropy group does not contain a dihedral subgroup of order 4. Then, by Lemma 2, it is of odd order so that it contains a subgroup  $Z_p$  for some odd prime  $p$ . Let  $F(Z_p)$  be the stationary point set of  $Z_p$  and let  $N$  be the normalizer of  $Z_p$ . Then  $N$  acts on  $F(Z_p)$  and  $F(Z_p)/N$  can be canonically imbedded into  $X/G$ .

Since  $F(Z_p)$  intersects every principal orbit at a 1-dimensional set,  $F(Z_p)$  is a mod  $p$  cohomology  $(n-2)$ -manifold and  $F(Z_p)/N = X/G$ . By hypothesis,  $H_c^n(X; Z_p) = Z_p$  and  $H_c^k(X; Z_p) = 0$  for  $k > n$  and for  $k = n-1, n-2$ . It follows from Smith sequence that

$$H_c^{n-2}(F(Z_p); Z_p) = Z_p$$

so that  $F(Z_p)$  is connected and orientable.

Let  $B$  be the union of all the singular orbits. Then  $F(Z_p) \cap B$  is the stationary point set of  $T$  so that it is of integral cohomology dimension  $\leq n-4$ . Hence

$$Y = F(Z_p) - B$$

is an orientable connected mod  $p$  cohomology  $(n-2)$ -manifold.

Let  $y \in Y$ . It is clear that a slice  $K$  of the transformation group  $(G, X)$  at  $y$  is an integral cohomology  $(n-3)$ -manifold and it is also a slice of  $(N, Y)$  at  $y$ . Since the isotropy group  $N_y$  is finite, there is a neighborhood of  $y$  in  $Y$  homeomorphic to the product of an open interval and  $K$ . Hence  $Y$  is an orientable connected integral coho-

mology  $(n - 2)$ -manifold and consequently an orientable connected rational cohomology  $(n - 2)$ -manifold.

Every  $h \in N - T$  preserves the orientation of  $Y$ . In fact, let  $f: [0, 1] \rightarrow G$  be a map such that

(i)  $f(0)$  is the identity,

(ii)  $f(1) = h$  and

(iii) whenever  $0 \leq t < t' < 1$ ,  $f(t)N \cap f(t')N = \phi$ . Then whenever  $0 \leq t < t' < 1$ ,  $f(t)Y \cap f(t')Y = \phi$ . Since  $f(0)Y = Y = hY = f(1)Y$ ,  $f([0, 1])Y$  is a connected integral cohomology  $(n - 1)$ -manifold. We have seen that  $\bar{Y} - Y = F(Z_p) \cap B$  is of integral cohomology dimension  $\leq n - 4$ . Therefore  $f([0, 1])(\bar{Y} - Y)$  is of integral cohomology dimension  $\leq n - 3$  so that  $X - f([0, 1])(\bar{Y} - Y)$  is an orientable cohomology  $n$ -manifold with  $H_c^{n-1}(X - f([0, 1])(\bar{Y} - Y); Z) = 0$ . Since  $f([0, 1])Y$  is closed in  $X - f([0, 1])(\bar{Y} - Y)$ , it follows that  $f([0, 1])Y$  is orientable. Hence  $h$  preserves the orientation of  $Y$ .

Now we may apply Lemma 1 to  $(N, Y)$  and conclude that  $Y/N$  is a nonorientable connected rational cohomology  $(n - 3)$ -manifold. Since  $F(Z_p)/N = X/G$ ,

$$Y/N = (X - B)/G.$$

However,  $X - B$  is an orientable connected rational cohomology  $n$ -manifold and the sheaf  $\mathcal{H}^3(Gx; R)$  on  $(X - B)/G$  is constant; we infer that  $(X - B)/G$  is orientable. Hence we have arrived at a contradiction. The proof of Theorem 1 is thus completed.

**THEOREM 2.** *Let  $X$  be an orientable connected integral cohomology  $n$ -manifold with  $H_c^k(X; Z) = 0$  for  $k = n - 1, n - 2, n - 3$  and let  $G = SO(3)$  act on  $X$  such that the principal isotropy group is finite. Then one of the following must hold.*

(1) *The principal isotropy group is trivial.*

(2) *The principal isotropy group is the dihedral group of order 4. There exists a 2-dimensional singular orbit and all the 2-dimensional singular orbits are projective planes. Moreover, the union of all the singular orbits is of integral cohomology dimension  $n - 2$ .*

(3) *The principal isotropy group is the icosahedral group. Every singular orbit is a stationary point of  $G$  and the stationary point set of  $G$  is an integral cohomology  $(n - 4)$ -manifold.*

*Proof.* Assume that the principal isotropy group is not trivial. Then, by Theorem 1, it contains a dihedral subgroup of order 4. Let  $Z_2$  be the cyclic subgroup of  $T$  of order 2 and let  $D_2$  be a dihedral subgroup of  $N$  of order 4. We have shown in Lemma 2 that the stationary point set  $F(Z_2)$  of  $Z_2$  is a connected mod 2 cohomology

$(n - 2)$ -manifold and that the stationary point set  $F(D_2)$  of  $D_2$  is a mod 2 cohomology  $(n - 3)$ -manifold. Since  $H_c^n(X; Z_2) = Z_2$  and  $H_c^k(X; Z_2) = 0$  for  $k > n$  and for  $k = n - 1, n - 2, n - 3$ , we infer that  $H_c^{n-3}(F(Z_2); Z_2) = 0$  and that  $H_c^{n-3}(F(D_2); Z_2) = Z_2$  so that  $F(D_2)$  is connected.

The transformation group  $(N, F(Z_2))$  satisfies the hypothesis of Lemma 3. In fact,  $F(Z_2)$  is a connected mod 2 cohomology  $(n - 2)$ -manifold with  $H_c^{n-3}(F(Z_2); Z_2) = 0$ , the principal orbits of  $(N, F(Z_2))$  are 1-dimensional and the fixed point set of  $h \in N - T$  in  $F(Z_2)$  is the stationary point set of the dihedral group generated by  $Z_2$  and  $h$  so that it is a mod 2 cohomology  $(n - 3)$ -manifold. By Lemma 3, the stationary point set  $F(N)$  of  $N$  is a mod 2 cohomology  $(n - 4)$ -manifold.

Suppose first that there exists a 2-dimensional singular orbit  $Gz$ . Since the isotropy group  $G_z$  at  $z$  contains the principal isotropy group which has been shown to contain a dihedral subgroup of order 4, it follows that  $G_z$  is isomorphic to  $N$  so that  $Gz$  is a projective plane. Therefore  $F(N)$  is the stationary point set of  $T$  and hence is an integral cohomology  $(n - 4)$ -manifold. The union of all the singular orbits is  $GF(N)$  which is clearly of integral cohomology dimension  $n - 2$ .

The principal isotropy group is a subgroup of  $G_z$  so that it is a dihedral group containing  $D_2$ . As in the proof of [3; (3.6)], one can easily show that if  $y$  is a point of  $F(D_2)$  such that  $Gy$  is a principal orbit, then the isotropy group  $G_y$  leaves every point of  $F(D_2)$  fixed. If  $G_y$  is not of order 4, then  $G_y$  leaves only one point of  $Gz$  fixed, contrary to the fact that  $F(D_2) \cap G_z$  contains three points.

Suppose next that no singular orbits is 2-dimensional. Then every singular orbit is a stationary point of  $G$ . Hence the union  $B$  of all the singular orbits is the stationary point set of  $T$  which is clearly an integral cohomology  $(n - 4)$ -manifold.

If  $n = 3$ , then  $X$  is an integral cohomology 3-sphere and  $G$  acts transitively on  $X$ . (Here  $H_c^?(X; Z)$  means the reduced group.) Hence the principal isotropy group is the icosahedral group.

If  $n > 3$ , then there is a point  $z$  of  $B$ . As in [1; Chapter XV], the principal orbits are integral cohomology 3-spheres. In fact, there is a neighborhood  $V$  of  $z$  in  $F(D_2)$  invariant under the normalizer  $C$  of  $D_2$ . Since  $V$  is a mod 2 cohomology  $(n - 3)$ -manifold and the stationary point set of  $C$  in  $V$  is a mod 2 cohomology  $(n - 4)$ -manifold, we may choose  $V$  such that  $V - B$  contains exactly two components each of which is a cross-section of  $(G, G(V - B))$ . Hence we may follow the argument of [1; p. 213] to show that the principal orbits are integral cohomology 3-spheres. Consequently the principal isotropy group is the icosahedral group.

REMARK. It is not hard to see that all three cases in Theorem 2 actually occur. In fact, we can have linear actions of  $SO(3)$  on spheres as examples for the first two cases as seen in [3] and a typical example for the third case is seen in [2]. In the third case, the stationary point set of the icosahedral group is not a cohomology manifold for  $n > 3$ ; in fact, it is an integral cohomology  $(n - 3)$ -manifold with the stationary point set of  $G$  as its boundary. Hence the third case never occurs when  $(G, X)$  is a differentiable transformation group and  $n > 3$ .

#### REFERENCES

1. A. Borel et al., *Seminar on Transformation Groups*, Annals of Mathematics Studies No. 46, Princeton University Press, 1960.
2. D. Montgomery and H. Samelson, *On the action of  $SO(3)$  on* Pacific J. Math., **12** (1962),
3. D. Montgomery and C. T. Yang, *A theorem on the action of  $SO(3)$* , Pacific J. Math., **12** (1962) 000-000.

UNIVERSITY OF PENNSYLVANIA AND  
THE INSTITUTE FOR ADVANCED STUDY





# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RALPH S. PHILLIPS

Stanford University  
Stanford, California

M. G. ARSOVE

University of Washington  
Seattle 5, Washington

J. DUGUNDJI

University of Southern California  
Los Angeles 7, California

LOWELL J. PAIGE

University of California  
Los Angeles 24, California

## ASSOCIATE EDITORS

E. F. BECKENBACH

T. M. CHERRY

D. DERRY

M. OHTSUKA

H. L. ROYDEN

E. SPANIER

E. G. STRAUS

F. WOLF

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

\* \* \*

AMERICAN MATHEMATICAL SOCIETY  
CALIFORNIA RESEARCH CORPORATION  
SPACE TECHNOLOGY LABORATORIES  
NAVAL ORDNANCE TEST STATION

Frantz Woodrow Ashley, Jr., <i>A cone of super-(L) functions</i> .....	1
Earl Robert Berkson, <i>Some metrics on the subspaces of a Banach space</i> ....	7
Felix Earl Browder and Walter Strauss, <i>Scattering for non-linear wave equations</i> .....	23
Edmond Darrell Cashwell and C. J. Everett, <i>Formal power series</i> .....	45
Frank Sydney Cater, <i>Continuous linear functionals on certain topological vector spaces</i> .....	65
John Douglas Dixon, <i>General group extensions</i> .....	73
Robert Pertsch Gilbert, <i>On harmonic functions of four variables with rational <math>p_4</math>-associates</i> .....	79
Irving Leonard Glicksberg, <i>On convex hulls of translates</i> .....	97
Simon Hellerstein, <i>On a class of meromorphic functions with deficient zeros and poles</i> .....	115
Donald William Kahn, <i>Secondary cohomology operations which extend the triple product</i> .....	125
G. K. Leaf, <i>A spectral theory for a class of linear operators</i> .....	141
R. Sherman Lehman, <i>Algebraic properties of the composition of solutions of partial differential equations</i> .....	157
Joseph Lehner, <i>On the generation of discontinuous groups</i> .....	169
S. P. Lloyd, <i>On certain projections in spaces of continuous functions</i> .....	171
Fumi-Yuki Maeda, <i>Generalized spectral operators on locally convex spaces</i> .....	177
Donald Vern Meyer, <i><math>E^3</math> modulo a 3-cell</i> .....	193
William H. Mills, <i>An application of linear programming to permutation groups</i> .....	197
Richard Scott Pierce, <i>Centers of purity in abelian groups</i> .....	215
Christian Pommerenke, <i>On meromorphic starlike functions</i> .....	221
Zalman Rubinstein, <i>Analytic methods in the study of zeros of polynomials</i> .....	237
B. N. Sahney, <i>On the Nörlund summability of Fourier series</i> .....	251
Tôru Saitô, <i>Regular elements in an ordered semigroup</i> .....	263
Lee Meyers Sonneborn, <i>Level sets on spheres</i> .....	297
Charles Andrew Swanson, <i>Asymptotic estimates for limit point problems</i> .....	305
Lucien Waelbroeck, <i>On the analytic spectrum of Arens</i> .....	317
Alvin (Murray) White, <i>Singularities of a harmonic function of three variables given by its series development</i> .....	321
Kôichi Yamamoto, <i>Decomposition fields of difference sets</i> .....	337
Chung-Tao Yang, <i>On the action of <math>SO(3)</math> on a cohomology manifold</i> .....	353