

Pacific Journal of Mathematics

**MAXIMUM MODULUS ALGEBRAS AND LOCAL
APPROXIMATION IN C^n**

ALBERT EMERSON HURD

MAXIMUM MODULUS ALGEBRAS AND LOCAL APPROXIMATION IN C^n

A. E. HURD

1. In [4] W. Rudin established an important result concerning maximum modulus algebras A of continuous complex-valued functions defined on the closure K of a Jordan domain in the complex plane (see also [5]). Rudin's result states, under the assumptions (a) A contains a function \mathcal{P} which is schlicht on K , and (b) A contains a non-constant function ϕ which is analytic in the interior, $\text{int } K$, of K , that every function in A is analytic in $\text{int } K$. In this note we will establish conditions under which assumption (b) alone yields the desired conclusion in a slightly more general setting. We assume that K is a compact set, with interior, of a Riemann surface, but also assume that $\text{int } K$ is essentially open in the maximal ideal space Σ_A of A (A being regarded as a Banach algebra with the sup norm $\|f\| = \sup_{p \in K} |f(p)|$; see [2]). This means that each point of $\text{int } K$, excepting a set of points having no limit point in $\text{int } K$, has a neighborhood in $\text{int } K$ which is open in Σ_A under the natural mapping of K into Σ_A . Under these assumptions it is easy to show, using the Local Maximum Modulus Principle of H. Rossi [3; Theorem 6.1] and Rudin's results, that (b) is sufficient to guarantee that A consists only of analytic functions. Our main purpose, however, is to establish the result by a geometric method, independent of Rudin's work, which is based on an appropriate local approximation in C^n . Unfortunately the geometric approach being used here only allows us to make the desired conclusion for twice continuously differentiable functions in A whereas the use of Rudin's results would give a proof valid for any function in A . However it is hoped that our method will be of some interest in itself.

The basic idea of the proof is as follows. For simplicity let K be the unit circle $\{z \in C: |z| \leq 1\}$ in the complex plane, and let f and g be nonconstant functions in the maximum modulus algebra A . Suppose that $\Sigma_A = K$. Use f and g to map K into C^2 (the space of 2 complex variables) in the obvious way. If f and g are twice continuously differentiable in the neighborhood of a given point in $\text{int } K$ then the image of this neighborhood in C^2 will be a two (real) dimensional surface possessing a tangent plane at the image p of the point. Let π be the two (real) dimensional tangent plane to this surface at p . If this plane is nonanalytic (Definition 1) then we can find a polynomial in the coordinates w_1 and w_2 of C^2 which locally peaks [3] at p when

Received November 1, 1962. This research was (partially) supported by the Air Force Office of Scientific Research

restricted to the surface. The results of Rossi and the Arens-Calderon Theorem [1] then show that this will contradict the maximum modulus property. Thus π cannot be nonanalytic. This gives a relation between the complex derivatives of f and g which, in particular, implies that both functions are analytic at the pre-image of p if one of them is analytic there. In § 2 the essential geometric lemma is established and in § 3 it is used to prove the main result.

2. Let $F: M \rightarrow R^n$ be an immersion (a regular map in the sense of Whitney [6]) of a two (real) dimensional twice continuously differentiable manifold M into real Euclidean n -space R^n . Let $p \in M$ and let (U, h) give local coordinates about p , where U is an open set in M , and h is a homeomorphism from U onto $D = \{(u, v) \in R^2: u^2 + v^2 < 1\}$, with $h(p) = (0, 0)$. If x_j ($j = 1, \dots, n$) is a coordinate function in R^n then the functions $\phi_j(u, v) = x_j \circ F \circ h^{-1}(u, v)$ ($j = 1, \dots, n$) are differentiable and give a map $\Phi: D \rightarrow R^n$ defined by $\Phi(u, v) = (\phi_1(u, v), \dots, \phi_n(u, v))$. Since F is an immersion, the 2 by n matrix

$$\left(\frac{\partial \phi_j}{\partial u}, \frac{\partial \phi_j}{\partial v} \right) = \begin{pmatrix} \frac{\partial \phi_1}{\partial u} & \frac{\partial \phi_1}{\partial v} \\ \vdots & \vdots \\ \frac{\partial \phi_n}{\partial u} & \frac{\partial \phi_n}{\partial v} \end{pmatrix}$$

has rank 2 and the mapping Φ is one-to-one in some disc $V = \{(u, v) \in R^2: u^2 + v^2 < r^2 < 1\}$. Further, the set $\Phi(V)$ is a surface element having a tangent plane at $\Phi(0, 0)$.

We can suppose for our purposes that $\Phi(0, 0)$ is the origin 0 in R^n . The tangent plane π to $\Phi(V)$ at 0 is then given parametrically by

$$(2.1) \quad x_j = \frac{\partial \phi_j}{\partial u} u + \frac{\partial \phi_j}{\partial v} v \quad (j = 1, \dots, n),$$

where the derivatives are evaluated at $u = v = 0$. A change of local parameters from u and v to $u' = u'(u, v)$ and $v' = v'(u, v)$ with $\partial(u', v')/\partial(u, v) \neq 0$ (the inverse transformation being given by $u = u(u', v')$ and $v = v(u', v')$ in some neighborhood of $u = v = 0$) yield new functions $\phi'_j(u', v') = \phi_j(u(u', v'), v(u', v'))$ and a new parametrization of the tangent plane, namely,

$$\begin{aligned} x_j &= \frac{\partial \phi'_j}{\partial u'} u' + \frac{\partial \phi'_j}{\partial v'} v' \\ &= \left(\frac{\partial \phi_j}{\partial u} \frac{\partial u}{\partial u'} + \frac{\partial \phi_j}{\partial v} \frac{dv}{\partial u'} \right) u' + \left(\frac{\partial \phi_j}{\partial u} \frac{\partial u}{\partial v'} + \frac{\partial \phi_j}{\partial v} \frac{\partial v}{\partial v'} \right) v' \end{aligned} \quad (j = 1, \dots, n).$$

Note that the rank of the matrix $(\partial\phi_j/\partial u, \partial\phi_j/\partial v)$ is the same as that of $(\partial\phi'_j/\partial u', \partial\phi'_j/\partial v')$ since $\partial(u', v')/\partial(u, v) \neq 0$.

Now u and v parametrize both the surface element $\Phi(V)$ and the tangent plane (given by (2.1)). Let η_j and η'_j ($j = 1, 2, \dots, n$) denote the coordinates in R^n of the points B and B' on π and $\Phi(V)$, respectively, corresponding to the parameters u and v ($u^2 + v^2 < r^2$). For sufficiently small u and v ,

$$\eta'_j = \frac{\partial\phi_j}{\partial u} u + \frac{\partial\phi_j}{\partial v} v + \frac{1}{2} \left(\frac{\partial^2\phi_j}{\partial u^2} u^2 + 2 \frac{\partial^2\phi_j}{\partial u\partial v} uv + \frac{\partial^2\phi_j}{\partial v^2} v^2 \right)$$

where the first derivatives are evaluated at $u = v = 0$ and the second derivatives are evaluated at $u' = \theta u, v' = \theta v$ for some θ satisfying $0 < \theta < 1$. Since M is twice continuously differentiable, the second derivatives of ϕ_j are bounded in absolute value in some sufficiently small neighborhood of $(0, 0)$ and we obtain

$$\sum_1^m (\eta_j - \eta'_j)^2 \leq K(|u| + |v|)^4$$

and so

$$(2.2) \quad |\eta_j - \eta'_j| \leq L(|u| + |v|)^2$$

where K and L are constants depending on these bounds and on n , and u and v are sufficiently small. These estimates will be used later.

Suppose now that $n = 2m$. One can define complex coordinates $w_j = x_{2j-1} + ix_{2j}$ making R^n into complex Euclidean space C^m . Also the (u, v) -plane can be formally complexified by writing $z = u + iv, \bar{z} = u - iv$. We then have a mapping $\Psi: V \rightarrow C^m$ defined by $\Psi(z, \bar{z}) = (w_1, \dots, w_m)$ where

$$w_j = \Psi_j(z, \bar{z}) = \phi_{2j-1} \left(\frac{z + \bar{z}}{2}, \frac{z - \bar{z}}{2i} \right) + i\phi_{2j} \left(\frac{z + \bar{z}}{2}, \frac{z - \bar{z}}{2i} \right) \quad (j = 1, \dots, m).$$

An elementary computation shows that in this formalism the tangent plane π to $\Psi(V)$ at the origin 0 is given parametrically by

$$w_j = \frac{\partial\Psi_j}{\partial z} z + \frac{\partial\Psi_j}{\partial \bar{z}} \bar{z} \quad (j = 1, \dots, m)$$

where the derivatives are evaluated at $z = \bar{z} = 0$. Furthermore, under a change of local coordinates in the parameter plane from z and \bar{z} to $z' = u' + iv'$ and $\bar{z}' = u' - iv'$, the tangent plane is given parametrically by

$$\begin{aligned} w_j &= \frac{\partial\Psi'_j}{\partial z'} z' + \frac{\partial\Psi'_j}{\partial \bar{z}'} \bar{z}' \\ &= \left(\frac{\partial\Psi_j}{\partial z} \frac{\partial z}{\partial z'} + \frac{\partial\Psi_j}{\partial \bar{z}} \frac{\partial \bar{z}}{\partial z'} \right) z' + \left(\frac{\partial\Psi_j}{\partial z} \frac{\partial z}{\partial \bar{z}'} + \frac{\partial\Psi_j}{\partial \bar{z}} \frac{\partial \bar{z}}{\partial \bar{z}'} \right) \bar{z}' \end{aligned}$$

where $\Psi'_j(z', \bar{z}') = \Psi_j(z(z'), \bar{z}(\bar{z}'))$. Since, as a short calculation shows,

$$\frac{\partial(z, \bar{z})}{\partial(z', \bar{z}')} = \frac{\partial(u, v)}{\partial(u', v')} \neq 0,$$

the complex rank of $(\partial\Psi_j/\partial z, \partial\Psi_j/\partial \bar{z})$ remains unaffected by a parameter change. We now make the following definition.

Definition 1. The two (real) dimensional plane in C^m defined parametrically by $w_j = \alpha_j z + \beta_j \bar{z}$ ($j = 1, 2, \dots, m$) is said to be *non-analytic* if the rank of the 2 by m (complex) matrix (α_j, β_j) is 2.

The preceding remarks show that if π is nonanalytic in one coordinate parametrization then it remains so under any change of coordinates in the parameter plane. We want to establish the following.

LEMMA. *Suppose the tangent plane π to $\Psi(V)$ at the origin in C^m is nonanalytic. Then there is a polynomial in the coordinates w_j whose absolute value takes on a local maximum at the origin when restricted to $\Psi(V)$.*

Proof. Since π is nonanalytic there exist new coordinates $w'_i = \sum_{j=1}^m \gamma_{ij} w_j$ ($i = 1, \dots, m$), where the matrix (γ_{ij}) is nonsingular, such that in the w'_i -coordinates π is given parametrically by $w'_1 = z, w'_2 = \bar{z}$, and $w'_j = 0$ ($m \geq j \geq 3$). Now let B and B' be points on π and $\Psi(V)$, respectively, corresponding to the parameters u and v . Let η_j and η'_j ($j = 1, \dots, 2n$) be the real coordinates of B and B' (with C^m regarded as R^{2m}) in the new coordinate system. Clearly $\eta_1 = u, \eta_2 = v, \eta_3 = u, \eta_4 = -v$, and $\eta_j = 0$ for $5 \leq j \leq 2n$. Let $\eta'_j - \eta_j = \varepsilon_j$ ($j = 1, \dots, 4$).

Now consider the function $P(w_i) = 1 - w'_1 w'_2$ (a polynomial in w_1, \dots, w_m). When restricted to π , $P(w_i)$ is real-valued and has a maximum in absolute value at the origin. We would like to show that $|P(w_i)|$ also has a local maximum at the origin when restricted to a sufficiently small neighborhood of the origin on $\Psi(V)$. This will be true essentially because π has a contact of order at least 1 with $\Psi(V)$ at the origin (here we will use the estimates (2.2)).

We have, at the point B' ,

$$\begin{aligned} |P(B')|^2 &= |1 - (\eta'_1 + i\eta'_2)(\eta'_3 + i\eta'_4)|^2 \\ &= 1 - 2(u^2 + v^2) + (u^2 + v^2) + 2Q(u, v)[u^2 + v^2 - 1] \\ &\quad + [Q(u, v)]^2 + [u(\varepsilon_2 + \varepsilon_4) + v(\varepsilon_3 - \varepsilon_1)]^2 \end{aligned}$$

where

$$Q(u, v) = u(\varepsilon_1 + \varepsilon_3) + v(\varepsilon_2 - \varepsilon_4) + \varepsilon_1 \varepsilon_3 - \varepsilon_2 \varepsilon_4.$$

Using inequalities (2.2) for $|\varepsilon_j|$ ($j = 1, \dots, 4$) we obtain

$$\begin{aligned} |P(B')|^2 &\leq 1 - 2(u^2 + v^2) + M(|u| + |v|)^3 \\ &\leq 1 - [1 - M(|u| + |v|)](|u| + |v|)^2 \end{aligned}$$

for u and v sufficiently small and some constant M . If $|u| + |v| < 1/M$ we see that $|P(B')| < 1$ unless $B' = 0$.

3. Let K be a compact subset, with nonempty interior, of a Riemann surface M .

DEFINITION 2. An algebra A of continuous functions on K is said to be a maximum modulus algebra on K if for every $f \in A$ there is a point p on the boundary ∂K of K such that $|f(q)| \leq |f(p)|$ for all $q \in K$.

As remarked in [4], we can suppose without loss of generality that A is uniformly closed and contains the constants and so is a Banach algebra with identity and norm $\|f\| = \sup_{p \in K} |f(p)|$. It is well known that there is a natural continuous mapping $i: K \rightarrow \Sigma_A$, where Σ_A is the maximal ideal space of A (with the usual Gelfand topology), defined by point evaluation (which is not 1 : 1 unless A separates points in Σ_A).

THEOREM. Let A be a uniformly closed algebra of continuous functions, containing the constants, on the compact subset K (with nonempty interior) of the Riemann surface M . Suppose that there is a set D of points in $\text{int } K$ having no limit point in $\text{int } K$, such that each $p \in \text{int } K - D$ has a neighborhood U for which $i(U)$ is open in Σ_A . Suppose further that A contains one nonconstant analytic function $g = g^1 + ig^2$. Then any function $f = f^1 + if^2$ in A such that f^1 and f^2 are twice continuously differentiable is analytic in $\text{int } K$.

Proof. Let S be the discrete subset of $\text{int } K$ on which the differential dg vanishes. For any point p in $\text{int } K - S$ there is a neighborhood containing p and contained in $\text{int } K - S$ and in which g is one-to-one. Thus for any point $p \in \text{int } K - (D \cup S)$ there exists a neighborhood U containing p which is mapped homeomorphically by i onto an open set W in Σ_A and hence local coordinates in U may be transferred to W . Define the mapping $F: \Sigma_A \rightarrow C^2$ by $F(q') = (f(q'), g(q'))$, $q' \in \Sigma_A$ (where we have used the letters f and g to denote the extension, via the Gelfand representation, of f and g , defined on $i(K)$, to Σ_A). For any point q' in W we have $f(q') = f(i^{-1}(q'))$ and $g(q') = g(i^{-1}(q'))$ so that F can be regarded as a mapping defined on U by $F(q) = (f(q), g(q))$, $q \in U$. F defines an immersion of W since in the local coordinates $z = u + iv$ the matrix

$$\begin{pmatrix} f_u^1 & f_u^2 & g_u^1 & g_u^2 \\ f_v^1 & f_v^2 & g_v^1 & g_v^2 \end{pmatrix}$$

(here the subscripts u and v denote partial differentiation) is of rank 2 due to the nonvanishing of the differential $dg = \partial g/\partial z dz$ —apply the Cauchy-Riemann equations to the matrix

$$\begin{pmatrix} g_u^1 & g_u^2 \\ g_v^1 & g_v^2 \end{pmatrix}.$$

Since A contains the constants we can suppose without loss of generality that $F(p)$ is the origin 0 in C^2 . We have thus a mapping $F: \Sigma_A \rightarrow C^2$ which maps a neighborhood W of $i(p)$ onto a two-dimensional surface element $F(W)$ having a tangent plane π at 0.

We now note that π cannot be nonanalytic. For if this were the case then by the lemma of § 2 there would be a polynomial in the coordinates w_1 and w_2 of C^2 taking on a local maximum in absolute value at 0 when restricted to $F(W)$. By the Arens-Calderon theorem [1; Theorem 3.3] there would then be a function $k \in A$ taking on a local maximum at $i(p)$, and finally, by Rossi's Local Peak-Point Theorem [3, Theorem 4.1] there would be a function $\tilde{k} \in A$ taking on its maximum value exactly at $i(p)$, contradicting the fact that A is a maximum modulus algebra.

Thus the rank of

$$\begin{pmatrix} \frac{\partial g}{\partial z} & 0 \\ \frac{\partial f}{\partial z} & \frac{\partial f}{\partial \bar{z}} \end{pmatrix}$$

(the derivatives being evaluated in the local coordinates at p) must be 1 and this implies that $\partial f/\partial \bar{z} = 0$. The same conclusion could be drawn for any $p \in \text{int } K - (D \cup S)$ and so by the theorem of Riemann on removable singularities, f is analytic in $\text{int } K$.

REFERENCES

1. R. Arens and A. P. Calderon, *Analytic functions of several Banach algebra elements*, Ann. of Math., **62** (1955), 204-216.
2. I. M. Gelfand, A. Raikov and G. Silov, *Commutative normed rings*, Uspehi Mat. Nauk (N.S.) 1, No. 2 (12) (1946), pp. 48-146 (Amer. Math. Soc. Translations, Ser. 2, Vol. 5, pp. 115-220).
3. H. Rossi, *The local maximum modulus principle*, Ann. of Math., **72** (1960), pp. 1-11.
4. W. Rudin, *Analyticity, and the maximum modulus principle*, Duke Math. J., **20** (1953), 449-457.
5. ———, *On the structure of maximum modulus algebras*, Proc. Amer. Math. Soc., **9** (1958), 708-712.
6. H. Whitney, *Differentiable manifolds*, Ann. of Math., **37** (1936), 645-680.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RALPH S. PHILLIPS

Stanford University
Stanford, California

M. G. ARSOVE

University of Washington
Seattle 5, Washington

J. DUGUNDJI

University of Southern California
Los Angeles 7, California

LOWELL J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

T. M. CHERRY

D. DERRY

M. OHTSUKA

H. L. ROYDEN

E. SPANIER

E. G. STRAUS

F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 13, No. 2

April, 1963

Rafael Artzy, <i>Solution of loop equations by adjunction</i>	361
Earl Robert Berkson, <i>A characterization of scalar type operators on reflexive Banach spaces</i>	365
Mario Borelli, <i>Divisorial varieties</i>	375
Raj Chandra Bose, <i>Strongly regular graphs, partial geometries and partially balanced designs</i>	389
R. H. Bruck, <i>Finite nets. II. Uniqueness and imbedding</i>	421
L. Carlitz, <i>The inverse of the error function</i>	459
Robert Wayne Carroll, <i>Some degenerate Cauchy problems with operator coefficients</i>	471
Michael P. Drazin and Emilie Virginia Haynsworth, <i>A theorem on matrices of 0's and 1's</i>	487
Lawrence Carl Eggan and Eugene A. Maier, <i>On complex approximation</i>	497
James Michael Gardner Fell, <i>Weak containment and Kronecker products of group representations</i>	503
Paul Chase Fife, <i>Schauder estimates under incomplete Hölder continuity assumptions</i>	511
Shaul Foguel, <i>Powers of a contraction in Hilbert space</i>	551
Neal Eugene Foland, <i>The structure of the orbits and their limit sets in continuous flows</i>	563
Frank John Forelli, Jr., <i>Analytic measures</i>	571
Robert William Gilmer, Jr., <i>On a classical theorem of Noether in ideal theory</i>	579
P. R. Halmos and Jack E. McLaughlin, <i>Partial isometries</i>	585
Albert Emerson Hurd, <i>Maximum modulus algebras and local approximation in C^n</i>	597
James Patrick Jans, <i>Module classes of finite type</i>	603
Betty Kvarda, <i>On densities of sets of lattice points</i>	611
H. Larcher, <i>A geometric characterization for a class of discontinuous groups of linear fractional transformations</i>	617
John W. Moon and Leo Moser, <i>Simple paths on polyhedra</i>	629
T. S. Motzkin and Ernst Gabor Straus, <i>Representation of a point of a set as sum of transforms of boundary points</i>	633
Rajakularaman Ponnuswami Pakshirajan, <i>An analogue of Kolmogorov's three-series theorem for abstract random variables</i>	639
Robert Ralph Phelps, <i>Čebyšev subspaces of finite codimension in $C(X)$</i>	647
James Dolan Reid, <i>On subgroups of an Abelian group maximal disjoint from a given subgroup</i>	657
William T. Reid, <i>Riccati matrix differential equations and non-oscillation criteria for associated linear differential systems</i>	665
Georg Johann Rieger, <i>Some theorems on prime ideals in algebraic number fields</i>	687
Gene Fuerst Rose and Joseph Silbert Ullian, <i>Approximations of functions on the integers</i>	693
F. J. Sansone, <i>Combinatorial functions and regressive isols</i>	703
Leo Sario, <i>On locally meromorphic functions with single-valued moduli</i>	709
Takayuki Tamura, <i>Semigroups and their subsemigroup lattices</i>	725
Pui-kei Wong, <i>Existence and asymptotic behavior of proper solutions of a class of second-order nonlinear differential equations</i>	737
Fawzi Mohamad Yaqub, <i>Free extensions of Boolean algebras</i>	761