

Pacific Journal of Mathematics

SIMPLE PATHS ON POLYHEDRA

JOHN W. MOON AND LEO MOSER

SIMPLE PATHS ON POLYHEDRA

J. W. MOON and L. MOSER

In Euclidean d -space ($d \geq 3$) consider a convex polytope whose n ($n \geq d + 1$) vertices do not lie in a $(d - 1)$ -space. By the "path length" of such a polytope is meant the maximum number of its vertices which can be included in any single simple path, i.e., a path along its edges which does not pass through any given vertex more than once. Let $p(n, d)$ denote the minimum path length of all such polytopes of n vertices in d -space. Brown [1] has shown that $p(n, 3) \leq (2n + 13)/3$ and Grünbaum and Motzkin [3] have shown that $p(n, d) < 2(d - 2)n^\alpha$ for some $\alpha < 1$, e.g., $\alpha = 1 - 2^{-19}$ and they have indicated how this last value may be improved to $\alpha = 1 - 2^{-16}$. The main object of this note is to derive the following result which, for sufficiently large values of n , represents an improvement upon the previously published bounds.

THEOREM.

$$p(n, d) < (2d + 3)((1 - 2/(d + 1))n - (d - 2))^{\log_2/\log d} - 1 < 3d n^{\log_2/\log d}.$$

When $d = 3$ the example we construct to imply our bound is built upon a tetrahedron which we denote by G_0 . Its 4 vertices, which will be called the 0th stage vertices, can all be included in a single simple path. Upon each of the 4 triangular faces of G_0 erect a pyramid in such a way that the resulting solid, G_1 , is a convex polyhedron with 12 triangular faces. This introduces 4 more vertices, the 1st stage vertices, which can be included in a single simple path involving all 8 vertices of G_1 . We may observe that it is impossible for a path to go from a 1st stage vertex to another 1st stage vertex without first passing through a 0th stage vertex.

The convex polyhedron G_2 is formed by erecting pyramids upon all the faces of G_1 . Of the 12 2nd stage vertices thus introduced at most 9 can be included in any single simple path since, as before, no path can join two 2nd stage vertices without passing through an intermediate vertex of a lower stage and there are only 8 such vertices available.

The procedure continues as follows: the convex polyhedron G_k , $k \geq 2$, is formed by erecting pyramids upon the $4 \cdot 3^{k-1}$ triangular faces of G_{k-1} . Making repeated use of the fact that the method of construction makes it impossible for a path to join two vertices of the j th stage, $j \geq 2$, without first passing through at least one vertex of a lower stage we find that at most $9 \cdot 2^{j-2}$ of the $4 \cdot 3^{j-1}$ vertices of the

j th stage, $j = 2, 3, \dots, k$, can be included in a single simple path along the edges of G_k . This and the earlier remarks imply that G_k , $k \geq 1$, has $2 \cdot 3^k + 2$ vertices and at most $9 \cdot 2^{k-1} - 1$ of these can be included in a single simple path.

For any integer $n > 4$ let k be the unique integer such that

$$(1) \quad 2 \cdot 3^k + 2 < n \leq 2 \cdot 3^{k+1} + 2.$$

Next consider the convex polyhedron with n vertices which can be obtained by erecting pyramids upon $n - (2 \cdot 3^k + 2)$ faces of G_k . Then, from considerations similar to those given before, it follows, using (1), that

$$(2) \quad p(n, 3) \leq 9 \cdot 2^k - 1 < 9((n - 2)/2)^{\log_2 / \log_3} - 1.$$

This suffices to complete the proof of the theorem when $d = 3$ since the result is trivially true when $n = 4$.

In the general case the construction starts with a d -dimensional simplex. Upon each of its $(d - 1)$ -dimensional faces is formed another d -dimensional simplex by the introduction of a new vertex on the side of the face opposite to the rest of the original simplex in such a way that the resulting polytope is convex. This process is repeated and the rest of the argument is completely analogous to that given for the case $d = 3$. It should be pointed out that the result of Grünbaum and Motzkin holds even for graphs all of whose vertices, but for a bounded number are incident with 3 edges, while in the polytopes described above the distribution of valences is quite different.

In closing we remark that the path length of any 3-dimensional convex polyhedron with n vertices is certainly greater than

$$(\log_2 n / \log_2 \log_2 n) - 1.$$

Suppose that there exists a vertex, q say, upon which at least $\log_2 n / \log_2 \log_2 n$ edges are incident. Let the vertices at the other ends of these edges be p_1, p_2, \dots, p_t , arranged in counterclockwise order. Each pair, (p_i, p_{i+1}) , $i = 1, \dots, t - 1$, of successive vertices in this sequence determines a unique polygonal face containing the edges $\overline{p_{i+1}q}$ and $\overline{qp_i}$. Traversing this face in a counterclockwise sense gives a path from p_i to p_{i+1} involving at least one edge. Since these faces all lie in different planes it is not difficult to see that these paths may be combined to give a simple path from q to p_1 to p_t whose length is at least $t \geq \log_2 n / \log_2 \log_2 n$. If there is no vertex upon which this many edges are incident then the required result follows from the type of argument used by Dirac [2; Theorem 5] in showing that the path length is at least of the magnitude of $\log n$ if only a bounded number of edges are incident upon any vertex.

BIBLIOGRAPHY

1. T. A. Brown, *Simple paths on convex polyhedra*, Pacific J. Math., **11** (1961), 1211-1214.
2. G. A. Dirac, *Some theorems on abstract graphs*, Proc. London Math. Soc., (3) **2** (1952), 69-81.
3. B. Grünbaum and T. S. Motzkin, *Longest simple paths in polyhedral graphs*, J. London Math. Soc., **37** (1962), 152-160.

UNIVERSITY OF ALBERTA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RALPH S. PHILLIPS

Stanford University
Stanford, California

M. G. ARSOVE

University of Washington
Seattle 5, Washington

J. DUGUNDJI

University of Southern California
Los Angeles 7, California

LOWELL J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

T. M. CHERRY

D. DERRY

M. OHTSUKA

H. L. ROYDEN

E. SPANIER

E. G. STRAUS

F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Rafael Artzy, <i>Solution of loop equations by adjunction</i>	361
Earl Robert Berkson, <i>A characterization of scalar type operators on reflexive Banach spaces</i>	365
Mario Borelli, <i>Divisorial varieties</i>	375
Raj Chandra Bose, <i>Strongly regular graphs, partial geometries and partially balanced designs</i>	389
R. H. Bruck, <i>Finite nets. II. Uniqueness and imbedding</i>	421
L. Carlitz, <i>The inverse of the error function</i>	459
Robert Wayne Carroll, <i>Some degenerate Cauchy problems with operator coefficients</i>	471
Michael P. Drazin and Emilie Virginia Haynsworth, <i>A theorem on matrices of 0's and 1's</i>	487
Lawrence Carl Eggan and Eugene A. Maier, <i>On complex approximation</i>	497
James Michael Gardner Fell, <i>Weak containment and Kronecker products of group representations</i>	503
Paul Chase Fife, <i>Schauder estimates under incomplete Hölder continuity assumptions</i>	511
Shaul Foguel, <i>Powers of a contraction in Hilbert space</i>	551
Neal Eugene Foland, <i>The structure of the orbits and their limit sets in continuous flows</i>	563
Frank John Forelli, Jr., <i>Analytic measures</i>	571
Robert William Gilmer, Jr., <i>On a classical theorem of Noether in ideal theory</i>	579
P. R. Halmos and Jack E. McLaughlin, <i>Partial isometries</i>	585
Albert Emerson Hurd, <i>Maximum modulus algebras and local approximation in C^n</i>	597
James Patrick Jans, <i>Module classes of finite type</i>	603
Betty Kvarda, <i>On densities of sets of lattice points</i>	611
H. Larcher, <i>A geometric characterization for a class of discontinuous groups of linear fractional transformations</i>	617
John W. Moon and Leo Moser, <i>Simple paths on polyhedra</i>	629
T. S. Motzkin and Ernst Gabor Straus, <i>Representation of a point of a set as sum of transforms of boundary points</i>	633
Rajakularaman Ponnuswami Pakshirajan, <i>An analogue of Kolmogorov's three-series theorem for abstract random variables</i>	639
Robert Ralph Phelps, <i>Čebyšev subspaces of finite codimension in $C(X)$</i>	647
James Dolan Reid, <i>On subgroups of an Abelian group maximal disjoint from a given subgroup</i>	657
William T. Reid, <i>Riccati matrix differential equations and non-oscillation criteria for associated linear differential systems</i>	665
Georg Johann Rieger, <i>Some theorems on prime ideals in algebraic number fields</i> ...	687
Gene Fuerst Rose and Joseph Silbert Ullian, <i>Approximations of functions on the integers</i>	693
F. J. Sansone, <i>Combinatorial functions and regressive isols</i>	703
Leo Sario, <i>On locally meromorphic functions with single-valued moduli</i>	709
Takayuki Tamura, <i>Semigroups and their subsemigroup lattices</i>	725
Pui-kei Wong, <i>Existence and asymptotic behavior of proper solutions of a class of second-order nonlinear differential equations</i>	737
Fawzi Mohamad Yaqub, <i>Free extensions of Boolean algebras</i>	761