

# Pacific Journal of Mathematics

**ON THE INTEGER SOLUTIONS OF  $y(y + 1) = x(x + 1)(x + 2)$**

LOUIS JOEL MORDELL

# ON THE INTEGER SOLUTIONS OF $y(y + 1) = x(x + 1)(x + 2)$

L. J. MORDELL

This paper contains a solution of the following problem proposed to me by Professor Burton Jones to whom it was given by Mr. Edgar Emerson.

*Problem.* To show that the only integer solutions of

$$(1) \quad y(y + 1) = x(x + 1)(x + 2)$$

are given by

$$(2) \quad x = 0, -1, -2, y = 0, -1; x = 1, y = 2, -3, x = 5, y = 14, -15 .$$

Put

$$2y + 1 = Y, \quad 2x + 2 = X .$$

Then

$$(3) \quad 2Y^2 = X^3 - 4X + 2 .$$

Obviously  $X$  in (3) cannot be odd so it must be shown that the only integer solutions of (3) are given by

$$(4) \quad X = 0, \pm 2, 4, 12 .$$

Diophantine equations of the form

$$(5) \quad Ey^2 = Ax^3 + Bx^2 + Cx + D$$

where  $A, B, C, D, E$  are integers are well known. I proved<sup>1,2</sup> in 1922 that the equation had only a finite number of integer solutions when the right hand side had no squared factor in  $x$ . In fact, this followed immediately from a result<sup>3</sup> I proved in 1913, by quoting Thue's result but which I did not know at that time. Finding these solutions may be a troublesome matter, involving many details, and usually rather difficult or even too difficult, to do.

One method requires a discussion of the field  $R(\theta)$  defined by

$$(6) \quad A\theta^3 + B\theta^2 + C\theta + D \equiv 0 .$$

---

Received November 12, 1962. I should like to thank the referee for his comments on my manuscript. This research was supported by the Air Force Office of Scientific Research.

<sup>1</sup> "Note on the integer solutions of the equation  $Ey^2 = Ax^3 + Bx^2 + Cx + D$ " Messenger of Math., **51** (1922), 169-171.

<sup>2</sup> "On the integer solutions of the equation  $ey^2 = ax^3 + bx^2 + cx + d$ " Proc. London Math. Soc., **21** (1923), 415-419.

<sup>3</sup> "Indeterminate Equations of the third and fourth degrees" Quarterly Journal of Mathematics **45** (1914).

Less arithmetical knowledge may be needed if the equation is reducible in the rational field  $R$ . In general, the field is a cubic field  $R(\theta)$  and then we require a knowledge of

- (1) The expression for the integers of  $R(\theta)$ .
- (2) The number  $h$  of classes of ideals in  $R(\theta)$ .
- (3) The fundamental unit when the equation (6) has only one real solution, and the two fundamental units when all the roots of (6) are real.

Ideal factorization theory may now be applied in (5) on noting that  $x - \theta$  is a factor of the right hand side. The results so found require the solution in integers  $P, Q, R$  of a finite number of pairs of simultaneous quadratic equations

$$(7) \quad F(P, Q, R) = 0, \quad G(P, Q, R) = 1,$$

where  $F$  and  $G$  are homogeneous polynomials of the second degree with rational integer coefficients. In general, it is difficult to find all the integer solutions of (7), but in some simple instances, they can be easily found. The difficulty is increased when there are two fundamental units.

In (3), the field  $R(\theta)$  is defined by

$$(8) \quad f(\theta) = \theta^3 - 4\theta + 2 = 0.$$

The arithmetical details are given in the table on page 112, of Delaunay and Faddeev's book (in Russian) on "The theory of the irrationalities of the third degree".

Thus

- (1) the integers are given by  $a + b\theta + c\theta^2$  where  $a, b, c$  are rational integers,
- (2) the class number  $h = 1$  and so unique factorization exists in  $R(\theta)$ ,
- (3) there are two fundamental units

$$\varepsilon = \theta - 1, \quad \text{and} \quad \eta = 2\theta - 1.$$

All the units are given by  $\pm \varepsilon^l \eta^m$  where  $l, m$ , run through all integer values positive, negative and zero.

We require the factorization of 2. Since  $f(x) \equiv x^3 \pmod{2}$ , the general theory shows that 2 is a perfect cube except for unit factors, and since

$$(9) \quad 2 = \theta(2 + \theta)(2 - \theta),$$

we would expect  $2 \pm \theta$  to be associated with  $\theta$ , namely that  $\varphi = 2/\theta \pm 1$  is an integral unit. In fact,

$$\frac{8}{(\varphi \mp 1)^3} - \frac{8}{\varphi \mp 1} + 2 = 0,$$

or

$$(\varphi \mp 1)^3 - 4(\varphi \mp 1)^2 + 4 = 0 ;$$

and this proves the result about  $\varphi$ . Hence  $2 = \zeta\theta^3$  where  $\zeta$  is a unit, and it is easily seen from this that  $\theta$  is a prime in  $R(\theta)$ .

Write (3) in the form

$$(10) \quad (X - \theta)(X^2 + \theta X + \theta^2 - 4) = \zeta\theta^3 Y^2 .$$

We now show that  $\theta$  is a common divisor of the two left hand factors of (10). Since  $X \equiv 0 \pmod{2}$ , and so  $X \equiv 0 \pmod{\theta^3}$ ,  $\theta$  divides  $X - \theta$  and  $X^2 + \theta X + \theta^2 - 4$  but  $\theta^2$  does not divide  $X - \theta$ . We show that  $4\theta - 3$  is a possible common divisor and note that  $4\theta - 3$  is a prime since its norm is given by  $N(4\theta - 3) = -37$ . Any common divisor must also divide

$$\theta^3 + \theta^2 + \theta^2 - 4 = 3\theta^2 - 4 = (3\theta^3 - 4\theta)/\theta = (8\theta - 6)/\theta = 2(4\theta - 3)/\theta ,$$

and so  $4\theta - 3$  may occur.

We have two cases according as  $X - \theta$  is not or is divisible by  $4\theta - 3$ .

The second case reduces very simply to the first case. Let  $(4\theta - 3)^n$  be the highest power of  $4\theta - 3$  dividing  $X - \theta$ . Then since unique factorization exists in the cubic field  $R(\theta)$ ,

$$X - \theta = \pm\theta(4\theta - 3)^n \varepsilon^l \eta^m (a + b\theta + c\theta^2)^2 ,$$

where  $l, m$  are any integers positive, negative or zero. Take the norm of both sides. Then

$$X^3 - 4X + 2 = \pm 2(37)^n z^2 ,$$

say. Put  $X = 2X_1$ , and then

$$4X_1^3 - 4X_1 + 1 = \pm(37)^n z^2 .$$

Since the left hand side is  $\equiv 1 \pmod{8}$ , we have an impossible congruence  $\pmod{8}$  if  $n$  is odd. If  $n$  is even, we can absorb  $(4\theta - 3)^n$  in  $(a + b\theta + c\theta^2)^2$ . We are then led to an equation of the same form as in the first case, that is,  $X - \theta$  is prime to  $4\theta - 3$ , and is divisible by  $\theta$  but not by  $\theta^2$ .

From (10), we have in the first case

$$(11) \quad X - \theta = \pm\theta\varepsilon^l \eta^m (a + b\theta + c\theta^2)^2 .$$

Clearly it suffices to consider only the four cases

$$(l, m) = (0, 0), (1, 0), (0, 1), (1, 1)$$

on absorbing even powers of  $\varepsilon, \eta$  in the square term. On using (8), we can write

$$(12) \quad (a + b\theta + c\theta^2)^2 = a^2 - 4bc + \theta(2ab + 8bc - 2c^2) \\ + \theta^2(b^2 + 2ac + 4c^2).$$

Take first  $(l, m) = (0, 0)$ . Then using (8) and equating terms in  $\theta, \theta^2$  on both sides of (11), we have

$$(13) \quad 0 = 2ab + 8bc - 2c^2, \\ \mp 1 = a^2 - 4bc + 4(b^2 + 2ac + 4c^2).$$

A congruence mod 4, shows that the plus sign should be taken. Write (13) as

$$b(a + 4c) = c^2, (a + 4c)^2 + 4b^2 - 4bc = 1.$$

When  $b = 0$ , this gives  $c = 0, a = \pm 1$ .

If  $b \neq 0$ ,

$$c^2/b^2 + 4b^2 - 4bc = 1.$$

The minimum value of the left hand side is  $b^2$  given when  $c = b$ , and so  $b = \pm 1, c = \pm 1, a = \mp 3$ .

Since from (11), we have

$$-X = -2(b^2 + 2ac + 4c^2),$$

we have

$$X = 0, -2.$$

Take next  $l = 0, m = 1$ . On multiplying (11) by  $\theta$ , absorbing  $\theta^2$  in the squared term, and with a slight change of notation, we have

$$\pm(\theta X - \theta^2) = (1 - 2\theta)(a^2 - 4bc + \theta(2ab + 8bc - 2c^2) \\ + \theta^2(2ac + b^2 + 4c^2)),$$

and so

$$(14) \quad 0 = a^2 - 4bc + 4(2ac + b^2 + 4c^2), \\ \mp 1 = 2ac + b^2 + 4c^2 - 2(2ab + 8bc - 2c^2), \\ \mp X = 2ab + 8bc - 2c^2 - 2(a^2 - 4bc) - 8(2ac + b^2 + 4c^2).$$

Clearly  $a$  is even and so  $\mp 1$  must be  $+1$ . Then

$$1 = a(2c - 4b) + b^2 - 16bc + 8c^2, \\ 0 = (a + 4c)^2 + 4b(b - c).$$

Suppose first that  $c = 2b$  and so  $b^2 = 1$ . Since solutions  $(a, b, c)$ ,  $(-a, -b, -c)$  give the same value for  $X$ , we need only take  $b = 1, c = 2$ , and then  $(a + 8)^2 - 4 = 0$  and so  $a = -6, -10$ . Then  $b = 1,$

$c = 2, a = -6$  gives  $X = 4$ .

Also  $b = 1, c = 2, a = -10$  gives from (14),  $X_1 = 12$ . Suppose next that  $c \neq 2b$ . Then

$$a = \frac{b^2 - 16bc + 8c^2 - 1}{4b - 2c}, a + 4c = \frac{b^2 - 1}{4b - 2c};$$

and

$$\left(\frac{b^2 - 1}{4b - 2c}\right)^2 + 4b(b - c) = 0.$$

Since  $b \mid (b^2 - 1)$ ,  $b = \pm 1, c = \pm 1, a = \mp 4$ . Then from (14),  $X = 2$ . All the solutions (4) have now been obtained.

We take next  $l = 1, m = 0$  and so

$$\begin{aligned} \pm(X - \theta) &= (\theta - \theta^2)(a^2 - 4bc + \theta(2ab + 8bc - 2c^2) \\ &\quad + \theta^2(2ac + b^2 + 4c^2)). \end{aligned}$$

Then

$$\begin{aligned} 0 &= 2ab + 8bc - 2c^2 - a^2 + 4bc - 4(2ac + b^2 + 4c^2), \\ \mp 1 &= a^2 - 4bc + 6(2ac + b^2 + 4c^2) - 4(2ab + 8bc - 2c^2). \end{aligned}$$

The first equation shows that  $a$  is even and the second that  $a$  is odd; and so no solutions arise.

Suppose finally that  $l = m = 1$ , so that (11) can be written as say,

$$\pm(X - \theta) = \theta(1 - \theta)(1 - 2\theta)F^2.$$

On multiplying by  $\theta$  and absorbing  $\theta^2$  in  $F^2$ , we can write this as, say

$$\begin{aligned} \pm(X\theta - \theta^2) &= (1 - 3\theta + 2\theta^2)(a^2 - 4bc + \theta(2ab + 8bc - 2c^2) \\ &\quad + \theta^2(2ac + b^2 + 4c^2)). \end{aligned}$$

Hence

$$\begin{aligned} 0 &= a^2 - 4bc + 6(2ac + b^2 + 4c^2) - 4(2ab + 8bc - 2c^2). \\ \mp 1 &= 9(2ac + b^2 + 4c^2) - 3(2ab + 8bc - 2c^2) + 2(a^2 - 4bc). \end{aligned}$$

The first equation shows that  $a$  is even and then that  $b$  is even since  $6b^2 \equiv 0 \pmod{4}$ . The second equation shows that  $b$  is odd. Hence no solutions arise.

This finishes the proof.



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RALPH S. PHILLIPS

Stanford University  
Stanford, California

M. G. ARSOVE

University of Washington  
Seattle 5, Washington

J. DUGUNDJI

University of Southern California  
Los Angeles 7, California

LOWELL J. PAIGE

University of California  
Los Angeles 24, California

## ASSOCIATE EDITORS

E. F. BECKENBACH  
T. M. CHERRY

D. DERRY  
M. OHTSUKA

H. L. ROYDEN  
E. SPANIER

E. G. STRAUS  
F. WOLF

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CALIFORNIA RESEARCH CORPORATION  
SPACE TECHNOLOGY LABORATORIES  
NAVAL ORDNANCE TEST STATION

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.



# Pacific Journal of Mathematics

Vol. 13, No. 4

June, 1963

Dallas O. Banks, <i>Bounds for eigenvalues and generalized convexity</i> . . . . .	1031
Jerrold William Bebernes, <i>A subfunction approach to a boundary value problem for ordinary differential equations</i> . . . . .	1053
Woodrow Wilson Bledsoe and A. P. Morse, <i>A topological measure construction</i> . . . . .	1067
George Clements, <i>Entropies of several sets of real valued functions</i> . . . . .	1085
Sandra Barkdull Cleveland, <i>Homomorphisms of non-commutative *-algebras</i> . . . . .	1097
William John Andrew Culmer and William Ashton Harris, <i>Convergent solutions of ordinary linear homogeneous difference equations</i> . . . . .	1111
Ralph DeMarr, <i>Common fixed points for commuting contraction mappings</i> . . . . .	1139
James Robert Dorroh, <i>Integral equations in normed abelian groups</i> . . . . .	1143
Adriano Mario Garsia, <i>Entropy and singularity of infinite convolutions</i> . . . . .	1159
J. J. Gergen, Francis G. Dressel and Wilbur Hallan Purcell, Jr., <i>Convergence of extended Bernstein polynomials in the complex plane</i> . . . . .	1171
Irving Leonard Glicksberg, <i>A remark on analyticity of function algebras</i> . . . . .	1181
Charles John August Halberg, Jr., <i>Semigroups of matrices defining linked operators with different spectra</i> . . . . .	1187
Philip Hartman and Nelson Onuchic, <i>On the asymptotic integration of ordinary differential equations</i> . . . . .	1193
Isidore Heller, <i>On a class of equivalent systems of linear inequalities</i> . . . . .	1209
Joseph Hersch, <i>The method of interior parallels applied to polygonal or multiply connected membranes</i> . . . . .	1229
Hans F. Weinberger, <i>An effectless cutting of a vibrating membrane</i> . . . . .	1239
Melvin F. Janowitz, <i>Quantifiers and orthomodular lattices</i> . . . . .	1241
Samuel Karlin and Albert Boris J. Novikoff, <i>Generalized convex inequalities</i> . . . . .	1251
Tilla Weinstein, <i>Another conformal structure on immersed surfaces of negative curvature</i> . . . . .	1281
Gregers Louis Krabbe, <i>Spectral permanence of scalar operators</i> . . . . .	1289
Shige Toshi Kuroda, <i>Finite-dimensional perturbation and a representation of scattering operator</i> . . . . .	1305
Marvin David Marcus and Afton Herbert Cayford, <i>Equality in certain inequalities</i> . . . . .	1319
Joseph Martin, <i>A note on uncountably many disks</i> . . . . .	1331
Eugene Kay McLachlan, <i>Extremal elements of the convex cone of semi-norms</i> . . . . .	1335
John W. Moon, <i>An extension of Landau's theorem on tournaments</i> . . . . .	1343
Louis Joel Mordell, <i>On the integer solutions of <math>y(y + 1) = x(x + 1)(x + 2)</math></i> . . . . .	1347
Kenneth Roy Mount, <i>Some remarks on Fitting's invariants</i> . . . . .	1353
Miroslav Novotný, <i>Über Abbildungen von Mengen</i> . . . . .	1359
Robert Dean Ryan, <i>Conjugate functions in Orlicz spaces</i> . . . . .	1371
John Vincent Ryff, <i>On the representation of doubly stochastic operators</i> . . . . .	1379
Donald Ray Sherbert, <i>Banach algebras of Lipschitz functions</i> . . . . .	1387
James McLean Sloss, <i>Reflection of biharmonic functions across analytic boundary conditions with examples</i> . . . . .	1401
L. Bruce Treybig, <i>Concerning homogeneity in totally ordered, connected topological space</i> . . . . .	1417
John Wermer, <i>The space of real parts of a function algebra</i> . . . . .	1423
James Juei-Chin Yeh, <i>Orthogonal developments of functionals and related theorems in the Wiener space of functions of two variables</i> . . . . .	1427
William P. Ziemer, <i>On the compactness of integral classes</i> . . . . .	1437