

# Pacific Journal of Mathematics

**A JORDAN-HÖLDER THEOREM**

CHARLES EDWARD WATTS

# A JORDAN-HÖLDER THEOREM

CHARLES E. WATTS

1. The purpose of this note is to present a certain general theorem, of the Jordan-Holder type, for finite groups. This theorem, although a simple and natural extension of the classical theorem, has we believe passed unnoticed before. The technique of proof is foreign to the usual methods of finite group theory, but seems well-suited to the situation.

2. A nonempty class  $\mathcal{D}$  of finite groups will be called a *genetic class* provided:

(1) If  $G_1$  belongs to  $\mathcal{D}$  and if  $G_2$  is isomorphic to  $G_1$ , then  $G_2$  belongs to  $\mathcal{D}$ .

(2) If  $G$  belongs to  $\mathcal{D}$ , then every normal subgroup and every quotient group of  $G$  also belongs to  $\mathcal{D}$ .

The following examples of genetic classes will be used as illustrations in the sequel:

The class  $\mathcal{G}$  of all finite groups.

The class  $\mathcal{E}$  of all one-element groups.

The class  $\mathcal{A}$  of all finite abelian groups.

The class  $\mathcal{O}$  of all groups of odd order.

The class  $\mathcal{G}_n$  of all groups of order  $\leq n$ .

Given any genetic class  $\mathcal{D}$ , we shall construct a "Grothendieck group" in the following way. Let  $\Sigma$  be the (countable) set of all isomorphism classes of finite groups, and let  $F$  be the free abelian group generated by  $\Sigma$ . If  $G$  is any finite group, its isomorphism class will be denoted by  $[G]$ , so that elements of  $F$  are finite sums

$$\sum \lambda_i [G_i], \quad \lambda_i \in \mathbb{Z},$$

where  $\mathbb{Z}$  denotes the ring of integers. We let  $N(\mathcal{D})$  be the subgroup of  $F$  generated by all elements of the form

$$[G] - [H] - [G/H]$$

such that  $H$  is a normal subgroup of  $G$  and  $G/H$  belongs to the genetic class  $\mathcal{D}$ . Finally we set  $K(\mathcal{D}) = F/N(\mathcal{D})$  and let  $k: F \rightarrow K(\mathcal{D})$  be the natural epimorphism. Our object is to determine the structure of the abelian group  $K(\mathcal{D})$ .

3. Let  $\mathcal{D}$  be a genetic class and let  $G$  be an arbitrary finite

---

Received April 3, 1963. This work was supported by the National Science Foundation under grant NSF-G 24155.

group. We say  $G$  is  $\mathcal{D}$ -simple provided that  $G$  has more than one element and that no proper quotient group of  $G$  belongs to  $\mathcal{D}$ , i.e., if  $H$  is normal in  $G$  and if  $G/H$  belongs to  $\mathcal{D}$ , then  $H = G$  or  $H = 1$ . In particular, if  $G$  itself belongs to  $\mathcal{D}$  and is  $\mathcal{D}$ -simple, then it is simple because of the second axiom for genetic classes. The following illustrations are based on the examples given in § 2 above.

$G$  is  $\mathcal{S}$ -simple if and only if it is simple in the classical sense.

Every finite group is  $\mathcal{S}$ -simple.

$G$  is  $\mathcal{A}$ -simple if and only if either  $G$  is cyclic of prime order, or else  $\neq 1$  and equal to its commutator subgroup.

$G$  is  $\mathcal{O}$ -simple if and only if  $G$  is simple and has odd order, or else has even order and no proper normal subgroups of odd index.

$G$  is  $\mathcal{E}_n$ -simple if and only if  $G$  is simple and has order  $\leq n$  or else has order  $> n$  and no proper normal subgroups of index  $\leq n$ .

Having given the definition of  $\mathcal{D}$ -simplicity, we can now state the theorem referred to in § 1:

**THEOREM.** *Given any genetic class  $\mathcal{D}$ ,  $K(\mathcal{D})$  is the free abelian group freely generated by the elements  $k[S]$ , where  $S$  is  $\mathcal{D}$ -simple.*

4. In this section we begin to prove the theorem above and show its relation to the Jordan-Holder theorem.

Let  $\mathcal{D}$  be a genetic class,  $G$  any finite group. If  $G$  is not  $\mathcal{D}$ -simple and if  $G \neq 1$ , we can find a normal subgroup  $G'_1$  of  $G$  such that  $1 \neq G'_1 \neq G$  and  $G/G'_1$  belongs to  $\mathcal{D}$ . Let  $G_1$  be a maximal proper normal subgroup containing  $G'_1$ . Then  $G/G_1$ , being a quotient of  $G/G'_1$ , is in  $\mathcal{D}$  and is simple (and a fortiori  $\mathcal{D}$ -simple). Now if  $G_1$  is not  $\mathcal{D}$ -simple and is  $\neq 1$ , we repeat the process to find a normal subgroup  $G_2$  of  $G_1$  such that  $G_1/G_2$  is in  $\mathcal{D}$  and is simple. Eventually we shall get a sequence

$$(1) \quad G = G_0, G_1, G_2, \dots, G_n,$$

where  $G_{i+1}$  is normal in  $G_i$ , where  $G_i/G_{i+1}$  is in  $\mathcal{D}$  and is simple ( $i = 0, 1, \dots, n-1$ ), and where either  $G_n$  is  $\mathcal{D}$ -simple and not in  $\mathcal{D}$  or else  $G_n = 1$ . Since  $G_i/G_{i+1}$  belongs to  $\mathcal{D}$ , we have

$$\begin{aligned} k[G] &= k[G_0] = k[G_0/G_1] + k[G_1] = \dots \\ &= k[G_0/G_1] + k[G_1/G_2] + \dots + k[G_{n-1}/G_n] + k[G_n]. \end{aligned}$$

Clearly if  $G_n = 1$ , then  $k[G_n] = 0$ . Thus we have shown that the elements  $k[S]$ ,  $S$   $\mathcal{D}$ -simple, generate the group  $K(\mathcal{D})$ .

We remark at this point that once we have shown the linear independence (over  $Z$ ) of these generators, it will follow that the  $\mathcal{D}$ -simple groups  $G_i/G_{i+1}$ ,  $G_n$  are uniquely determined (up to isomorphism) by  $G$ , and are independent of the sequence 1) used to compute

them. Thus in the case  $\mathcal{D} = \mathcal{G}$ , we get precisely the classical Jordan-Holder theorem. In the general case, the groups  $G_i/G_{i+1}$  are of course among the composition factors of  $G$ , but the group  $G_n$  (if it is not 1) is something new. It is a subnormal subgroup of  $G$  which depends, up to isomorphism, only on  $G$  and on  $\mathcal{D}$ .

Continuing our digression from the proof, let us say that two finite groups  $G$  and  $G'$  are  $\mathcal{D}$ -equivalent if they represent the same element of  $K(\mathcal{D})$ . Thus  $G$  and  $G'$  are  $\mathcal{G}$ -equivalent if and only if they have the same composition factors, while to be  $\mathcal{G}$ -equivalent it is clear that they must be isomorphic. In general, the smaller the genetic class  $\mathcal{D}$ , the sharper is the notion of  $\mathcal{D}$ -equivalence.

5. We return to the proof of the theorem; it remains to show that the generators  $k[S]$ ,  $S$   $\mathcal{D}$ -simple, are linearly independent over  $Z$ . We shall show that for each  $\mathcal{D}$ -simple group  $S$  there exists an integer-valued function  $f$  defined on  $\Sigma$  (and depending on  $S$ ) such that:

- (1)  $f[S] = 1$ ;
- (2)  $f[T] = 0$  if  $T$  is any  $\mathcal{D}$ -simple group not isomorphic to  $S$ ;
- (3) If  $H$  is a normal subgroup of  $G$  and if  $G/H$  is in  $\mathcal{D}$ , then  $f[G] = f[H] + f[G/H]$ .

Because of (3) such a function induces a homomorphism  $K(\mathcal{D}) \rightarrow Z$ , vanishing on  $k[T]$  if  $T$  is as in (2), but equal to 1 on  $k[S]$ . The linear independence of the generators is an immediate consequence of the existence of such homomorphisms.

We construct  $f$  inductively. Let  $\Sigma_r$  be the set of isomorphism classes of groups of order  $\leq r$ . Define  $f = 0$  on  $\Sigma_1$ . Now suppose that  $f$  has been defined on  $\Sigma_r$  in such a way that (1), (2), (3) hold whenever  $S, T, G$  have orders  $\leq r$ . Next suppose that  $G$  has order  $r + 1$ . If  $G$  is  $\mathcal{D}$ -simple, then the value of  $f[G]$  is forced by (1) or by (2). If  $G$  is not  $\mathcal{D}$ -simple, then it has a normal subgroup  $H$  with  $G/H$  in  $\mathcal{D}$  and with  $H$  and  $G/H$  in  $\Sigma_r$ . Consequently, the value of  $f[G]$  must be given by (3), and it remains to show that  $f[H] + f[G/H]$  is independent of the choice of  $H$  as long as  $H$  has order  $\leq r$  and  $G/H$  is in  $\mathcal{D}$ .

Thus let  $K$  be another such subgroup. Then  $G/HK$  is in  $\mathcal{D}$ , since it is a quotient of  $G/H$ , and  $H/H \cap K$  is in  $\mathcal{D}$ , since it is isomorphic to  $HK/H$ , which is normal in  $G/H$ . Hence using the Noether isomorphisms we get

$$\begin{aligned} f[H] + f[G/H] &= f[H] + f[G/HK] + f[HK/H] \\ &= f[H] + f[G/HK] + f[K/H \cap K] \\ &= f[H/H \cap K] + f[H \cap K] + f[G/HK] + f[K/H \cap K]. \end{aligned}$$

This last expression being symmetric in  $H$  and  $K$ , it follows that  $f[H] + f[G/H] = f[K] + f[G/K]$ . Thus we have shown how to extend  $f$  unambiguously to  $\Sigma_{r+1}$  in such a way that (1), (2), (3) still hold on this enlarged domain. Therefore  $f$  can be defined on all of  $\Sigma$  so as to have the desired properties, and this completes the proof.

UNIVERSITY OF ROCHESTER

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

ROBERT OSSERMAN  
Stanford University  
Stanford, California

J. DUGUNDJI  
University of Southern California  
Los Angeles 7, California

M. G. ARSOVE  
University of Washington  
Seattle 5, Washington

LOWELL J. PAIGE  
University of California  
Los Angeles 24, California

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CALIFORNIA RESEARCH CORPORATION  
SPACE TECHNOLOGY LABORATORIES  
NAVAL ORDNANCE TEST STATION

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and on submission, must be accompanied by a separate author's résumé. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Tom M. (Mike) Apostol and Herbert S. Zuckerman, <i>On the functional equation</i> $F(mn)F((m, n)) = F(m)F(n)f((m, n))$ .....	377
Reinhold Baer, <i>Irreducible groups of automorphisms of abelian groups</i> .....	385
Herbert Stanley Bear, Jr., <i>An abstract potential theory with continuous kernel</i> .....	407
E. F. Beckenbach, <i>Superadditivity inequalities</i> .....	421
R. H. Bing, <i>The simple connectivity of the sum of two disks</i> .....	439
Herbert Busemann, <i>Length-preserving maps</i> .....	457
Heron S. Collins, <i>Characterizations of convolution semigroups of measures</i> .....	479
Paul F. Conrad, <i>The relationship between the radical of a lattice-ordered group and complete distributivity</i> .....	493
P. H. Doyle, III, <i>A sufficient condition that an arc in <math>S^n</math> be cellular</i> .....	501
Carl Clifton Faith and Yuzo Utumi, <i>Intrinsic extensions of rings</i> .....	505
Watson Bryan Fulks, <i>An approximate Gauss mean value theorem</i> .....	513
Arshag Berge Hajian, <i>Strongly recurrent transformations</i> .....	517
Morisuke Hasumi and T. P. Srinivasan, <i>Doubly invariant subspaces. II</i> .....	525
Lowell A. Hinrichs, Ivan Niven and Charles L. Vanden Eynden, <i>Fields defined by polynomials</i> .....	537
Walter Ball Laffer, I and Henry B. Mann, <i>Decomposition of sets of group elements</i> .....	547
John Albert Lindberg, Jr., <i>Algebraic extensions of commutative Banach algebras</i> .....	559
W. Ljunggren, <i>On the Diophantine equation <math>Cx^2 + D = y^n</math></i> .....	585
M. Donald MacLaren, <i>Atomic orthocomplemented lattices</i> .....	597
Moshe Marcus, <i>Transformations of domains in the plane and applications in the theory of functions</i> .....	613
Philip Miles, <i><math>B^*</math> algebra unit ball extremal points</i> .....	627
W. F. News, <i>On the difference and sum of a basic set of polynomials</i> .....	639
Barbara Osofsky, <i>Rings all of whose finitely generated modules are injective</i> .....	645
Calvin R. Putnam, <i>Toeplitz matrices and invertibility of Hankel matrices</i> .....	651
Shoichiro Sakai, <i>Weakly compact operators on operator algebras</i> .....	659
James E. Simpson, <i>Nilpotency and spectral operators</i> .....	665
Walter Laws Smith, <i>On the elementary renewal theorem for non-identically distributed variables</i> .....	673
T. P. Srinivasan, <i>Doubly invariant subspaces</i> .....	701
J. Roger Teller, <i>On the extensions of lattice-ordered groups</i> .....	709
Robert Charles Thompson, <i>Unimodular group matrices with rational integers as elements</i> .....	719
J. L. Walsh and Ambikeshwar Sharma, <i>Least squares and interpolation in roots of unity</i> .....	727
Charles Edward Watts, <i>A Jordan-Hölder theorem</i> .....	731
Kung-Wei Yang, <i>On some finite groups and their cohomology</i> .....	735
Adil Mohamed Yaqub, <i>On the ring-logic character of certain rings</i> .....	741
Paul Ruel Young, <i>A note on pseudo-creative sets and cylinders</i> .....	749