

Pacific Journal of Mathematics

RINGS OF ARITHMETIC FUNCTIONS

L. CARLITZ

RINGS OF ARITHMETIC FUNCTIONS

L. CARLITZ

1. **Introduction.** Let F denote a fixed but arbitrary field and let Z denote the set of positive integers. By an *arithmetic function* f is meant a function from Z to F , that is to say $f(n) \in F$ for all $n \in Z$. If f, g are two arithmetic functions, the sum $h = f + g$ is defined by means of

$$(1) \quad h(n) = f(n) + g(n) \quad (n \in Z).$$

There are two products that are of interest, the *ordinary* product defined by

$$(2) \quad h(n) = f(n)g(n) \quad (n \in Z),$$

and the Dirichlet product defined by

$$(3) \quad h(n) = \sum_{rs=n} f(r)g(s) \quad (n \in Z),$$

where the summation on the right is extended over all factorizations $rs = n$. We shall denote the ordinary product by $f \circ g$ and the Dirichlet product by $f * g$.

Let S denote the set of arithmetic functions as defined above. It is well known and easy to prove that the system

$$(4) \quad \Omega = (S, f, \circ)$$

is a commutative ring. The multiplicative identity of Ω is defined by

$$(5) \quad v(n) = 1 \quad (n \in Z).$$

Clearly Ω is not a domain of integrity; note however that there are no nilpotent elements in Ω . On the other hand the system

$$(6) \quad \mathcal{A} = (S, f, *)$$

is a domain of integrity. The multiplicative identity of \mathcal{A} is given by

$$(7) \quad u(n) = \begin{cases} 1 & (n = 1) \\ 0 & (n > 1). \end{cases}$$

Moreover the function f has an inverse (relative to $*$) if and only if

$$(8) \quad f(1) \neq 0;$$

Received September 15, 1963, and in revised form November 12, 1963. Supported in part by NSF grant G16485.

the set of functions that satisfy (8) evidently constitute an abelian group with respect to $*$.

If $\lambda \in F$ we define the function λf by means of

$$(9) \quad (\lambda f)(n) = \lambda \cdot f(n) \quad (n \in Z) .$$

It follows at once that S is a vector space over F of infinite dimension. Also we have

$$\lambda(f \circ g) = (\lambda f) \circ g = f \circ (\lambda g) , \quad \lambda(f * g) = (\lambda f) * g = f * (\lambda g) .$$

If in place of Z we employ a semigroup J that has no units except the identity, a countable infinity of primes, and which has the unique factorization property, the resulting systems Ω and Δ are not essentially different. Indeed if $\bar{p}_1, \bar{p}_2, \bar{p}_3, \dots$ denote the primes of J we may set up the correspondence $f \leftrightarrow \bar{f}$ by means of $f(n) = \bar{f}(\bar{n})$, where

$$(10) \quad n = \prod p_j^{e_j} , \quad \bar{n} = \prod \bar{p}_j^{e_j} ,$$

where the first half of (10) is the usual factorization of n into primes. There is therefore little loss in generality in restricting the discussion to Z .

In view of the above it is of interest to consider the system

$$(11) \quad \Phi = (S, +, \circ, *)$$

with three binary operations and in particular to attempt to give an abstract formulation of such systems. Since \circ and $*$ do not combine in any very obvious way, it is perhaps not clear how this can be done. We shall obtain such a characterization by making use of *minimal* functions. A function f is minimal provided there exists an integer k (depending on f) such that

$$(12) \quad f(n) = 0 \quad (n \neq k) ; \quad f(k) \neq 0 .$$

We remark that Cashwell and Everett [1] have proved that Δ is a unique factorization domain. However this result will not be required in what follows.

2. As above let F denote a fixed but arbitrary field. Let \bar{S} denote a vector space over F . The elements of \bar{S} will be denoted by small italic letters, the elements of F by small Greek letters; addition in \bar{S} will be denoted by $+$. Moreover we have two "multiplications" denoted by \circ and $*$. The following assumptions will be made.

S1. The system

$$(13) \quad \Omega = (\bar{S}, +, \circ)$$

is a commutative ring with multiplicative identity \bar{v} . Moreover

$$\alpha(\bar{f} \circ \bar{g}) = (\alpha\bar{f}) \circ \bar{g} = \bar{f} \circ (\alpha\bar{g}) \quad (\bar{f}, \bar{g} \in \bar{S}, \alpha \in F).$$

S2. The system

$$(14) \quad \bar{A} = (\bar{S}, +, *)$$

is a domain of integrity with multiplicative identity \bar{u} . Moreover

$$\alpha(\bar{f} * \bar{g}) = (\alpha\bar{f}) * \bar{g} = \bar{f} * (\alpha\bar{g}) \quad (\bar{f}, \bar{g} \in \bar{S}, \alpha \in F).$$

DEFINITION. Two elements $\bar{f}, \bar{g} \in \bar{S}$ are *associates* provided $\bar{f} = \lambda\bar{g}$, where $\lambda \in F$, $\lambda \neq 0$.

DEFINITION. An element $\bar{f} \in \bar{S}$, $\bar{f} \neq 0$, is *minimal* provided

$$(15) \quad \bar{f} \circ \bar{g} = \lambda(\bar{f}, \bar{g})\bar{f} \quad (\bar{g} \in \bar{S})$$

where \bar{g} is any element of \bar{S} and $\lambda(\bar{f}, \bar{g})$ is a number of F . It is evident that $\lambda(\bar{f}, \bar{g})$ is unique.

Clearly the associate of a minimal element is also minimal. Also it is evident that if \bar{f}, \bar{g} are two minimal elements that are not associates then

$$(16) \quad \bar{f} \circ \bar{g} = 0.$$

S3. For each minimal element \bar{f} there exists a nonzero number $\lambda(\bar{f})$ of F such that

$$(17) \quad \bar{f} \circ \bar{f} = \lambda(\bar{f})\bar{f}.$$

DEFINITION. A minimal element $\bar{f} \in \bar{S}$ is *normalized* provided

$$(18) \quad \bar{f} \circ \bar{f} = \bar{f}.$$

S4. If \bar{g} is an arbitrary nonzero element of \bar{S} there exists at least one minimal element \bar{f} such that $\lambda(\bar{f}, \bar{g}) \neq 0$, where $\lambda(\bar{f}, \bar{g})$ is defined by (15).

Let M denote the set of normalized minimal elements.

S5. M is a semigroup with respect to $*$; the identity element of M coincides with \bar{u} , the multiplicative identity of \bar{A} . Moreover M contains no units except the identity.

DEFINITION. An element \bar{f} of M , $\bar{f} \neq \bar{u}$, is *prime* provided $\bar{f} = \bar{g} * \bar{h}$ implies $\bar{g} = \bar{u}$ or $\bar{h} = \bar{u}$.

S6. M contains a countable number of primes. Any element of M , different from \bar{u} , can be expressed as a product of primes in essentially only one way.

DEFINITION. Let $\bar{f}_1, \bar{f}_2, \bar{f}_3, \dots$ denote the elements of M . If \bar{g} is an arbitrary element of \bar{S} the numbers

$$\lambda_j(\bar{g}) = \lambda(\bar{f}_j, \bar{g})$$

defined by

$$(19) \quad \bar{f}_j \circ \bar{g} = \lambda(\bar{f}_j, \bar{g})\bar{f}_j$$

may be called the (Dirichlet) coefficients of \bar{g} .

S7. If $\bar{g} \neq \bar{h}$ then for at least one value of j we have $\lambda_j(\bar{g}) \neq \lambda_j(\bar{h})$.

It evidently follows that two elements of \bar{S} are equal if and only if the respective sets of coefficients are equal.

S8. If \bar{g} and \bar{h} are arbitrary elements of \bar{S} while \bar{f} is an element of M , then

$$\bar{f} \circ (\bar{g} * \bar{h}) = \Sigma(\bar{f}_r \circ \bar{g}) * (\bar{f}_s \circ \bar{h})$$

where the summation is over all $\bar{f}_r, \bar{f}_s \in M$ such that $\bar{f}_r * \bar{f}_s = \bar{f}$.

Finally we have

S9. For every sequence $\lambda_1, \lambda_2, \lambda_3, \dots, \lambda_j \in F$, there exists a $\bar{g} \in \bar{S}$ such that

$$\bar{f}_j \circ \bar{g} = \lambda_j \bar{f}_j \quad (j = 1, 2, 3, \dots).$$

3. LEMMA 1. If \bar{f}_i, \bar{f}_j are distinct elements of M then

$$(20) \quad \bar{f}_i \circ \bar{f}_j = 0 \quad (i \neq j).$$

This is immediate from (16).

LEMMA 2. Let \bar{g}, \bar{h} be two arbitrary elements of \bar{S} and let $\lambda_j(\bar{g}), \lambda_j(\bar{h})$ denote the respective sets of coefficients of \bar{g} and \bar{h} . Then

$$(21) \quad \lambda_j(\bar{g} \circ \bar{h}) = \lambda_j(\bar{g})\lambda_j(\bar{h}) \quad (j = 1, 2, 3, \dots).$$

Indeed we have by (18) and (19)

$$\begin{aligned} \lambda_j(\bar{g} \circ \bar{h})\bar{f}_j &= \bar{f}_j \circ (\bar{g} \circ \bar{h}) = (\bar{f}_j \circ \bar{g}) \circ (\bar{f}_j \circ \bar{h}) = (\lambda_j(\bar{g})\bar{f}_j) \circ (\lambda_j(\bar{h})\bar{f}_j) \\ &= \lambda_j(\bar{g})\lambda_j(\bar{h})(\bar{f}_j \circ \bar{f}_j) = \lambda_j(\bar{g})\lambda_j(\bar{h})\bar{f}_j \end{aligned}$$

and (21) follows at once.

LEMMA 3. Let \bar{g}, \bar{h} be two arbitrary elements of \bar{S} and let $\lambda_j(\bar{g}), \lambda_j(\bar{h})$ denote the respective sets of coefficients of \bar{g} and \bar{h} . Then

$$(22) \quad \lambda_j(\bar{g} * \bar{h}) = \sum \lambda_r(\bar{g}) \lambda_s(\bar{h}) \quad (j = 1, 2, 3, \dots),$$

where the summation is over all pairs r, s such that

$$(23) \quad \bar{f}_r * \bar{f}_s = \bar{f}_j.$$

Proof. We have by S8

$$\begin{aligned} \lambda_j(\bar{g} * \bar{h}) \bar{f}_j &= \bar{f}_j \circ (\bar{g} * \bar{h}) = \sum_{\bar{f}_r * \bar{f}_s = \bar{f}_j} (\bar{f}_r \circ \bar{g}) * (\bar{f}_s \circ \bar{h}) \\ &= \sum_{\bar{f}_r * \bar{f}_s = \bar{f}_j} (\lambda_r(\bar{g}) \bar{f}_r) * (\lambda_s(\bar{h}) \bar{f}_s) \\ &= \left\{ \sum_{\bar{f}_r * \bar{f}_s = \bar{f}_j} \lambda_r(\bar{g}) \lambda_s(\bar{h}) \right\} \bar{f}_j. \end{aligned}$$

This evidently implies (22).

Let $\bar{p}_1, \bar{p}_2, \bar{p}_3, \dots$ denote the primes of M and let p_1, p_2, p_3, \dots denote the ordinary primes. We assume to begin with that the number of primes in M is infinite and set up the correspondence

$$(24) \quad p_j \leftrightarrow \bar{p}_j \quad (j = 1, 2, 3, \dots).$$

If

$$n = p_1^{e_1} p_2^{e_2} \dots p_r^{e_r}$$

is an arbitrary positive integer, we put

$$(25) \quad \bar{f}_n = \bar{p}_1^{e_1} * \bar{p}_2^{e_2} * \dots * \bar{p}_r^{e_r},$$

where

$$\bar{g}^e = \bar{g} * \dots * \bar{g},$$

with e factors on the right. By means of (25) we have the one-to-one correspondence between Z and M

$$(26) \quad n \leftrightarrow \bar{f}_n \quad (n = 1, 2, 3, \dots).$$

Let \bar{g} be an arbitrary element of \bar{S} and let $\lambda_j(\bar{g})$ denote the set of coefficients of \bar{g} . Corresponding to \bar{g} we have the function g in S defined by

$$(27) \quad g(n) = \lambda_n(\bar{g}).$$

Conversely if g is any function in S then by S9 and S7 the element \bar{g} of \bar{S} is uniquely determined by means of (27), so that we have obtained a one-to-one correspondence between S and \bar{S} .

Now if $\alpha \in F$ it follows at once from (27) that

$$(28) \quad \alpha g(n) = \lambda_n(\alpha \bar{g}) ,$$

so that scalar multiplication is consistent with the correspondence defined by (27). Again if $h \in S$ and $\bar{h} \in \bar{S}$ satisfy

$$(29) \quad h(n) = \lambda_n(\bar{h})$$

it is clear that

$$(30) \quad g(n) + h(n) = \lambda_n(\bar{g} + \bar{h}) .$$

In the next place, if (27) and (29) hold, it follows from Lemma 2 that

$$(31) \quad g(n)h(n) = \lambda_n(\bar{g})\lambda_n(\bar{h}) = \lambda_n(\bar{g} \circ \bar{h}) .$$

Thus if \bar{g} corresponds to g and \bar{h} corresponds to h then $\bar{g} \circ \bar{h}$ corresponds to the "ordinary" product of g and h .

Next we observe that if

$$r \leftrightarrow \bar{f}_r , \quad s \leftrightarrow \bar{f}_s$$

under the correspondence (26), then

$$(32) \quad rs \leftrightarrow \bar{f}_r * \bar{f}_s .$$

Thus, assuming (27) and (29), we get

$$\sum_{rs=n} g(r)h(s) = \sum_{rs=n} \lambda_r(\bar{g})\lambda_s(\bar{h}) = \sum_{\bar{f}_r * \bar{f}_s = \bar{f}_n} \lambda_r(\bar{g})\lambda_s(\bar{h}) .$$

Therefore, by Lemma 3,

$$(33) \quad \sum_{rs=n} g(r)h(s) = \lambda_n(\bar{g} * \bar{h}) .$$

Thus if \bar{g} corresponds to g and \bar{h} corresponds to h then $\bar{g} * \bar{h}$ corresponds to the Dirichlet product of g and h .

Combining (27), (28), (29), (30), (31), (32) and (33) we have the following result.

THEOREM 1. *Let Φ denote the system of arithmetic functions from the integers to an arbitrary but fixed field F as defined in §1. Let $\bar{\Phi}$ be a structure with the three binary operations $+$, \circ , $*$ that satisfies the assumptions S1-S9 of §2. Also let the number of primes in M be infinite. Then $\bar{\Phi}$ is isomorphic to Φ , all operations being preserved under the isomorphism.*

4. We have assumed in the above result that the number of

prime elements in M is infinite. The conclusion of the theorem is no longer valid when the number of primes is finite. However it is easily verified that in this case $\bar{\Phi}$ is isomorphic to a subset of Φ . More precisely, we have the following result.

Let $\bar{p}_1, \bar{p}_2, \dots, \bar{p}_k$ denote the primes of M and let p_1, p_2, \dots, p_k be a set of k distinct primes, for example the first k primes. Then the correspondence (26) holds except that n is now restricted to the set of integers Z_k whose prime divisors are in the set p_1, p_2, \dots, p_k . Consider the set of functions g such that

$$(34) \quad g(n) = 0 \quad (n \in Z - Z_k),$$

while $g(n)$ is an arbitrary number of F when $n \in Z_k$. It is easily verified that the set of functions satisfying (34) is closed under scalar, ordinary and Dirichlet multiplication. We denote the system by Φ_k . Then we have

THEOREM 2. *Let Φ_k denote the system of arithmetic functions that satisfy (34). Let $\bar{\Phi}$ be a structure with three binary operations $+, \circ, *$ that satisfies the assumptions S1–S9 of § 2 but let the number of primes in M equal k . Then $\bar{\Phi}$ is isomorphic to Φ_k .*

It is evident that Φ_k is isomorphic to $F\{x_1, x_2, \dots, x_k\}$, the ring of formal power series in k indeterminates with coefficients in F .

REMARK. The referee has pointed out that S4 and S7 are equivalent, in the presence of the other assumptions. First, S7 implies S4. For $\bar{g} \neq 0$, by S7 there exists a j such that $\lambda_j(\bar{g}) \neq \lambda_j(0) = 0$. Hence S4 holds with $\bar{f} = \bar{f}_j$.

Conversely, S4 implies S7. For if $\bar{g} \neq \bar{h}$, then $\bar{d} = \bar{g} - \bar{h} \neq 0$. By S4 there exists a minimal \bar{f} such that $\bar{f} \circ \bar{d} = \lambda(\bar{f}, \bar{d})\bar{f}$, where $\lambda(\bar{f}, \bar{d}) \neq 0$. Since \bar{f} is minimal, $\bar{f} \circ \bar{f} = \lambda(\bar{f})\bar{f}$, where $\lambda(\bar{f}) \neq 0$ by S3. Hence there exists a minimal

$$\bar{f} = (\lambda(\bar{f}))^{-1}\bar{f}$$

(an associate of the minimal element \bar{f}) which is also normalized. Thus

$$\begin{aligned} \bar{f}_j \circ \bar{d} &= \lambda(\bar{f}, \bar{d})\bar{f}_j = \bar{f}_j \circ (\bar{g} - \bar{h}) = \bar{f}_j \circ \bar{g} - \bar{f}_j \circ \bar{h} \\ &= \lambda_j(\bar{g})\bar{f}_j - \lambda_j(\bar{h})\bar{f}_j = [\lambda_j(\bar{g}) - \lambda_j(\bar{h})]\bar{f}_j. \end{aligned}$$

Hence

$$\lambda_j(\bar{g}) - \lambda_j(\bar{h}) = \lambda(\bar{f}, \bar{d}) \neq 0.$$

REFERENCE

1. E. D. Cashwell and C. J. Everett, *The ring of number-theoretic functions*, Pacific J. Math., **9** (1959), 975–985.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

ROBERT OSSERMAN

Stanford University
Stanford, California

M. G. ARSOVE

University of Washington
Seattle 5, Washington

J. DUGUNDJI

University of Southern California
Los Angeles 7, California

LOWELL J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and on submission, must be accompanied by a separate author's résumé. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 14, No. 4

August, 1964

Homer Franklin Bechtell, Jr., <i>Pseudo-Frattini subgroups</i>	1129
Thomas Kelman Boehme and Andrew Michael Bruckner, <i>Functions with convex means</i>	1137
Lutz Bungart, <i>Boundary kernel functions for domains on complex manifolds</i>	1151
L. Carlitz, <i>Rings of arithmetic functions</i>	1165
D. S. Carter, <i>Uniqueness of a class of steady plane gravity flows</i>	1173
Richard Albert Dean and Robert Harvey Oehmke, <i>Idempotent semigroups with distributive right congruence lattices</i>	1187
Lester Eli Dubins and David Amiel Freedman, <i>Measurable sets of measures</i>	1211
Robert Pertsch Gilbert, <i>On class of elliptic partial differential equations in four variables</i>	1223
Harry Gonshor, <i>On abstract affine near-rings</i>	1237
Edward Everett Grace, <i>Cut points in totally non-semi-locally-connected continua</i>	1241
Edward Everett Grace, <i>On local properties and G_δ sets</i>	1245
Keith A. Hardie, <i>A proof of the Nakaoka-Toda formula</i>	1249
Lowell A. Hinrichs, <i>Open ideals in $C(X)$</i>	1255
John Rolfe Isbell, <i>Natural sums and abelianizing</i>	1265
G. W. Kimble, <i>A characterization of extremals for general multiple integral problems</i>	1283
Nand Kishore, <i>A representation of the Bernoulli number B_n</i>	1297
Melven Robert Krom, <i>A decision procedure for a class of formulas of first order predicate calculus</i>	1305
Peter A. Lappan, <i>Identity and uniqueness theorems for automorphic functions</i>	1321
Lorraine Doris Lavalley, <i>Mosaics of metric continua and of quasi-Peano spaces</i>	1327
Mark Mahowald, <i>On the normal bundle of a manifold</i>	1335
J. D. McKnight, <i>Kleene quotient theorems</i>	1343
Charles Kimbrough Megibben, III, <i>On high subgroups</i>	1353
Philip Miles, <i>Derivations on B^* algebras</i>	1359
J. Marshall Osborn, <i>A generalization of power-associativity</i>	1367
Theodore G. Ostrom, <i>Nets with critical deficiency</i>	1381
Elvira Rapaport Strasser, <i>On the defining relations of a free product</i>	1389
K. Rogers, <i>A note on orthogonal Latin squares</i>	1395
P. P. Saworotnow, <i>On continuity of multiplication in a complemented algebra</i>	1399
Johanan Schonheim, <i>On coverings</i>	1405
Victor Lenard Shapiro, <i>Bounded generalized analytic functions on the torus</i>	1413
James D. Stafney, <i>Arens multiplication and convolution</i>	1423
Daniel Sterling, <i>Coverings of algebraic groups and Lie algebras of classical type</i>	1449
Alfred B. Willcox, <i>Šilov type C algebras over a connected locally compact abelian group. II</i>	1463
Bertram Yood, <i>Faithful $*$-representations of normed algebras. II</i>	1475
Alexander Zabrodsky, <i>Covering spaces of paracompact spaces</i>	1489