

# Pacific Journal of Mathematics

**CUT POINTS IN TOTALLY NON-SEMI-LOCALLY-CONNECTED  
CONTINUA**

EDWARD EVERETT GRACE

# CUT POINTS IN TOTALLY NON-SEMI-LOCALLY-CONNECTED CONTINUA

EDWARD E. GRACE

1. **Introduction.** F. Burton Jones has shown [4, Theorem 15] that the set of weak cut points of a compact metric continuum that is not semi-locally-connected at any point of an open set  $U$  of  $M$  is dense in  $U$  (the proofs apply to locally peripherally compact complete metric continua). It is the purpose of this paper to extend Jones' results by establishing stronger cutting properties.

The theory is given here for locally compact metric spaces but applies, with appropriate modifications, to locally peripherally bicomact, regular spaces of the general class mentioned in [2].

2. **Definitions and preliminary theorems.** A point  $p$  of a continuum  $M$  is a *weak cut point* of  $M$  (or *cuts  $M$  weakly*) if there are two points  $x$  and  $y$  of  $M - \{p\}$  such that each subcontinuum of  $M$  that contains both  $x$  and  $y$  contains  $p$  also. In this case  $p$  *cuts  $x$  from  $y$  weakly* in  $M$ .

A continuum  $M$  is *semi-locally-connected* at a point  $p$  of  $M$  if each open subset  $U$  of  $M$  containing  $p$  contains an open subset  $V$  of  $M$  containing  $p$ , the complement of which relative to  $M$  consists of a finite number of components. A continuum  $M$  is *totally nonsemi-locally-connected* (on a point set  $A$ ) if  $M$  is not semi-locally-connected at any point (of  $A$ ). A continuum  $M$  is *locally peripherally aposyndetic* at a point  $p$  of  $M$  if each open subset  $U$  of  $M$  containing  $p$  contains an open subset  $V$  of  $M$  containing  $p$  such that, for some collection  $(H_1, \dots, H_n)$  of subcontinua of  $M$ ,  $(\bigcap_{i=1}^n H_i) \cap (\bar{V} - V) = \phi$  and  $p$  is in the (nonvoid) interior  $W$  of  $(\bigcap_{i=1}^n H_i) \cap V$ , relative to  $M$ . In this case  $W$  is a *peripheral aposyndesis subset* of  $U$  and  $V$  is a set *associated with  $W$  and  $U$* .

**EXAMPLE.** The Cartesian product of a cantor set and a simple closed curve, with one of the cantor sets shrunk to a point, is locally peripherally aposyndetic at each point but aposyndetic at only one point.

**THEOREM 1.** *A locally compact metric continuum  $M$  is locally peripherally aposyndetic on a dense  $G_\delta$  subset of an open subset  $D$*

---

Received July 9, 1962, and in revised form October 23, 1963. This work was completed while the author was on leave of absence from Emory University as an NSF Science Faculty Fellow at the University of Wisconsin.

of  $M$  if it is locally peripherally aposyndetic on a dense subset of  $D$ .

Theorem 1 is a corollary of [2, Theorem 1].

**THEOREM 2.** *If  $M$  is a locally compact metric continuum that is not semi-locally-connected at the point  $x$ , then  $x$  is a limit point of the set of points of  $M$  at which  $M$  is nonaposyndetic with respect to  $x$ .*

Theorem 2 follows from the fact that any open subset  $D$  of  $M$  which has a compact boundary and contains the point  $x$ , contains an open subset, containing  $x$ , whose complement with respect to  $M$  consists of a finite number of subcontinua of  $M$ , if  $M$  is aposyndetic at each point of  $\bar{D} - D$  with respect to  $x$ .

**THEOREM 3.** *If  $M$  is a locally compact metric continuum that is not locally peripherally aposyndetic at the point  $x$  then  $x$  is a limit point of the set of all points  $y$  of  $M$  such that  $M$  is nonaposyndetic at  $x$  with respect to  $y$ .*

Theorem 3 follows from the Heine-Borel Theorem by an indirect proof.

### 3. Cut point theorems.

**THEOREM 4.** *If  $M$  is a locally compact metric continuum which is locally peripherally aposyndetic and nonsemi-locally connected at each point of a  $G_\delta$  subset  $I$  dense in an open subset  $D$ , then for each point  $p$  of  $M$  there is a dense  $G_\delta$  subset  $J_p$  of  $D$  such that for each point  $x$  of  $J_p$  there is a point  $y$  at which  $M$  is nonaposyndetic with respect to  $x$  and  $x$  cuts  $p$  weakly from each such  $y$ .*

*Proof.* A point  $x$  of  $D$  has the properties required for membership in  $J_p$  if  $x \in I - \{p\}$  and each open subset  $D'$  of  $D$ , containing  $x$ , contains an open subset  $D_1$ , containing  $x$ , and a subset  $S$  such that (1)  $M$  is aposyndetic at each point of  $M - D'$  with respect to each point of  $D_1$  or (2)  $S$  separates  $p$  from each point  $y$  of  $M - D'$  at which  $M$  is nonaposyndetic with respect to some point of  $D_1$ . This observation suggests consideration of distinguished subsets defined as follows (see [2, p.     ]). An open subset  $D_1$  of an open subset  $D'$  of  $D$  is a distinguished subset of  $D'$  if

- (1)  $M$  is aposyndetic at each point of  $M - D'$  with respect to each point of  $D_1$  or
- (2) there is a subset  $S$  of  $D'$  which separates  $p$  from each point

$y$ , in  $M - D'$ , at which  $M$  is nonaposyndetic with respect to some point of  $D_1$ .

If each open set in  $D$  contains a distinguished open subset then [2, Theorem 1] the set of distinguished points of  $D$  is a dense  $G_\delta$  subset of  $D$  the intersection of which with  $I$  is the desired set  $J_p$ . To prove that each open set in  $D$  contains a distinguished open subset, let

(1)  $D'$  be any open subset of  $D$ ,

(2)  $S$  be a peripheral aposyndesis subset of  $D'$  that does not contain  $p$  (such a set exists since  $M$  is locally peripherally aposyndetic on a dense subset of  $D$ ) and

(3)  $D''$  be a set associated with  $S$  and  $D'$ . If  $M - S$  is connected then  $D_1 = S$  is a distinguished subset of  $D'$ , since  $M$  is aposyndetic at each point of  $M - D'$  with respect to each point of  $D_1$ . If  $M - S = A \cup B$ , where  $A$  and  $B$  are mutually separated and  $p \in A$ , then  $B \cap (D'' - S) \neq \phi$ . Let  $H$  be a continuum containing  $S$  but not containing  $B \cap (D'' - S)$ . Let  $D_1 = (M - H) \cap B \cap (D'' - S)$ . Then  $M$  is aposyndetic at each point of  $A$  with respect to each point of  $D_1$ , since  $H \cup A$  is a continuum containing  $A$  in its interior. But  $S$  separates  $p$  from each point of  $B$  and  $A \cup B \supset M - D'$ . It follows that  $D_1$  is a distinguished subset of  $D'$ .

That there is a space satisfying the hypothesis of Theorem 4 is seen from the example [3, Example 2] of a bounded plane continuum which is both connected im kleinen and nonsemi-locally-connected at each point of a dense  $G_\delta$  subset.

Local peripheral aposyndesis is used instead of aposyndesis in Theorem 4 in order that the complementary case (covered in Theorem 5) will be such that each  $x$  in  $J_p$  will be a limit point of  $A_x$ .

**COROLLARY 4.1.** *If a compact metric continuum  $M$  is aposyndetic, but not semi-locally-connected, at each point of a dense  $G_\delta$  subset of  $M$  and  $P$  is a countable, dense subset of  $M$ , then there is a dense  $G_\delta$  subset  $J$  of  $M$  each point  $x$  of which cuts each point  $p$  of  $P$  from each point  $y$  at which  $M$  is nonaposyndetic with respect to  $x$ .*

*Proof.* First, if  $M$  is aposyndetic at  $x$  then  $M$  is locally peripherally aposyndetic at  $x$ . Second, if  $p(1), p(2), \dots$  is a counting of  $P$  let  $J = \bigcap_{n=1}^{\infty} J_{p(n)}$ , where  $J_{p(n)}$  is as given in Theorem 4.

Theorem 5 is a corollary of [1, Theorem 2].

**THEOREM 5.** *If  $M$  is a locally compact metric continuum and  $D$  is an open subset of  $M$  such that  $M$  is nonlocally peripherally aposyndetic at each point of a dense  $G_\delta$  subset of  $D$ , then for each*

point  $p$  of  $M$  there is a dense  $G_\delta$  subset  $J_p$  of  $D$  such that each point  $x$  of  $J_p$

(1) is cut from  $p$  weakly by each point of the set  $A_x$ , of all points  $y \neq p$  such that  $M$  is nonaposyndetic at  $x$  with respect to  $y$ , and

(2) is a limit point of  $A_x$ .

The following theorem is a consequence of Theorems 4 and 5.

**THEOREM 6.** *If  $M$  is a locally compact metric continuum which is totally nonsemi-locally-connected on some dense  $G_\delta$  subset of an subset  $D$  of  $M$  then  $D$  is contained in the union of the closures of two (possibly void) open subsets  $D_1$  and  $D_2$  such that  $M$  is locally peripherally aposyndetic at each point of a  $G_\delta$  set dense in  $D_1$  but not at any point of  $D_2$  and for each point  $p$  of  $M$  there is a dense  $G_\delta$  subset  $J_p$  of  $D$  such that*

(1) each point  $x$  of  $J_p \cap D_1$  cuts  $p$  weakly from each point  $y$  such that  $M$  is nonaposyndetic at  $y$  with respect to  $x$ , and

(2) each  $x$  in  $J_p \cap D_2$  is cut from  $p$  weakly by each point of  $A_x = \{y \mid y \neq p \text{ and } M \text{ is nonaposyndetic at } x \text{ with respect to } y\}$  and is a limit point of  $A_x$ .

**COROLLARY 6.1.** *If  $M$  is a compact metric continuum which is totally nonsemi-locally-connected on some dense  $G_\delta$  set and  $P$  is a countable dense subset of  $M$  then  $M$  contains a dense set  $A$  each point of which cuts some point of  $M$  weakly from each point of  $P$ .*

*Proof.* Let  $p(1), p(2), \dots$  be a counting of  $P$  and let  $D_1$  and  $D_2$  be as given in Theorem 6. For each natural number  $n$ , let  $J_{p(n)}$  be as given in Theorem 6. Let  $J = \bigcap_{n=1}^{\infty} J_{p(n)}$ . Let  $B = J \cap D_1$  and  $C$  be the set of all  $x \in M$  such that  $M$  is nonaposyndetic at some point  $y$  of  $J \cap D_2$  with respect to  $x$ . Then  $\bar{C} \supset J \cap D_2$  and each point of  $C$  cuts some point of  $J \cap D_2$  weakly from each point of  $P$ . The union of  $B$  and  $C$  is the desired dense set of points.

*Question.* Does each totally nonsemi-locally-connected, compact metric continuum contain a dense  $G_\delta$  set of weak cut points?

#### REFERENCES

1. E. E. Grace, *Cut sets in totally nonaposyndetic continua*, Proc. Amer. Math. Soc., **9** (1958), 98-104.
2. ———, *On local properties and  $G_\delta$  sets*, to appear in Pacific J. Math., **14** (1964), 1245-1248.
3. ———, *Certain questions related to the equivalence of local connectedness and connectedness im kleinen*, to appear.
4. F. B. Jones, *Concerning non-aposyndetic continua*, Amer. J. Math., **70** (1948), 403-413.

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

ROBERT OSSERMAN

Stanford University  
Stanford, California

M. G. ARSOVE

University of Washington  
Seattle 5, Washington

J. DUGUNDJI

University of Southern California  
Los Angeles 7, California

LOWELL J. PAIGE

University of California  
Los Angeles 24, California

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

\* \* \*

AMERICAN MATHEMATICAL SOCIETY  
CALIFORNIA RESEARCH CORPORATION  
SPACE TECHNOLOGY LABORATORIES  
NAVAL ORDNANCE TEST STATION

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and on submission, must be accompanied by a separate author's résumé. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal but they are not owners or publishers and have no responsibility for its content or policies.

# Pacific Journal of Mathematics

Vol. 14, No. 4

August, 1964

Homer Franklin Bechtell, Jr., <i>Pseudo-Frattini subgroups</i> .....	1129
Thomas Kelman Boehme and Andrew Michael Bruckner, <i>Functions with convex means</i> .....	1137
Lutz Bungart, <i>Boundary kernel functions for domains on complex manifolds</i> .....	1151
L. Carlitz, <i>Rings of arithmetic functions</i> .....	1165
D. S. Carter, <i>Uniqueness of a class of steady plane gravity flows</i> .....	1173
Richard Albert Dean and Robert Harvey Oehmke, <i>Idempotent semigroups with distributive right congruence lattices</i> .....	1187
Lester Eli Dubins and David Amiel Freedman, <i>Measurable sets of measures</i> .....	1211
Robert Pertsch Gilbert, <i>On class of elliptic partial differential equations in four variables</i> .....	1223
Harry Gonshor, <i>On abstract affine near-rings</i> .....	1237
Edward Everett Grace, <i>Cut points in totally non-semi-locally-connected continua</i> .....	1241
Edward Everett Grace, <i>On local properties and <math>G_\delta</math> sets</i> .....	1245
Keith A. Hardie, <i>A proof of the Nakaoka-Toda formula</i> .....	1249
Lowell A. Hinrichs, <i>Open ideals in <math>C(X)</math></i> .....	1255
John Rolfe Isbell, <i>Natural sums and abelianizing</i> .....	1265
G. W. Kimble, <i>A characterization of extremals for general multiple integral problems</i> .....	1283
Nand Kishore, <i>A representation of the Bernoulli number <math>B_n</math></i> .....	1297
Melven Robert Krom, <i>A decision procedure for a class of formulas of first order predicate calculus</i> .....	1305
Peter A. Lappan, <i>Identity and uniqueness theorems for automorphic functions</i> .....	1321
Lorraine Doris Lavalley, <i>Mosaics of metric continua and of quasi-Peano spaces</i> .....	1327
Mark Mahowald, <i>On the normal bundle of a manifold</i> .....	1335
J. D. McKnight, <i>Kleene quotient theorems</i> .....	1343
Charles Kimbrough Megibben, III, <i>On high subgroups</i> .....	1353
Philip Miles, <i>Derivations on <math>B^*</math> algebras</i> .....	1359
J. Marshall Osborn, <i>A generalization of power-associativity</i> .....	1367
Theodore G. Ostrom, <i>Nets with critical deficiency</i> .....	1381
Elvira Rapaport Strasser, <i>On the defining relations of a free product</i> .....	1389
K. Rogers, <i>A note on orthogonal Latin squares</i> .....	1395
P. P. Saworotnow, <i>On continuity of multiplication in a complemented algebra</i> .....	1399
Johanan Schonheim, <i>On coverings</i> .....	1405
Victor Lenard Shapiro, <i>Bounded generalized analytic functions on the torus</i> .....	1413
James D. Stafney, <i>Arens multiplication and convolution</i> .....	1423
Daniel Sterling, <i>Coverings of algebraic groups and Lie algebras of classical type</i> .....	1449
Alfred B. Willcox, <i>Šilov type <math>C</math> algebras over a connected locally compact abelian group. II</i> .....	1463
Bertram Yood, <i>Faithful <math>*</math>-representations of normed algebras. II</i> .....	1475
Alexander Zabrodsky, <i>Covering spaces of paracompact spaces</i> .....	1489