

Pacific Journal of Mathematics

ON LOCAL PROPERTIES AND G_δ SETS

EDWARD EVERETT GRACE

ON LOCAL PROPERTIES AND G_δ SETS

EDWARD E. GRACE

1. **Introduction.** For complete metric spaces, R. H. Bing has shown [2, Theorem 2] that being not connected im kleinen at each point of a dense G_δ set implies the existence of an open set that contains uncountably many components. The proof, in fact, shows that there is an open set no component of which contains an open set. For Baire topological continua, the author has shown [4, Lemma 2] that being not connected im kleinen, at each point of a dense-domain intersection set, implies the existence of an open set no component of which contains an open set.

For a certain class of II_ϕ spaces [3, p. 642], including complete metric spaces and Baire topological spaces, there is a general theorem about local properties which has both of these theorems as corollaries. For simplicity, the theory is presented here only as it applies to the special case of complete metric spaces. Generalization to II_ϕ spaces, which have "developments" [1, p. 180] consisting of ϕ (perhaps not \aleph_0) collections of open sets, is rather straightforward. The pattern of this generalization is indicated to some extent by [4].

This theory applies to those local properties which the space has at a point x if and only if x is a distinguished point in the following sense. There is a relation "is a *distinguished subset* of" having the following two properties. (1) D' is a distinguished subset of D only if D' is open and is a subset of D . (2) If D contains D' and D'' is a distinguished subset of D' , then D'' is a distinguished subset of D . A point x is said to be a *distinguished point* if each open set containing x contains a distinguished subset containing x .

Several of the corollaries given here have been known for some time.

2. **Theorems.** The proof of Theorem 1 is complicated slightly in order that it will be clearer how it generalizes to apply to the general class of spaces referred to above.

THEOREM 1. *If M is a complete metric space each open set of which contains a distinguished open subset, then the set of distinguished points of M is a dense G_δ set.*

Received July 9, 1962, and in revised form September 26, 1963. Presented to the American Mathematical Society, November 18, 1961, under the title Local properties and Z category. This research was done while the author was on leave of absence from Emory University as an NSF Science Faculty Fellow at the University of Wisconsin.

Proof. For each $t < \aleph_0$, let U_t be the collection of open sets of diameter less than $1/t$ and let V_t be the union of all distinguished open subsets of members of U_t . The set V_t is dense and open since $U_t^* = \{x | x \in N \in U_t\}$ is dense and each open set contained in a member of U_t contains a distinguished open subset of that member. Let x be a point of the dense G_δ set $J = \bigcap_{t < \aleph_0} V_t$ and let D be an open set containing x . Then for some $t < \aleph_0$, the open set D contains $U_t^*(x)$. However, $x \in V_t$ and consequently x is contained in a distinguished open subset of some member N of U_t (containing x). But any distinguished open subset of N is a distinguished open subset of D , since $D \supset N$. It follows that J is a subset of the desired G_δ set. Clearly J contains each of the desired points. Hence, J is the desired dense G_δ set.

COROLLARY 1.1. *If each open set in a complete metric space M has a component with a nonvoid interior, then M is connected im kleinen at each point of some dense G_δ set.*

Proof. Let D' be a distinguished subset of D if D' is an open subset of a component of D .

COROLLARY 1.2. *If f is a function (not necessarily continuous) on a complete metric space and each open set D contains an open set D' such that $D \supset f(D')$, then $f(x) = x$ for each x in some dense G_δ set.*

Proof. Let D' be a distinguished subset of D if D' is an open subset of D and $D \supset f(D')$.

COROLLARY 1.3. *If M is a connected, compact metric space and each open set D contains a point p such that M is aposyndetic [5, p. 138] at p with respect to each point of $M - D$, then M is aposyndetic at each point of some dense G_δ set.*

Proof. Let D' be a distinguished subset of D if D' is open and D contains the intersection of a finite number of continua, each of which contains D' .

COROLLARY 1.4. *If M is a connected, complete metric space and the complement of each open set is contained in the union of a finite number of continua the union of which is not M , then M is semi-locally-connected [7, p. 19] at each point of a dense G_δ set.*

Proof. Let D' be a distinguished subset of D if $D \supset D'$ and the

complement of D' is the union of a finite number of continua.

Theorem 2 follows from (part of) the proof of Theorem 1.

THEOREM 2. *In a metric space, the set of distinguished points is a G_δ set.*

COROLLARY 2.1. *The set of points at which a metric space is connected im kleinen is a G_δ set.*

There are also corollaries to Theorem 2 corresponding of Corollaries 1.2, 1.3 and 1.4.

THEOREM 3. *In a complete metric space, if some dense G_δ set contains no distinguished point then there is an open set which contains no distinguished subset.*

Proof. Assume the conclusion is false. Then by Theorem 1 the set of distinguished points is a dense G_δ set. But any two dense G_δ subsets of a complete metric space have a nonvoid intersection. This contradicts the hypothesis.

COROLLARY 3.1. *If a complete metric space is nonconnected im kleinen at each point of a dense G_δ set then it contains an open set no component of which contains an open set (and which, therefore, has uncountably many components).*

There are also corollaries to Theorem 3 corresponding to Corollaries 1.2, 1.3 and 1.4.

Theorem 4 is proved by repeated application of Theorem 3 to open subspaces.

THEOREM 4. *In a complete metric space, if (1) some dense G_δ set contains no distinguished point and (2) the union of any collection of disjoint open sets contains a distinguished subset only if one of the members of the collection contains a distinguished subset, then some dense open subset contains no distinguished subset.*

COROLLARY 4.1. *If a complete metric space is nonconnected im kleinen at each point of a dense G_δ set then there is a dense open set no component of which contains an open set.*

COROLLARY 4.2. *Let M be a complete, metrizable, open subspace of a connected topological space T such that, at each point of some*

dense G_δ subset of M , the space T fails to be semi-locally-connected. Then there is an open set D , dense in M , such that the complement of each open subset of D has infinitely many components.

Proof. Let V be called a distinguished subset of an open subset W of M , if V is an open subset of W and the complement of V , in T , consists of a finite number of components. Here the union of a collection of disjoint open subsets of M is a distinguished subset of an open subset U of M containing it only if *each* member of the collection is a distinguished subset of U . Hence the corollary follows from Theorem 4.

REFERENCES

1. R. H. Bing, *Metrization of topological spaces*, Canad. J. Math., **3** (1951), 175-186.
2. ———, *A simple closed curve is the only homogeneous bounded plane continuum that contains an arc*, Canad. J. Math., **12** (1960), 209-230.
3. Nelson Dunford, *On continuous mapping*, Ann. of Math., **41** (1940), 639-661.
4. E. E. Grace, *Totally nonconnected im kleinen continua*, Proc. Amer. Math. Soc., **9** (1958), 818-821.
5. F. B. Jones, *Concerning aposyndetic and non-aposyndetic continua*, Bull. Amer. Math. Soc., **58** (1952), 137-151.
6. J. W. Tukey, *Convergence and uniformity in topology*, Annals of Mathematic Studies, no. 3, Princeton University Press, 1940.
7. G. T. Whyburn, Amer. Math. Soc. Colloquium Publications, vol. **28**, New York, 1942.

ARIZONA STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

ROBERT OSSERMAN

Stanford University
Stanford, California

M. G. ARSOVE

University of Washington
Seattle 5, Washington

J. DUGUNDJI

University of Southern California
Los Angeles 7, California

LOWELL J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and on submission, must be accompanied by a separate author's résumé. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 14, No. 4

August, 1964

Homer Franklin Bechtell, Jr., <i>Pseudo-Frattini subgroups</i>	1129
Thomas Kelman Boehme and Andrew Michael Bruckner, <i>Functions with convex means</i>	1137
Lutz Bungart, <i>Boundary kernel functions for domains on complex manifolds</i>	1151
L. Carlitz, <i>Rings of arithmetic functions</i>	1165
D. S. Carter, <i>Uniqueness of a class of steady plane gravity flows</i>	1173
Richard Albert Dean and Robert Harvey Oehmke, <i>Idempotent semigroups with distributive right congruence lattices</i>	1187
Lester Eli Dubins and David Amiel Freedman, <i>Measurable sets of measures</i>	1211
Robert Pertsch Gilbert, <i>On class of elliptic partial differential equations in four variables</i>	1223
Harry Gonshor, <i>On abstract affine near-rings</i>	1237
Edward Everett Grace, <i>Cut points in totally non-semi-locally-connected continua</i>	1241
Edward Everett Grace, <i>On local properties and G_δ sets</i>	1245
Keith A. Hardie, <i>A proof of the Nakaoka-Toda formula</i>	1249
Lowell A. Hinrichs, <i>Open ideals in $C(X)$</i>	1255
John Rolfe Isbell, <i>Natural sums and abelianizing</i>	1265
G. W. Kimble, <i>A characterization of extremals for general multiple integral problems</i>	1283
Nand Kishore, <i>A representation of the Bernoulli number B_n</i>	1297
Melven Robert Krom, <i>A decision procedure for a class of formulas of first order predicate calculus</i>	1305
Peter A. Lappan, <i>Identity and uniqueness theorems for automorphic functions</i>	1321
Lorraine Doris Lavalley, <i>Mosaics of metric continua and of quasi-Peano spaces</i>	1327
Mark Mahowald, <i>On the normal bundle of a manifold</i>	1335
J. D. McKnight, <i>Kleene quotient theorems</i>	1343
Charles Kimbrough Megibben, III, <i>On high subgroups</i>	1353
Philip Miles, <i>Derivations on B^* algebras</i>	1359
J. Marshall Osborn, <i>A generalization of power-associativity</i>	1367
Theodore G. Ostrom, <i>Nets with critical deficiency</i>	1381
Elvira Rapaport Strasser, <i>On the defining relations of a free product</i>	1389
K. Rogers, <i>A note on orthogonal Latin squares</i>	1395
P. P. Saworotnow, <i>On continuity of multiplication in a complemented algebra</i>	1399
Johanan Schonheim, <i>On coverings</i>	1405
Victor Lenard Shapiro, <i>Bounded generalized analytic functions on the torus</i>	1413
James D. Stafney, <i>Arens multiplication and convolution</i>	1423
Daniel Sterling, <i>Coverings of algebraic groups and Lie algebras of classical type</i>	1449
Alfred B. Willcox, <i>Šilov type C algebras over a connected locally compact abelian group. II</i>	1463
Bertram Yood, <i>Faithful $*$-representations of normed algebras. II</i>	1475
Alexander Zabrodsky, <i>Covering spaces of paracompact spaces</i>	1489