

# Pacific Journal of Mathematics

**FAITHFUL \*-REPRESENTATIONS OF NORMED ALGEBRAS. II**

BERTRAM YOOD

# FAITHFUL \*-REPRESENTATIONS OF NORMED ALGEBRAS II

BERTRAM YOOD

1. **Introduction.** Let  $A$  be a complex Banach algebra with an involution  $x \rightarrow x^*$ . By the positive cone  $P$  of  $A$  is meant the closure, in the set  $H$  of self-adjoint elements of  $A$ , of the set of all finite sums of elements of the form  $x^*x$ . Kelley and Vaught [5] have shown that, if  $A$  has an identity,<sup>1</sup>  $A$  has a faithful \*-representation (as bounded linear operators on a Hilbert space) if and only if (1)  $x \rightarrow x^*$  is continuous and (2)  $P \cap (-P) = (0)$ . Consider the (incomplete) normed algebra case. Examples exist with a faithful \*-representation and both conditions false, with (1) true and (2) false, and with (1) false and (2) true. Moreover, even if (1) holds so that  $x \rightarrow x^*$  extends to the completion  $A_c$  of  $A$ , one can have a continuous faithful \*-representation for  $A$  when none exists for  $A_c$ . It follows that the results which we now describe, even for the normed algebra case, can *not* be deduced from the theory of Banach algebras.

These facts led us to consider the development of a theory of \*-representations of a complex algebra  $A$  with involution (with or without an identity) under minimal assumptions on  $A$  but with results sufficiently definitive to illuminate the counter-examples mentioned above. We suppose that the real linear space  $H$  has a norm in terms of which it is a real normed linear space such that

(a) the real subalgebra generated by each  $h \in H$  is a normed algebra and

(b) the Jordan product  $x \cdot h = xh + hx$  is a continuous function on  $H$  for each fixed  $h \in H$ .

It is shown that  $A$  has a faithful \*-representation continuous on  $H$  if and only if  $A$  is semi-simple and  $P \cap (-P) = (0)$ . If  $A$  is a normed \*- $Q$ -algebra, any \*-representation is automatically continuous on  $H$  so that these conditions are necessary and sufficient there for a faithful \*-representation. As already noted, this can fail if the  $Q$ -algebra hypothesis is dropped.

For previous work on \*-representations we refer to [5], [7], [8], and [10].

2. **Preliminaries.** Let  $A$  be an algebra over the complex field

---

Received October 23, 1963. This research was supported by the National Science Foundation Grant NSF-G-25219.

<sup>1</sup> As pointed out in [10, p. 352] this statement is incorrect if  $A$  has no identity. For a version covering that case see [10, Theorem 3.4]. Theorem 4.3 below shows that  $A$  has a faithful \*-representation if and only if  $A$  is semi-simple and  $P \cap (-P) = (0)$ .

with an involution  $x \rightarrow x^*$ . The set of self-adjoint (s.a.) elements of  $A$  is denoted by  $H$ . By a  $*$ -representation of  $A$  we mean a homomorphism  $x \rightarrow T_x$  of  $A$  into the algebra of bounded linear operators on a Hilbert space where, for each  $x$ ,  $T_{x^*}$  is the adjoint of  $T_x$ . A  $*$ -representation which is one to one is called *faithful*. A general representation procedure of Gelfand and Naimark [7] which we adapt to our needs leads to  $*$ -representations via positive linear functionals.

A complex linear functional  $f$  on  $A$  is called *positive* if  $f(x^*x) \geq 0$  for all  $x \in A$ . We call  $f$  *hermitian* if  $f(x^*) = \overline{f(x)}$  for all  $x \in A$  or equivalently if  $f$  is a real linear functional when restricted to the real linear space  $H$ . As in [8, p. 200] we define  $L_f = \{x : f(zx) = 0 \text{ for all } z \in A\} = \{x : f(x^*x) = 0\}$ ;  $L_f$  is a left ideal of  $A$ . Let  $X_f$  be the linear space  $A - L_f$  and  $\pi$  be the natural homomorphism of  $A$  onto  $X_f$ . Then, [8, p. 212],  $(\pi(x), \pi(y)) = f(y^*x)$  defines an inner product on  $X_f$  in terms of which  $X_f$  is a pre-Hilbert space. Let  $H_f$  be the completion of  $X_f$  in the pre-Hilbert space norm. As in [7, p. 120] we associate with  $y \in A$  a linear operator  $T_y^f$  defined on  $X_f$  by the rule  $T_y^f[\pi(x)] = \pi(yx)$ . In order that every  $T_y^f$ ,  $y \in A$ , be extendable to a bounded linear operator  $U_y^f$  on  $H_f$  it is necessary and sufficient [8, p. 213] that  $f$  be *admissible*, that is, to each  $x \in A$  there corresponds a number  $K(x) < \infty$  such that  $f(y^*x^*xy) \leq K(x)f(y^*y)$  for all  $y \in A$ . If  $f$  is admissible, the mapping  $x \rightarrow U_x^f$  is a  $*$ -representation of  $A$ .

For any positive linear functional  $f$  and any  $y \in A$  we define the positive linear functional  $f_y(x) = f(y^*xy)$ .

**2.1. LEMMA.** *Let  $f$  be a positive linear functional on  $A$ . Then  $f$  is admissible if and only if*

$$(2.1) \quad \sup_n [f_y(h^{2^n})]^{2^{-n}} < \infty,$$

for each  $y \in A$ ,  $h \in H$ , where the sup is taken over the set of positive integers.

Suppose that  $f$  is admissible. Then, for  $h \in H$ ,  $U_h^f$  is a bounded s.a. operator on the Hilbert space  $H_f$ . For convenience, let  $U_z^f$  where  $z = h^{2^n}$  be denoted by  $V_n$ . For each  $y \in A$ ,

$$f_y(h^{2^{n+1}}) = \|V_n \pi(y)\|^2 \leq \|U_h^f\|^{2^{n+1}} f(y^*y)$$

for  $n = 0, 1, 2, \dots$ . This implies (2.1).

For the converse we make use of an inequality due to Kaplansky [4, p. 55] concerning a positive linear functional  $f$  which asserts that

$$(2.2) \quad f_y(x^*x) \leq f(y^*y)^{1-2^{-n}} [f_y((x^*x)^{2^n})]^{2^{-n}}$$

for all  $x, y \in A$  and all positive integers  $n$ . Assume (2.1). It is clearly sufficient to show that  $T_h^f$  is a bounded operator on  $X_f$  for each  $h$  s.a. Using (2.2) we have

$$\|T_h^f \pi(y)\|^2 = f_y(h^2) \leq f(y^*y)^{1-2^{-n}} [f_y(h^{2^{n+1}})]^{2^{-n}}$$

so that  $\|T_h^f\|$  cannot exceed the sup of (2.1).

**2.2. LEMMA.** *Suppose  $H$  is given a topology in which it is a real linear topological space. Then the mapping  $p \rightarrow (1+h)x(1+h)$  is continuous on  $H$ , for each  $h \in H$ , if and only if the Jordan product  $x \cdot h = hx + xh$  is continuous on  $H$  for each  $h \in H$ .*

Let  $a(x, h) = x + xh + hx + h x h$ . Then  $x \cdot h = [a(x, h) - a(x, -h)]/2$  and  $a(x, h) = x + x \cdot h + [(x \cdot h) \cdot h - x \cdot h^2]/2$  from which the lemma is immediate.

We now state metric requirements which we put on the algebra  $A$  with involution. We suppose given a norm  $\|h\|$  on  $H$  in terms of which  $H$  is a real normed linear space and, for each  $h \in H$ , the real subalgebra generated by  $h$  is a normed algebra. No assumptions are made about the elements not in  $H$  nor are there any requirements of completeness or identity element. We assume that the Jordan product  $x \cdot h$  is continuous on  $H$  for each  $h \in H$ . We call  $A$  a *normed \*-algebra* if,  $A$  is a normed algebra. Following [3] we say that the normed \*-algebra  $A$  is a *normed  $Q$ -\*-algebra* if the set of quasi-regular elements of  $A$  is open. If  $A$  is a Banach algebra it has this property [3, p. 155].

For  $h \in H$ ,  $\lim \|h^n\|^{1/n} = \nu(h)$  exists. Clearly  $\nu(h) \leq \|h\|$  and  $\nu(h^2) = [\nu(h)]^2$  (see [8, p. 10]).

**2.3. LEMMA.** *Let  $f$  be a positive linear functional on  $A$ . The following statements are equivalent.*

- (a) *Each  $f_y$  is continuous on  $H$ .*
- (b)  *$f_y(x^*x) \leq \nu(x^*x)f(y^*y)$  for all  $x, y \in A$ .*
- (c)  *$f$  is admissible and the mapping  $x \rightarrow T_x^f$  is continuous on  $H$ .*

Suppose (a) holds. From the inequality (2.2) we obtain

$$f_y(x^*x) \leq f(y^*y)^{1-2^{-n}} (\|f_y\|_H \|x^*x\|^{2^n})^{2^{-n}}.$$

If we let  $n \rightarrow \infty$  we obtain (b).

Suppose (b) holds. Clearly  $f$  is admissible. For  $h \in H$  we have  $f_y(h^2) \leq \nu(h)^2 f(y^*y)$  so that  $\|T_h^f \pi(y)\| \leq \|h\| \|\pi(y)\|$  and  $\|T_h^f\| \leq \|h\|$ .

Suppose (c) with  $\|T_h^f\| \leq k\|h\|$ ,  $h \in H$ . Then, by the Cauchy-Schwarz inequality,

$$|f_y(h)|^2 \leq f(y^*y)f_y(h^2) = f(y^*y) \|T_h^f \pi(y)\|^2 \leq k^2 \|h\|^2 [f(y^*y)]^2$$

so that  $f_y$  is continuous on  $H$ .

We note that, under these conditions, the norm of the mapping  $x \rightarrow T_x^f$  on  $H$  does not exceed one.

**2.4. LEMMA.** *Any \*-representation of a normed \*-Q-algebra  $A$  is continuous on  $H$ .*

Let  $x \rightarrow T_x$  be a \*-representation of  $A$ . Let  $\rho(u)$  denote the spectral radius of  $u$  [8, p. 30]. For  $h \in H$  we have  $\|T_h\| = \rho(T_h) \leq \rho(h) \leq \|h\|$  by [9, p. 373]. Thus in the  $Q$ -algebra case the admissible positive linear functionals are those satisfying (b) of Lemma 2.3; if also  $A$  has an identity the admissible positive linear functionals are those continuous on  $H$ .

**2.5. LEMMA.** *Suppose  $f$  is positive linear functional on  $A$  which is continuous on  $H$ . Then  $f_y$  is continuous on  $H$  for each  $y \in A$ .*

It follows from Lemma 2.2 that the mapping  $x \rightarrow h x h$  is continuous on  $H$  for each  $h \in H$ . Therefore the functional  $f_h$  is continuous on  $H$  for each  $h \in H$ . Now, if  $y = u + iv$ ,  $u, v \in H$  we have  $f_y(x) = f_u(x) + f_v(x) + if(uxv - vxu)$ . But, by the Cauchy-Schwarz inequality, for any  $x \in H$ ,  $|f(uxv)|^2 \leq f(u^2)f_v(x^2) \leq f(u^2)\|f_v\|\|x\|^2$  where  $\|f_v\|$  is the norm of  $f_v$  considered as a linear functional on  $H$ . This makes  $\|f_y(x)\| \leq K\|x\|$ ,  $x \in H$ , where

$$K = \|f_u\| + \|f_v\| + 2[f(u^2)\|f_v\| + f(v^2)\|f_u\|]^{1/2}.$$

In view of Lemma 2.3,  $f$  is admissible.

We give an example of a normed \*-algebra  $A$  whose involution is continuous with the following properties.

- (1)  $A$  has a faithful \*-representation.
- (2) Every \*-representation of  $A$  other than the zero representation is discontinuous on  $H$ .
- (3) The completion  $A_c$  of  $A$  has only the zero \*-representation.

Let  $A$  be the set of all polynomials in the complex variable  $z$  which vanish at the origin. For  $p(z) = \sum \alpha_k z^k$  we define  $p^*(z) = \sum \bar{\alpha}_k z^k$  and  $\|p(z)\| = \sum |\alpha_k|/k!$ . Then (see [3, p. 158])  $A$  is a normed \*-algebra. That (1) holds will be pointed out in §4. Let  $p \rightarrow T_p$  be a \*-representation of  $A$  continuous on  $H$ . The polynomial  $z$  is s.a.. For each real scalar  $\lambda$ ,  $\|\lambda^n z^n\| \rightarrow 0$ . Therefore  $\|\lambda^n T_z^n\| = |\lambda|^n \|T_z\|^n \rightarrow 0$ . This makes  $T_z = 0$  so that  $T_p = 0$  on  $A$ . Now the involution on  $A$ , being bicontinuous, extends to an involution on  $A_c$ . Any \*-representation  $x \rightarrow V_x$  of the Banach algebra  $A_c$  must be continuous by [8, Theorem

4.1.20]. Therefore, by the above,  $V_x = 0$  for all  $x \in A_*$ .

Let  $F$  be a set of admissible positive linear functionals on  $A$ . We call  $F$  a *compatible set* if for each  $x \in A$  there exists a real number  $K(x)$  such that  $\|U_x^f\| \leq K(x)$  for all  $f \in F$ . This is equivalent to requiring that, for each  $x \in A$ , there exists  $C(x) < \infty$ , such that  $f_y(x^*x) \leq C(x)f(y^*y)$  for all  $y \in A$  and all  $f \in F$ . By Lemmas 2.3 and 2.4 the set of all admissible positive linear functionals on a normed \*-Q-algebra is a compatible set.

For each  $f$  in the compatible set  $F$  consider the Hilbert space  $H_f$  and the corresponding \*-representation  $x \rightarrow U_x^f$ . Let  $H$  be the Hilbert space direct sum of the Hilbert spaces  $H_f$ . Since  $\|U_x^f\| \leq K(x)$  for all  $f \in F$  we can take [7, p. 113] the direct sum  $x \rightarrow U_x$  of the \*-representations  $x \rightarrow U_x^f$ ,  $f \in F$  where  $U_x$  is a bounded operator on  $H$  and  $\|U_x\| \leq K(x)$ . We call this \*-representation the *canonical \*-representation of  $A$  induced by  $F$* . For a left ideal  $L$  of  $A$  we use the notation  $(L : A)$  as in [8, p. 53] to denote the set of all  $x \in A$  such that  $xA \subset L$ . The kernel of the canonical \*-representation induced by  $F$  is given by  $\bigcap (L_f : A)$  where the intersection is taken over all  $f \in F$ .

**3. On \*-representations.** For our purposes we wish to define the \*-radical  $\mathfrak{R}^*$  of  $A$  as the intersection of the kernels of all \*-representations of  $A$  which are continuous on  $H$ . Let  $A^*$  denote the set of all positive linear functionals on  $A$ . At the outset we consider three subsets of  $A^*$ . Let  $\mathfrak{B} = \{f \in A^* : f_y(x^*x) \leq \nu(x^*x)f(y^*y), \text{ for all } x, y \in A\}$ . Let  $\mathfrak{D}$  be the set of *dual functionals* by which we mean  $\{f \in A^* : f \text{ is hermitian and } f \text{ is continuous on } H\}$ . Let  $\mathfrak{E} = \{f \in D : |f(x)|^2 \leq f(x^*x) \text{ for all } x \in A\}$ . By Lemmas 2.3 and 2.5 we see that  $\mathfrak{B} \supset \mathfrak{D} \supset \mathfrak{E}$  and that these are compatible sets. Let  $\mathfrak{B}_0$ ,  $\mathfrak{D}_0$ , and  $\mathfrak{E}_0$  be the kernels of the canonical \*-representations of  $A$  induced by  $\mathfrak{B}$ ,  $\mathfrak{D}$ , and  $\mathfrak{E}$  respectively. Then  $\mathfrak{E}_0 \supset \mathfrak{D}_0 \supset \mathfrak{B}_0$ .

**3.1. LEMMA.**  $\mathfrak{R}^* = \mathfrak{E}_0 = \mathfrak{D}_0 = \mathfrak{B}_0$ .  $A/\mathfrak{R}^*$  is semi-simple.

For any  $f \in \mathfrak{B}$ , and  $x, y \in A$ ,  $\|T_x^f \pi(y)\|^2 \leq \nu(x^*x) \|\pi(y)\|^2$  so that  $\|T_x^f\| \leq \nu(x^*x)^{1/2}$ . Consequently  $\|T_h^f\| \leq \nu(h) \leq \|h\|$ ,  $h \in H$ . Therefore if  $x \rightarrow T_x$  is any of the canonical \*-representations in question,  $\|T_h\| \leq \|h\|$ ,  $h \in H$ , and the \*-representation is continuous on  $H$ . This proves that  $\mathfrak{R}^* \subset \mathfrak{B}_0 \subset \mathfrak{D}_0 \subset \mathfrak{E}_0$ . We show that  $\mathfrak{E}_0 \subset \mathfrak{R}^*$ .

Let  $x \rightarrow V_x$  be any \*-representation of  $A$  continuous on  $H$ , say as operators on the Hilbert space  $M$ . For each  $\alpha \in M$  the functional  $g^\alpha(x) = (V_x(\alpha), \alpha)$  is continuous on  $H$  and is a dual functional. For  $\alpha$  in the unit ball  $\Sigma$  of  $M$ ,  $|g^\alpha(x)|^2 \leq \|V_x(\alpha)\|^2 = g^\alpha(x^*x)$  so that  $g^\alpha \in \mathfrak{E}$ . We have

$$\begin{aligned}
\mathfrak{G}_0 &= \bigcap_{f \in E} (L_f : A) \subset \bigcap_{\alpha \in \Sigma} (L_{g^\alpha} : A) \\
&= \bigcap_{\alpha \in \Sigma} \{z \in A : g^\alpha((zy)^*(zy)) = 0 \text{ for all } y \in A\} \\
&= \bigcap_{\alpha \in \Sigma} \{z \in A : V_{zy}(\alpha) = 0 \text{ for all } y \in A\} \\
&= \{z \in A : V_{zy} = 0 \text{ for all } y \in A\} \\
&\subset \{z \in A : \|V_{zz^*}\| = \|V_z\|^2 = 0\}.
\end{aligned}$$

Therefore  $\mathfrak{G}_0 \subset \mathfrak{R}^*$ .

Since  $A/\mathfrak{R}^* = A/\mathfrak{G}_0$  is algebraically \*-isomorphic to a \*-subalgebra of the algebra of all bounded linear operators on a Hilbert space, we see from [8, Theorem 4.1.19] that  $A/\mathfrak{R}^*$  is semi-simple. From this we see also that the radical of  $A$  is contained in  $\mathfrak{R}^*$ .

**3.2. LEMMA.** *A normed \*-algebra  $A$  has a faithful \*-representation continuous on  $H$  if and only if  $\mathfrak{R}^* = (0)$ .*

Suppose  $\mathfrak{R}^* = (0)$ . The preceding Lemma 3.1 then asserts the canonical \*-representations induced by  $\mathfrak{B}$ ,  $\mathfrak{D}$ , or  $\mathfrak{G}$  are faithful. As noted above, these \*-representations are continuous on  $H$ . We naturally seek conditions on  $A$  which force  $\mathfrak{R}^* = (0)$ .

We set forth notation which will be used below. Let  $R_0$  be the collection of all finite sums of elements of  $A$  of the form  $x^*x$  and let  $P$  be the closure of  $R_0$  in  $H$ . The set  $P$  will be considered as a closed cone in the real normed linear space  $H$ . Let  $A_1$  be the algebra obtained by adjoining an identity  $e$  to  $A$ . As usual the involution on  $A$  is extended to  $A_1$  by  $(\lambda e + x)^* = \bar{\lambda}e + x^*$  where  $\lambda$  is a scalar and  $x \in A$ . We shall have occasion to consider the sets  $H$ ,  $\mathfrak{R}^*$ ,  $\mathfrak{D}$ ,  $R_0$ , and  $P$  in  $A$  simultaneously with the corresponding sets defined for  $A_1$ . When we do so, we denote the latter sets by  $H_1$ ,  $\mathfrak{R}_1^*$ ,  $\mathfrak{D}_1$ ,  $R_{01}$ , and  $P_1$  respectively. The given norm on  $H$  leads to a norm on  $H_1$  via  $\|\lambda e + h\| = |\lambda| + \|h\|$ ,  $\lambda$  real,  $h \in H$ .  $A_1$  satisfies the requirements of our theory.

We set  $Z(\mathfrak{D}) = \bigcap f^{-1}(0)$ ,  $f \in \mathfrak{D}$  and  $Z(\mathfrak{G}) = \bigcap f^{-1}(0)$ ,  $f \in E$ . We define two versions of the *reducing ideal* [7, p. 130] suitable for this setting. Let  $\bigcap L_f$  where  $f$  runs over  $\mathfrak{D}(\mathfrak{G})$  be denoted by  $RI(\mathfrak{D})$  and  $RI(\mathfrak{G})$  respectively.

Let  $g$  be a continuous real linear functional on  $H$ ,  $g(P) \geq 0$ . If we extend  $g$  to  $A$  by the rule  $g(x) = g(h_1) + ig(h_2)$  for  $x = h_1 + ih_2$ ,  $h_1, h_2$  s.a., we obtain an element of  $\mathfrak{D}$ . Conversely the restriction to  $H$  of any  $f \in D$  has the property that  $g(P) \geq 0$ . From the theory of closed cones in a normed linear space [5, Lemma 1.2] it follows that  $P \cap (-P)$  is the s.a. part of  $Z(\mathfrak{D})$  so that  $Z(\mathfrak{D}) = P \cap (-P) + iP \cap (-P)$ . In a more restrictive context, this was pointed out and used in [1].

It will appear that  $Z(\mathfrak{E})$  can differ from  $Z(\mathfrak{D})$ ;  $Z(\mathfrak{E})$  does not seem to have as neat an interpretation as  $Z(\mathfrak{D})$ . For that reason the results of § 4 involving  $\mathfrak{D}$  are more interesting than the theory for  $\mathfrak{E}$ .

3.3. LEMMA.  $\mathfrak{R}^* = (RI(\mathfrak{D}) : A) = (RI(\mathfrak{E}) : A)$

We see, by Lemma 3.1, that  $\mathfrak{R}^*$  is the kernel of the canonical \*-representation induced by  $\mathfrak{D}$ . Thus

$$\begin{aligned}\mathfrak{R}^* &= \bigcap_{f \in \mathfrak{D}} (L_f : A) = \bigcap_{f \in \mathfrak{D}} \{x : xy \in L_f, \text{ for all } y \in A\} \\ &= \left\{ x : xy \in \bigcap_{f \in \mathfrak{D}} L_f, \text{ for all } y \in A \right\} = (RI(\mathfrak{D}) : A) .\end{aligned}$$

Let  $x \rightarrow V_x$  be the canonical \*-representation induced by  $\mathfrak{E}$ . We show, by direct computation, (see [7, p. 132]) that

$$(3.1) \quad \|V_x\|^2 = \sup_{f \in \mathfrak{E}} f(x^*x), \quad x \in A .$$

Let  $\beta(x)$  denote the right hand side of (3.1). Take  $f \in \mathfrak{E}$ . Then  $|f_y(x)|^2 = |f(y^*xy)|^2 \leq f(y^*y)f_y(x^*x)$  by the Cauchy-Schwarz inequality. Therefore  $f_y \in \mathfrak{E}$  whenever  $f(y^*y) \leq 1$ . Now  $\|T'_x \pi(y)\|^2 = f_y(x^*x)$  so that  $\|T'_x\|^2 \leq \beta(x)$  from which we see that  $\|V_x\|^2 \leq \beta(x)$ . On the other hand, for  $f \in \mathfrak{E}$ ,

$$[f(x^*x)]^2 \leq f(x^*xx^*x) = \|T'_x \pi(x)\|^2 \leq \|T'_x\|^2 f(x^*x)$$

which shows that  $\beta(x) \leq \|V_x\|^2$ .

From Lemma 3.1 we observe that  $\mathfrak{R}^* = \{x : f(x^*x) = 0, \text{ for all } f \in \mathfrak{E}\} = RI(\mathfrak{E})$ . This formula, as we shall see in § 4, can be invalid if  $\mathfrak{E}$  is replaced by  $\mathfrak{D}$ .

We consider next a version of Kelley and Vaught's result [5, Theorem 4.4].

3.4. THEOREM. Let  $x \rightarrow V_x$  be the conical \*-representation of  $A$  induced by  $\mathfrak{E}$ . Then  $\|V_x\|^2 = \text{dist}(-x^*x, P_1)$ .

Let  $h \in H$ ,  $\|h\| \leq 1$ . In the algebra  $A_1$  let  $B$  be the real subalgebra generated by  $e$  and  $h$  and let  $B_o$  be its completion. For  $m = 1, 2, \dots$  let

$$(3.2) \quad w_m = \sum_{k=1}^m \binom{1/2}{k} (-1)^k h^k .$$

Clearly  $w_m \in H$ . In  $B_o$  we have  $(e - h) = [\lim (e + w_m)]^2$  so that, in  $A_1$ , we get

$$(3.3) \quad e - h = \lim_m (e + w_m)^2 .$$



This shows that, in  $H_1$ ,  $e$  is an interior point of the cone  $P_1$ .

The discussion in [6, p. 96] shows that any  $f \in \mathfrak{E}$  is extendable to  $A_1$  so as to belong to  $\mathfrak{D}_1$  where  $f(e) \leq 1$ . On the other hand if  $g \in D_1$ ,  $g(e) \leq 1$  then its restriction to  $A$  lies in  $\mathfrak{E}$  by the Cauchy-Schwarz inequality. Since  $\|e\| = 1$  and  $e$  is an interior point of  $P_1$  we see, from Lemma 1.3 of [5], that, for each  $x \in A$ ,

$$\text{dist}(-x^*x, P_1) = \sup_{\substack{f \in \mathfrak{D}_1 \\ f(e) \leq 1}} f(x^*x) = \sup_{f \in \mathfrak{E}} f(x^*x).$$

An application of formula (3.1) completes the proof.

#### 4. Faithful \*-representations.

##### 4.1. THEOREM.

- (a)  $Z(\mathfrak{D})$  is a two-sided ideal of  $A$ .
- (b)  $Z(\mathfrak{D}) \subset RI(\mathfrak{D}) \subset \mathfrak{R}^*$  and the inclusions can be proper.
- (c) If  $A$  has an identity,  $Z(\mathfrak{D}) = RI(\mathfrak{D}) = \mathfrak{R}^*$ .
- (d) If  $x \in \mathfrak{R}^*$  then  $x^3 \in Z(\mathfrak{D})$ .
- (e)  $\mathfrak{R}^*$  is the complete inverse image of the radical of  $A/Z(\mathfrak{D})$  under the natural homomorphism of  $A$  onto  $A/Z(\mathfrak{D})$ .

We refer to formula (3.2) for notation. For each  $m = 1, 2, \dots$  we define the operator  $\alpha_m$  on  $H$  by the rule  $\alpha_m(x) = (e + w_m)x(e + w_m)$ . Since  $\alpha_m(x^*x) = (x + xw_m)^*(x + xw_m)$  we see that  $\alpha_m(R_0) \subset R_0$ . Because  $\alpha_m$  is continuous on  $H$  by Lemma 2.2, we also get  $\alpha_m(P) \subset P$ .

Suppose next that also  $h \in P$ . Then  $(e + w_m)h(e + w_m) = h(e + w_m)^2 \in P$ . Passing to the limit as  $m \rightarrow \infty$  we see from (3.3) that  $h - h^2 \in P$ . We have established that, for any  $h \in P$  whatever its norm,  $f(h)\|h\| \geq f(h^2) \geq 0$ ,  $f \in \mathfrak{D}$ . By the Cauchy-Schwarz inequality  $|f(hx)|^2 \leq f(h^2)f(x^*x)$  and  $|f(xh)|^2 \leq f(xx^*)f(h^2)$ ,  $f \in \mathfrak{D}$ . Now  $P \cap (-P) = \{y \in H : f(y) = 0, f \in \mathfrak{D}\}$ , so that  $f(yx) = 0 = f(xy)$  for all  $x \in A$ ,  $f \in \mathfrak{D}$ . Next let  $w \in Z(\mathfrak{D})$ . We can write  $w = y_1 + iy_2$  where each  $y_k \in P \cap (-P)$ . We then see that  $f(wx) = 0 = f(xw)$  for all  $f \in \mathfrak{D}$ ,  $x \in A$ , so that  $wx$  and  $xw$  lie in  $Z(\mathfrak{D})$ . This establishes (a).

Let  $x \in Z(\mathfrak{D})$ . By (a) we see that  $x^*x \in Z(\mathfrak{D})$  so that  $f(x^*x) = 0$  for all  $f \in \mathfrak{D}$ . Thus  $Z(\mathfrak{D}) \subset RI(\mathfrak{D})$ . Next let  $x \in RI(\mathfrak{D})$ ,  $y \in A$ . Then  $xy \in RI(\mathfrak{D})$  so that  $x \in (RI(\mathfrak{D}) : A) = \mathfrak{R}^*$  by Lemma 3.3.

We now produce an example for which  $Z(\mathfrak{D}) \neq RI(\mathfrak{D})$ . Let  $A = C([0, 1])$  with the usual norm and involution but considered as a zero algebra. Then all linear functionals on  $A$  are positive. This implies that  $RI(\mathfrak{D}) = A$ . On the other hand it is trivial that  $Z(\mathfrak{D}) = (0)$ .

We now provide an instance where  $RI(\mathfrak{D}) \neq \mathfrak{R}^*$ . Let  $q(w)$  be the function  $q(w) = w$  on  $[0, 1]$ . Again we take  $A = C([0, 1])$  with the

usual norm and involution but define the product by the rule  $xy = x(0)y(0)q$ . Under these definitions  $A$  is a Banach algebra and  $A^3 = (0)$ . Since the radical of  $A$  is contained in  $\mathfrak{R}^*$  by Lemma 3.1, we see that  $\mathfrak{R}^* = A$ . Now for any linear functional  $f$  on  $A$ ,  $f(x^*x) = |x(0)|^2 f(q)$ . By the Hahn-Banach theorem, there exists a continuous real linear functional  $g$  on  $H$  such that  $g(q) = 1$ . We extend  $g$  to  $A$  by the rule  $g(h_1 + ih_2) = g(h_1) + ig(h_2)$  where  $h_1, h_2 \in H$ . Then  $g \in \mathfrak{D}$ . If  $x \in RI(\mathfrak{D})$ ,  $g(x^*x) = |x(0)|^2 = 0$ . Thus  $RI(\mathfrak{D}) = \{x \in A : x(0) = 0\}$ . This completes the proof of (b).

Suppose that  $A$  has an identity  $e$ . For any  $x \in \mathfrak{R}^*$ ,  $x = xe \in RI(\mathfrak{D})$  by Lemma 3.3. Next take  $x \in RI(\mathfrak{D})$ . Since  $|f(x)|^2 \leq f(e)f(x^*x) = 0$ , for all  $f \in \mathfrak{D}$ , we see that  $x \in Z(\mathfrak{D})$ . Combining this information with the set inequalities of (b) we obtain (c).

By Lemma 3.1 there exists a \*-representation of  $A_1$  continuous on  $H_1$  with kernel  $\mathfrak{R}_1^*$ . By restricting this \*-representation to  $A$  we see that

$$(4.1) \quad \mathfrak{R}^* \subset A \cap \mathfrak{R}_1^*.$$

Let  $\lambda$  be a scalar and  $x, y \in A$ . Then  $y^*(\lambda e + x)^*(\lambda e + x)y = (\lambda y + xy)^*(\lambda y + xy) \in R_0$ . Thus  $y^*R_{01}y \subset R_0$  for each  $y \in A$ . From Lemma 2.2 it can be seen that, for  $h \in H$ , the mapping  $x \rightarrow h x h$  is continuous on  $H$ . It is easily shown that  $x \rightarrow h x h$  is also continuous on  $H_1$ . It then follows that  $hP_1h \subset P$ . This shows that  $h[P_1 \cap (-P_1) + iP_1 \cap (-P)]h \subset P \cap (-P) + iP \cap (-P)$ . By (c) this gives

$$(4.2) \quad h\mathfrak{R}_1^*h \subset Z(\mathfrak{D}), h \in H.$$

From (4.1) and (4.2) we have  $h\mathfrak{R}^*h \subset Z(\mathfrak{D})$ . It follows readily that  $uzw + wzu \in Z(\mathfrak{D})$  for all  $u, w \in H$  and  $z \in \mathfrak{R}^*$ . Let  $x = u + iv \in \mathfrak{R}^*$ ,  $u, v \in H$  and note that  $u, v \in \mathfrak{R}^*$ . Writing  $x^3 = u^3 - iv^3 + i(u^2v + vu^2) + iuvu - vuv - (v^2u + uv^2)$  we see that the individual terms of the expansion lie in  $Z(\mathfrak{D})$ .

We turn to (e). Let  $\gamma$  be the natural homomorphism of  $A$  onto  $A/Z(\mathfrak{D})$ . For  $x \in R^*$ ,  $[\gamma(x)]^3 = 0$  by (d) so that  $\gamma(\mathfrak{R}^*) \subset W$ , the radical of  $A/Z(\mathfrak{D})$ . Inasmuch as  $A/R^*$  is semi-simple by Lemma 3.1, so is  $[A/Z(\mathfrak{D})]/[\mathfrak{R}^*/Z(\mathfrak{D})]$ . Therefore  $\mathfrak{R}^*/Z(\mathfrak{D}) \supset W$ .

**4.2. THEOREM.** *The following statements are equivalent.*

- (a) *There exists a faithful \*-representation of  $A$  continuous on  $H$ .*
- (b)  *$A$  is semi-simple and  $P \cap (-P) = (0)$ .*
- (c)  *$A$  is semi-simple and  $RI(\mathfrak{D}) = (0)$ .*

Suppose (a).  $A$  is semi-simple by [8, Theorem 4.1.19]. Lemma 3.2 gives  $\mathfrak{R}^* = (0)$  so that  $P \cap (-P) = (0)$  from Theorem 4.1 (b).

Suppose (b). Then  $Z(\mathfrak{D}) = (0)$  so that, by Theorem 4.1 (d),  $x^3 = 0$  for each  $x \in \mathfrak{R}^*$ . Since  $A$  is semi-simple,  $\mathfrak{R}^* = (0)$ . Then Theorem 4.1 (b) shows that  $RI(\mathfrak{D}) = (0)$ .

Suppose (c). Again  $Z(\mathfrak{D}) = (0)$  by Theorem 4.1 (b). As just seen this implies that  $\mathfrak{R}^* = (0)$  so that (a) follows from Lemma 3.2.

**4.3. COROLLARY.** *Let  $A$  be a normed  $*$ - $Q$ -algebra. Then  $A$  has faithful  $*$ -representation if and only if  $A$  is semi-simple and  $P \cap (-P) = (0)$ .*

This follows immediately from Theorem 4.2 and Lemma 2.4.

We now exhibit a normed  $*$ -algebra with a faithful  $*$ -representation but for which  $P \cap (-P) \neq (0)$ . Let  $A$  be the algebra of all polynomials in the complex variable  $z$ . For  $p(z) = \sum \alpha_n z^n$  set  $p^*(z) = \sum \bar{\alpha}_n z^n$ . First consider  $A$  in the norm

$$\|p\| = \sup_{0 \leq t \leq 1} |p(t)|.$$

Here, for each  $t$ ,  $0 \leq t \leq 1$  the functional  $f_t(p) = p(t)$  is a positive linear functional continuous on  $A$  and real-valued on  $H$ . Thus  $Z(\mathfrak{D}) = (0)$ . By Theorem 4.2 we see that  $A$  has a faithful  $*$ -representation. This also justifies a remark following Lemma 2.5.

Next consider  $A$  in the norm  $\|p\| = \sum |\alpha_k|/k!$  (see § 2). For  $p(z) = \alpha_0 + \cdots + \alpha_n z^n$  let  $f(p) = \alpha_0$ . This gives us a continuous  $*$ -representation of  $A$  as operators on one-dimensional Hilbert space with kernel  $M = \{p : p(0) = 0\}$  so that  $M \supset \mathfrak{R}^*$ . The arguments of § 2 following Lemma 2.5 show that any  $*$ -representation of  $A$  continuous on  $H$  must vanish on  $M$ . Therefore  $M = \mathfrak{R}^*$ . Via Theorem 4.1 we see that  $P \cap (-P)$  is the set of all polynomials with real coefficients vanishing at the origin. We investigate the commutative case more closely in § 5.

**4.4. LEMMA.**  $\mathfrak{R}^* = A \cap \mathfrak{R}_1^*$ .

We already have  $\mathfrak{R}^* \subset A \cap \mathfrak{R}_1^*$  by (4.1). Let  $\mathfrak{R} = A \cap \mathfrak{R}_1^*$ . By (4.2),  $h\mathfrak{R}h \subset Z(\mathfrak{D})$  for each  $h \in H$ . Reasoning exactly as in the proof of Theorem 4.1 (d) we obtain  $x^3 \in Z(\mathfrak{D})$  for each  $x \in \mathfrak{R}$ . Let  $\beta$  be the natural homomorphism of  $A$  onto  $A/\mathfrak{R}^*$ . Since  $Z(\mathfrak{D}) \subset \mathfrak{R}^*$  by Theorem 4.1 (b), we see that  $[\beta(x)]^3 = 0$  for each  $x \in \mathfrak{R}$ . From Lemma 3.1 we obtain  $\beta(\mathfrak{R}) = (0)$ . We now derive another formula for  $\mathfrak{R}^*$ .

**4.5. THEOREM.**  $\mathfrak{R}^* = Z(\mathfrak{E})$ .

As noted in the proof of Theorem 3.4,  $\mathfrak{E}$  is the set of positive

linear functionals on  $A$  which are extendable to positive linear functionals  $f$  on  $A_1$ , lying in  $\mathfrak{D}_1$ , with  $f(e) \leq 1$ . Now any  $g \in \mathfrak{D}_1$  is a multiple of such a functional. Therefore  $Z(\mathfrak{C}) = A \cap Z(\mathfrak{D}_1) = A \cap \mathfrak{R}_1^* = \mathfrak{R}^*$  by way of Theorem 4.1 and Lemma 4.4.

We have  $\mathfrak{R}^* = RI(\mathfrak{C}) = Z(\mathfrak{C})$ , a situation which differs from what can happen for  $\mathfrak{D}$ . In particular  $Z(\mathfrak{C}) \neq Z(\mathfrak{D})$  can occur.

5. The commutative case. Let  $A$  be a commutative algebra with an involution. By commutativity,  $H$  is a real subalgebra of  $A$ . We suppose in § 5 that  $H$  has a norm in terms of which it is a real normed algebra. Let  $\mathfrak{M}$  be the set of modular maximal ideals of  $A_1$ . We call  $M \in \mathfrak{M}$  *symmetric* if  $M = M^*$  and single out for special attention the set of symmetric  $M$  for which  $M \cap H$  is closed in  $H$ .

5.1. LEMMA. *Let  $\mu$  be a homomorphism of  $H$  into the reals. Define, for each  $x = h + ik$ ,  $h, k \in H$  the functional  $\mu_a$  by the rule  $\mu_a(x) = \mu(h) + i\mu(k)$ . Then  $\mu_a$  is a multiplicative (complex) linear functional on  $A$ .*

This can be verified in a straight forward way.

5.2. LEMMA. *Let  $M$  be a symmetric modular maximal ideal of  $A$  where  $M \cap H$  is closed in  $H$ . Then there exists a continuous homomorphism  $\mu$  of  $H$  onto the reals such that  $\mu_a^{-1}(0) = M$ .*

Let  $j$  be an identity for  $A$  modulo  $M$ . Then so is  $(j + j^*)/2$  so without loss of generality we can take  $j$  s.a. Then  $ju - u \in M \cap H$  for all  $u \in H$  and therefore  $M \cap H$  is a modular ideal of  $H$ . Since  $M = M \cap H \oplus i(M \cap H)$  it is clear the  $M \cap H \neq H$ . We claim that  $M \cap H$  is a modular maximal ideal of  $H$ . For otherwise there exists a modular maximal ideal  $K$  of  $H$  containing  $M \cap H$ ,  $K \neq M \cap H$ . An easy computation shows that  $K \oplus iK$  is an ideal of  $A$  containing  $M$ . Then  $K \oplus iK = A$  which is impossible as  $j \notin K$  (otherwise  $K = H$ ). Inasmuch as  $M \cap H$  is closed in  $H$ ,  $H/M \cap H$  is a normed field in the quotient algebra norm. By Mazur's theorem,  $H/M \cap H$  is a copy of the real or complex field. We rule out the latter possibility. If  $H/M \cap H$  were a copy of the complexes then it would be two-dimensional over the real field and there would be a two-dimensional real subspace  $L$  of  $H$  such that  $H = M \cap H \oplus L$ . Then  $A = H \oplus iH = M \oplus L \oplus iL$  which compels  $A/M$  to be four-dimensional over the reals. But surely  $A/M$  is a division algebra over the reals. Thus a well-known theorem of Frobenius makes  $A/M$  a copy of the quaternions. This is impossible in view of commutativity. Consequently there is a continuous homomorphism  $\mu$  of  $H$  onto the reals with kernel  $M \cap H$ .

5.3. THEOREM.  $\mathfrak{R}^*$  is the intersection of those symmetric modular maximal ideals  $M$  of  $A$  such that  $M \cap H$  is closed in  $H$ .

Take  $x = u + iv$ ,  $u, v \in H$ . By commutativity,  $x^*x = (u^2 + v^2)/2$ . Thus  $P$  is the closure in  $H$  of finite sums of squares of elements of  $H$ . Suppose first that  $A$  has an identity  $e$ . The proof of Theorem 3.4 shows that  $e \in \text{Int}(P)$ . Let  $\Sigma$  represent the set of all continuous real linear functionals  $g$  on  $H$  where  $g(P) \geq 0$  and  $g(e) \leq 1$ . The arguments of [5, Theorem 2.1] show that the set  $\Sigma_e$  of extreme points of  $\Sigma$  is the set the continuous homomorphisms of  $H$  into the reals. As in [5, Remark 2.3] it follows that  $P \cap (-P) = \bigcap f^{-1}(0)$  where  $f$  ranges over  $\Sigma_e$ . Let  $S$  be the intersection of the symmetric modular maximal ideals  $M$  of  $A$  with  $M \cap H$  closed in  $H$ . Lemmas 5.1 and 5.2 show that  $H \cap S = \bigcap f^{-1}(0) = P \cap (-P)$  and Lemma 4.1 (b) shows that  $S = \mathfrak{R}^*$ .

Now suppose that  $A$  has no identity. Each multiplicative linear functional on  $A$  which is real and continuous on  $H$  extends, as is easily verified, to a multiplicative linear functional on  $A_1$  which is real and continuous on  $H_1$ . Applying the result for the case with the identity we get  $S = A \cap \mathfrak{R}_1^* = \mathfrak{R}^*$  with the aid of Lemma 4.4.

6. An example. We give an example of a normed  $*$ -algebra  $A$  which has a continuous faithful  $*$ -representation and a continuous involution but for which the completion<sup>2</sup>  $A_c$  has no faithful  $*$ -representation. This demonstrates conclusively that our results in the case of a normed  $*$ -algebra (e.g. Theorem 4.2 and Corollary 4.3) cannot possibly be deduced from the theory of Banach algebras.

The algebra  $A$  which we use is a subalgebra of an algebra devised for other purposes by C. Feldman [2]. His algebra is the commutative algebra  $B$  which is the completion of the algebra of all finite sums

$$\sum_{i=1}^n \alpha_i e_i + \beta r$$

where  $\alpha_i$  and  $\beta$  are complex, the  $e_i$  are mutually orthogonal idempotents,  $r^2 = 0 = e_i r = r e_i$  for all  $i$  and

$$\|\Sigma \alpha_i e_i + \beta r\| = \max \{(\Sigma |\alpha_i|^2)^{1/2}, |\beta - \Sigma \alpha_i|\}.$$

Consider the subalgebra  $A$  consisting of all finite sums  $\Sigma \alpha_i e_i$ . The involution

$$(\Sigma \alpha_i e_i)^* = \Sigma \bar{\alpha}_i e_i$$

on  $A$  is an isometry. For each integer  $n > 1$  let  $s(n)$  be the smallest

---

<sup>2</sup>The involution on  $A$  extends to an involution on  $A_c$ .

integer of the form  $n + k$  such that  $\sum_{j=0}^k (n + j)^{-1} > 1$ . Let  $z_n = n^{-1}e_n + (n + 1)^{-1}e_{n+1} + \cdots + [s(n)]^{-1}e_{s(n)}$ . It is readily verified that  $\|r - z_n\| \rightarrow 0$ . Therefore  $B$  is the completion of  $A$ . For each  $n = 1, 2, \dots$  the functional  $f_n$  defined on  $A$  by the rule  $f_n(\sum \alpha_i e_i) = \alpha_n$  is a continuous multiplicative linear functional on  $A$ . Moreover  $\bigcap f_n^{-1}(0) = (0)$  so that  $A$  is semi-simple and, by Theorem 5.3 and Lemma 3.2,  $A$  has a faithful \*-representation continuous on  $H$ . The continuity of the involution allows us to assert that this \*-representation is continuous on  $A$ . However, the completion  $B$  of  $A$  is not semi-simple [2] and so has no faithful \*-representation [8, Theorem 4.1.19].

## REFERENCES

1. P. C. Curtis, Jr., *Order and commutativity in Banach algebras*, Proc. Amer. Math. Soc., **9** (1958), 643-646.
2. C. Feldman, *The Wedderburn principal theorem in Banach algebras*, Proc. Amer. Math. Soc., **2** (1951), 771-777.
3. I. Kaplansky, *Topological rings*, Amer. J. Math., **69** (1947), 153-183.
4. ———, *Topological algebra*, Dept. of Math. University of Chicago, 1952 (mimeographed notes).
5. J. L. Kelley and R. L. Vaught, *The positive cone in Banach algebras*, Trans. Amer. Math. Soc., **74** (1953), 44-55.
6. L. H. Loomis, *An introduction to abstract harmonic analysis*, D. Van Nostrand Co., New York 1953.
7. M. A. Naimark, *Involutive Algebren*, Sowjetische Arbeiten Zur Funktional-Analyse, Berlin (1954), 89-196.
8. C. E. Rickart, *General theory of Banach algebras*, D. Van Nostrand Co., New York, 1960.
9. B. Yood *Homomorphisms on normed algebras*, Pacific J. Math., **8** (1958), 373-381.
10. ———, *Faithful \*-representations of normed algebras*, Pacific J. Math., **10** (1960), 345-363.

UNIVERSITY OF OREGON



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

ROBERT OSSERMAN

Stanford University  
Stanford, California

M. G. ARSOVE

University of Washington  
Seattle 5, Washington

J. DUGUNDJI

University of Southern California  
Los Angeles 7, California

LOWELL J. PAIGE

University of California  
Los Angeles 24, California

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

\* \* \*

AMERICAN MATHEMATICAL SOCIETY  
CALIFORNIA RESEARCH CORPORATION  
SPACE TECHNOLOGY LABORATORIES  
NAVAL ORDNANCE TEST STATION

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and on submission, must be accompanied by a separate author's résumé. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal but they are not owners or publishers and have no responsibility for its content or policies.



# Pacific Journal of Mathematics

Vol. 14, No. 4

August, 1964

Homer Franklin Bechtell, Jr., <i>Pseudo-Frattini subgroups</i> .....	1129
Thomas Kelman Boehme and Andrew Michael Bruckner, <i>Functions with convex means</i> .....	1137
Lutz Bungart, <i>Boundary kernel functions for domains on complex manifolds</i> .....	1151
L. Carlitz, <i>Rings of arithmetic functions</i> .....	1165
D. S. Carter, <i>Uniqueness of a class of steady plane gravity flows</i> .....	1173
Richard Albert Dean and Robert Harvey Oehmke, <i>Idempotent semigroups with distributive right congruence lattices</i> .....	1187
Lester Eli Dubins and David Amiel Freedman, <i>Measurable sets of measures</i> .....	1211
Robert Pertsch Gilbert, <i>On class of elliptic partial differential equations in four variables</i> .....	1223
Harry Gonshor, <i>On abstract affine near-rings</i> .....	1237
Edward Everett Grace, <i>Cut points in totally non-semi-locally-connected continua</i> .....	1241
Edward Everett Grace, <i>On local properties and <math>G_\delta</math> sets</i> .....	1245
Keith A. Hardie, <i>A proof of the Nakaoka-Toda formula</i> .....	1249
Lowell A. Hinrichs, <i>Open ideals in <math>C(X)</math></i> .....	1255
John Rolfe Isbell, <i>Natural sums and abelianizing</i> .....	1265
G. W. Kimble, <i>A characterization of extremals for general multiple integral problems</i> .....	1283
Nand Kishore, <i>A representation of the Bernoulli number <math>B_n</math></i> .....	1297
Melven Robert Krom, <i>A decision procedure for a class of formulas of first order predicate calculus</i> .....	1305
Peter A. Lappan, <i>Identity and uniqueness theorems for automorphic functions</i> .....	1321
Lorraine Doris Lavalley, <i>Mosaics of metric continua and of quasi-Peano spaces</i> ....	1327
Mark Mahowald, <i>On the normal bundle of a manifold</i> .....	1335
J. D. McKnight, <i>Kleene quotient theorems</i> .....	1343
Charles Kimbrough Megibben, III, <i>On high subgroups</i> .....	1353
Philip Miles, <i>Derivations on <math>B^*</math> algebras</i> .....	1359
J. Marshall Osborn, <i>A generalization of power-associativity</i> .....	1367
Theodore G. Ostrom, <i>Nets with critical deficiency</i> .....	1381
Elvira Rapaport Strasser, <i>On the defining relations of a free product</i> .....	1389
K. Rogers, <i>A note on orthogonal Latin squares</i> .....	1395
P. P. Saworotnow, <i>On continuity of multiplication in a complemented algebra</i> .....	1399
Johanan Schonheim, <i>On coverings</i> .....	1405
Victor Lenard Shapiro, <i>Bounded generalized analytic functions on the torus</i> .....	1413
James D. Stafney, <i>Arens multiplication and convolution</i> .....	1423
Daniel Sterling, <i>Coverings of algebraic groups and Lie algebras of classical type</i> .....	1449
Alfred B. Willcox, <i>Šilov type <math>C</math> algebras over a connected locally compact abelian group. II</i> .....	1463
Bertram Yood, <i>Faithful <math>*</math>-representations of normed algebras. II</i> .....	1475
Alexander Zabrodsky, <i>Covering spaces of paracompact spaces</i> .....	1489