

Pacific Journal of Mathematics

ON A CONJECTURE OF R. J. KOCH

L. E. WARD

ON A CONJECTURE OF R. J. KOCH

L. E. WARD, JR.¹

*Dedicated to Professor Alexander Doniphan Wallace
 on the occasion of his sixtieth birthday*

R. J. Koch proved that if X is a compact, continuously partially ordered space and if W is an open subset of X which has no local minima, then each point of W is the supremum of an order arc which meets $X - W$. More recently he extended this result to quasi ordered spaces in which the sets $E(x) = \{y: x \leq y \leq x\}$ are assumed to be totally disconnected and W is a chain. He conjectured that the latter hypothesis is superfluous, and we show here that Koch's conjecture is correct.

As a corollary it follows that if X is a compact, continuously quasi ordered space with zero (i.e., a unique minimal element), if each set $E(x)$ is totally disconnected, and if each set $L(x) = \{y: y \leq x\}$ is connected, then X is arcwise connected.

We begin by recalling a few definitions (see [1], [2], [3] and [4]). We say that $X = (X, \Gamma)$ is a continuously quasi ordered space provided X is a Hausdorff space, Γ is a quasi order (= reflexive, transitive relation) on X and the graph of Γ is a closed subset of $X \times X$. We identify Γ with its graph and regard the symbols $x \leq y$, and $x \Gamma y$ and $(x, y) \in \Gamma$ as synonyms.

A *chain* of a quasi ordered space X is a subset C of X such that $a \leq b$ or $b \leq a$ holds for each a and b in C . We also define

$$\begin{aligned} L(a, \Gamma) &= \{x \in X: (x, a) \in \Gamma\}, \\ M(a, \Gamma) &= \{x \in X: (a, x) \in \Gamma\}, \\ E(a, \Gamma) &= L(a, \Gamma) \cap M(a, \Gamma), \end{aligned}$$

for each $a \in X$. It is also convenient to define

$$I(a, b, \Gamma) = M(a, \Gamma) \cap L(b, \Gamma),$$

the closed "interval" from a to b . Where there is no ambiguity we shall write $(L(a)$ (resp., $M(a)$, $E(a)$, $I(a, b)$) for $L(a, \Gamma)$, (resp., $M(a, \Gamma)$, $E(a, \Gamma)$, $I(a, b, \Gamma)$). It is well known [3] that if X is a continuously quasi ordered space then the sets $L(a)$, $M(a)$, $E(a)$ and $I(a, b)$ are closed and, if X is compact, then X contains a minimal element, that is, an element m such that $L(m) - E(m)$ is empty.

¹ Received September 2, 1964. Presented to the American Mathematical Society, November 14, 1964. This research was supported by a grant from the National Science Foundation.

A subset Y of the quasi ordered space (X, Γ) is said to have no local Γ -minima if, for each $x \in Y$ and each neighborhood U of x , the set

$$Y \cap U \cap L(x, \Gamma) - E(x, \Gamma)$$

is nonempty. This definition is due to Koch [2].

In case the relation Γ is a partial order, it is known that a connected chain joining two distinct points is an arc. (Here we use the term *arc* to describe a continuum with precisely two non-cutpoints.) An arc which is also a chain is termed an *order arc*.

The following two lemmas will be of later use.

LEMMA 1. *Let X be a compact, continuously quasi ordered space, let a and b be members of X , and let K be a closed subset of X such that $I(a, b) \cap K = 0$. Then there exist open sets U and V such that $a \in U$, $b \in V$ and for each $a' \in U$ and $b' \in V$ it follows that $I(a', b') \cap K = 0$.*

Proof. Suppose, on the contrary, that for all neighborhoods U and V of a and b , respectively, there exists $a' \in U$ and $b' \in V$ such that $I(a' b') \cap K \neq 0$. Then

$$\Gamma \cap (\bar{U} \times K) \cap (K \times \bar{V}) \neq 0.$$

These sets form a family of nonempty closed sets with the finite intersection property and hence their intersection is nonempty:

$$\Gamma \cap (\{a\} \times K) \cap (K \times \{b\}) \neq 0,$$

that is to say, $I(a, b) \cap K \neq 0$, contrary to the hypothesis.

LEMMA 2. *If R is an open subset of the compact, continuously quasi ordered space X , then the set*

$$F = \{(a, b) \in X \times X: I(a, b) - R \neq 0\}$$

is closed.

Proof. If $(a, b) \notin F$ then $I(a, b) \cap (X - R) = 0$. By Lemma 1, there are open sets U and V with $a \in U$ and $b \in V$ such that for each $a' \in U$ and $b' \in V$ it follows that $I(a', b') \subset R$, and hence $(U \times V) \cap F = 0$. Therefore, F is closed.

2. Koch's theorem for quasi ordered spaces. The crux of our proof is embodied in the following theorem.

THEOREM. *Let $X = (X, \Gamma)$ be a compact, continuously quasi*

ordered space and let W be an open subset of X . If

(i) $E(x, \Gamma)$ is totally disconnected for each $x \in X$,

(ii) W has no local Γ -minima, then X admits a minimal quasi order which has a closed graph and satisfies (i) and (ii). Moreover, this minimal quasi order is a partial order.

Proof. Let $\{\Gamma_\alpha\}$ be a maximal nest of quasi orders on X such that each Γ_α has a closed graph and satisfies (i) and (ii), and let $\Gamma = \bigcap \{\Gamma_\alpha\}$. Clearly (X, Γ) is a continuously quasi ordered space and $E(x, \Gamma)$ is totally disconnected. We will show that W has no local Γ -minima.

Let $x \in W$ and let U be a neighborhood of x ; since W is open and $E(x, \Gamma)$ is totally disconnected, we may assume that $U \subset W$ and that $E(x, \Gamma) \cap U$ is closed. Since X is normal there exist open sets V and R such that

$$\begin{aligned} E(x, \Gamma) \cap U &\subset V \subset \bar{V} \subset U, \\ X - U &\subset R \subset \bar{R} \subset X - \bar{V}. \end{aligned}$$

For each α , the compact set $L(x, \Gamma_\alpha) \cap \bar{V}$ has a Γ_α -minimal element which we denote x_α . And since W has no local Γ_α -minima there exists

$$y_\alpha \in (X - \bar{R}) \cap L(x_\alpha, \Gamma_\alpha) - E(x_\alpha, \Gamma_\alpha).$$

It follows that

$$y_\alpha \in L(x, \Gamma_\alpha) - \bar{R} \cup \bar{V}$$

so that the sets $L(x, \Gamma_\alpha) - \bar{R} \cup \bar{V}$ are compact, nonempty and nested. Consequently there exists

$$y \in L(x, \Gamma) - \bar{R} \cup \bar{V}$$

and it is clear that $y \notin E(x, \Gamma)$. That is, W has no local Γ -minima.

Now suppose that Γ is not a partial order; then there exists a nondegenerate set $E(x, \Gamma)$. Since $E(x, \Gamma)$ is compact and totally disconnected, there exist nonempty, closed and disjoint sets A and B whose union is $E(x, \Gamma)$. Since X is normal there exist disjoint open sets P and Q such that $A \subset P$ and $B \subset Q$. Let

$$F = \{(a, b) : I(a, b) - P \cup Q \neq \emptyset\}.$$

By Lemma 2, F is a closed subset of $X \times X$ and hence

$$\Delta = \Gamma - ((P \times Q) - F)$$

is also closed. Since P and Q are disjoint, Δ is a reflexive relation on X .

We claim that Δ is a quasi order. For suppose $p \Delta q$ and $q \Delta r$ but $(p, r) \in (X \times X) - \Delta$. Now $(p, r) \in \Gamma$ so that $(p, r) \in (P \times Q) - F$ and hence $q \in P$ or $q \in Q$. If $q \in P$ then, since $r \in Q$ and $(q, r) \in \Delta$ we infer that $(q, r) \in F$ and thus $I(q, r) - P \cup Q \neq \emptyset$. But $I(q, r) \subset I(p, r)$ and hence $I(p, r) - P \cup Q \neq \emptyset$, contrary to the fact that $(p, r) \in (P \times Q) - F$. A similar contradiction ensues if $q \in Q$, and thus Δ is a quasi order.

Since $\Delta \subset \Gamma$ it is obvious that each set $E(x, \Delta)$ is totally disconnected. Now suppose $z \in W$ and that O is a neighborhood of z , $O \subset W$. If $z \in W - Q$ then

$$L(z, \Delta) = L(z, \Gamma)$$

and hence there exists

$$y \in O \cap L(z, \Delta) - E(z, \Delta).$$

And if $z \in Q$, the fact that W has no local Γ -minima insures the existence of

$$y \in O \cap Q \cap L(z, \Gamma) - E(z, \Gamma).$$

But $y \notin P$ implies $y \in L(z, \Delta)$, so that in any event W has no local Δ -minima.

Finally we note that Δ contradicts the minimality of Γ , for if $a \in A$ and $b \in B$ then $(a, b) \in \Gamma - \Delta$. Therefore Γ is a partial order.

COROLLARY 1. *Let X be a compact, continuously quasi ordered space and let W be an open subset of X . If conditions (i) and (ii) of the theorem are satisfied, then each point of W is the supremum of an order arc which meets $X - W$.*

Proof. By the preceding theorem we may assume that the quasi order is a partial order. Thus Koch's theorem for partially ordered spaces applies.

An element 0 of the quasi ordered space X is a *zero* of X provided

$$0 = E(0) = \cap \{L(x) : x \in X\}.$$

COROLLARY 2. *If X is a compact, continuously quasi ordered space with zero, if each set $E(x)$ is totally disconnected and if each set $L(x)$ is connected, then X is arcwise connected.*

Proof. Let $W = X - \{0\}$; the connectedness of the sets $L(x)$ guarantees that W has no local minima and therefore each point of W lies in arc containing 0 .

Following Koch we say that a subset C of the quasi ordered space X is *biconnected* if C is connected and if each of the sets $E(x) \cap C$ is

connected.

COROLLARY 3. *Let X be a compact, continuously quasi ordered space and suppose there exists $a \in X$ such that*

$$E(a) = \bigcap \{L(x) : x \in X\} .$$

If $X - E(a)$ has no local minima then each element of X can be joined to $E(a)$ by a biconnected chain.

Proof. Let Z denote the compact, continuously partially ordered space which is obtained when $E(x)$ is identified with a point, for each $x \in X$. Let $\phi(X) = Z$ be the canonical quotient map and let

$$X \xrightarrow{m} Y \xrightarrow{l} Z$$

be the monotone-light factorization of ϕ . It is easy to see that Y inherits a quasi order from Z which has a closed graph and is such that $E(y)$ is totally disconnected, for each $y \in Y$. Moreover, $Y - m(E(a))$ has no local minima and hence, by the theorem, there are order arcs joining points of Y to $m(E(a))$. Since m is monotone, the corollary follows at once.

REFERENCES

1. R. J. Koch, *Arcs in partially ordered spaces*, Pacific J. Math. **9** (1959), 723-728.
2. ———, *Connected chains in quasi ordered spaces*, Fund. Math. **56** (1965), 245-249.
3. L. E. Ward, Jr., *Partially ordered topological spaces*, Proc. Amer. Math. Soc. **5** (1954), 144-161.
4. ———, *Concerning Koch's theorem on the existence of arcs*, Pacific J. Math. **15** (1965), 347-355.

UNIVERSITY OF OREGON

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California

R. M. BLUMENTHAL

University of Washington
Seattle, Washington 98105

J. DUGUNDJI

University of Southern California
Los Angeles, California 90007

*RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. No separate author's resumé is required. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

* Basil Gordon, Acting Managing Editor until February 1, 1966.

Robert James Blattner, <i>Group extension representations and the structure space</i>	1101
Glen Eugene Bredon, <i>On the continuous image of a singular chain complex</i>	1115
David Hilding Carlson, <i>On real eigenvalues of complex matrices</i>	1119
Hsin Chu, <i>Fixed points in a transformation group</i>	1131
Howard Benton Curtis, Jr., <i>The uniformizing function for certain simply connected Riemann surfaces</i>	1137
George Wesley Day, <i>Free complete extensions of Boolean algebras</i>	1145
Edward George Effros, <i>The Borel space of von Neumann algebras on a separable Hilbert space</i>	1153
Michel Mendès France, <i>A set of nonnormal numbers</i>	1165
Jack L. Goldberg, <i>Polynomials orthogonal over a denumerable set</i>	1171
Frederick Paul Greenleaf, <i>Norm decreasing homomorphisms of group algebras</i>	1187
Fletcher Gross, <i>The 2-length of a finite solvable group</i>	1221
Kenneth Myron Hoffman and Arlan Bruce Ramsay, <i>Algebras of bounded sequences</i>	1239
James Patrick Jans, <i>Some aspects of torsion</i>	1249
Laura Ketchum Kodama, <i>Boundary measures of analytic differentials and uniform approximation on a Riemann surface</i>	1261
Alan G. Konheim and Benjamin Weiss, <i>Functions which operate on characteristic functions</i>	1279
Ronald John Larsen, <i>Almost invariant measures</i>	1295
You-Feng Lin, <i>Generalized character semigroups: The Schwarz decomposition</i>	1307
Justin Thomas Lloyd, <i>Representations of lattice-ordered groups having a basis</i>	1313
Thomas Graham McLaughlin, <i>On relative coimmunity</i>	1319
Mitsuru Nakai, <i>Φ-bounded harmonic functions and classification of Riemann surfaces</i>	1329
L. G. Nova, <i>On n-ordered sets and order completeness</i>	1337
Fredos Papan gelou, <i>Some considerations on convergence in abelian lattice-groups</i>	1347
Frank Albert Raymond, <i>Some remarks on the coefficients used in the theory of homology manifolds</i>	1365
John R. Ringrose, <i>On sub-algebras of a C^*-algebra</i>	1377
Jack Max Robertson, <i>Some topological properties of certain spaces of differentiable homeomorphisms of disks and spheres</i>	1383
Zalman Rubinstein, <i>Some results in the location of zeros of polynomials</i>	1391
Arthur Argyle Sagle, <i>On simple algebras obtained from homogeneous general Lie triple systems</i>	1397
Hans Samelson, <i>On small maps of manifolds</i>	1401
Annette Sinclair, <i>$\varepsilon(z)$-closeness of approximation</i>	1405
Edsel Ford Stiel, <i>Isometric immersions of manifolds of nonnegative constant sectional curvature</i>	1415
Earl J. Taft, <i>Invariant splitting in Jordan and alternative algebras</i>	1421
L. E. Ward, <i>On a conjecture of R. J. Koch</i>	1429
Neil Marchand Wigley, <i>Development of the mapping function at a corner</i>	1435
Horace C. Wiser, <i>Embedding a circle of trees in the plane</i>	1463
Adil Mohamed Yaqub, <i>Ring-logics and residue class rings</i>	1465
John W. Lamperti and Patrick Colonel Suppes, <i>Correction to: Chains of infinite order and their application to learning theory</i>	1471
Charles Vernon Coffman, <i>Correction to: Non-linear differential equations on cones in Banach spaces</i>	1472
P. H. Doyle, III, <i>Correction to: A sufficient condition that an arc in S^n be cellular</i>	1474
P. P. Saworotnow, <i>Correction to: On continuity of multiplication in a complemented algebra</i>	1474
Basil Gordon, <i>Correction to: A generalization of the coset decomposition of a finite group</i>	1474