

Pacific Journal of Mathematics

FAMILIES OF PARALLELS ASSOCIATED WITH SETS

EUGENE ELLIOT ROBKIN AND F. A. VALENTINE

FAMILIES OF PARALLELS ASSOCIATED WITH SETS

E. E. ROBKIN and F. A. VALENTINE

There exist sets S in Euclidean space E_n which have an interesting association with a family \mathcal{P} of parallel lines. For instance S and \mathcal{P} may be so related that each point of S lies on a member of \mathcal{P} which intersects S in either a line segment or a point. There exist compact sets $S \subset E_2$ such that every finite collection of points in S is contained in some collection of parallel lines each of which intersects S in a single point, and yet no infinite family \mathcal{P} of parallel lines exists having the same property and covering S . This paper contains a theorem which enables one to determine the existence of a family of parallel lines each of which intersects S in a line segment or point and which as a family covers S .

Secondly we show that the points, the closed line segments, the closed convex triangular regions, and the closed convex sets bounded by parallelograms are the only compact convex sets B in E_2 which have the following property. If A is a closed connected set disjoint from B and if every 3 or fewer points of A lie on parallel lines intersecting B , then A is covered by a family of parallel lines each of which intersects B .

Finally, we obtain a theorem of Krasnoselskii type. Intuitively, this may be stated as follows. Suppose S is a compact set in E_n and suppose there exists a plane H such that every n points of S can see H via S along parallel lines. Then all the points of S can see H via S along parallel lines.

The above results appear in Theorems 3, 2, 1 in that order. The appendix at the end contains the theorems of Helly, Krasnosel'skii and other results used. Furthermore, the reader is recommended to consult the compendium "Helly's theorem and its relatives" by Danzer, Klee and Grünbaum [1]. In order to proceed logically we adopt the following notations.

NOTATION. If S is a set in n -dimensional Euclidean space E_n , then closure of $S = \text{cl } S$, interior of $S = \text{int } S$, boundary of $S = \text{bd } S$, convex hull of $S = \text{conv } S$. If $x \in E_n, y \in E_n, x \neq y$, then $L(x, y) =$ line containing x and y , $xy =$ closed segment joining x and y , $\text{intv } xy =$ relative interior of the segment xy , $R(x, y) =$ ray having x as endpoint and containing y . The empty set is indicated by \emptyset and the origin of E_n by \mathcal{O} . Set union, intersection and difference are denoted by \cup, \cap and \sim respectively.

Received July 6, 1964. The preparation of this paper was sponsored (in part) by the National Science Foundation Grants GP 1368 and 1988.

Parallels.

THEOREM 1. *Let S be a closed set in n -dimensional Euclidean space E_n .*

(a) *Suppose there exists a hyperplane H such that $S \cap H$ is compact.*

(b) *Suppose for each integer s such that $1 \leq s \leq n$ and for each set of distinct points x_1, x_2, \dots, x_s in $S \sim H$ there exist points y_1, y_2, \dots, y_s in H such that $x_i y_i \subset S$ ($i = 1, \dots, s$) and such that $x_1 y_1, x_2 y_2, \dots, x_s y_s$ are parallel. (The y_1, \dots, y_s need not be distinct.)*

Then there exists a family \mathcal{P} of parallel lines such that for each point $x \in S \sim H$ there exists a point $y \in H$ such that $xy \subset S$ and such that the line $L(x, y)$ belongs to \mathcal{P} .

Proof. To each $x \in S \sim H$, let $C(x)$ denote the union of all lines $L(x, y)$ where $y \in H$ such that $xy \subset S$. Choose a point \emptyset as origin in E_n with $\emptyset \notin H$, and let $D(x)$ be that translate of $C(x)$ so that x goes to \emptyset . Define $M(x)$ as follows, when $x \in S \sim H$,

$$M(x) \equiv \text{conv}(H \cap D(x)).$$


Since $H \cap D(x) \neq \emptyset$, we must have $M(x) \neq \emptyset$ for each $x \in S \sim H$. Hypothesis (b) implies that every n or fewer members of the collection $\{M(x), x \in S \sim H\}$ have a point in common. Since the dimension of H is $n - 1$, and since the members of $\{M(x), x \in S \sim H\}$ are compact convex sets in H , Helly's theorem [2] (see Appendix) for $(n - 1)$ -dimensional space implies

$$(1) \quad \bigcap_{x \in S \sim H} M(x) \neq \emptyset.$$

Since $\emptyset \notin H$, condition (1) implies there exists a line L through \emptyset such that $L \cap M(x) \neq \emptyset$ for each $x \in S \sim H$. We let \mathcal{P} denote the family of all lines parallel to L and intersecting S . We will prove that \mathcal{P} has the desired property stated in the theorem. To do this choose a point $x \in S \sim H$, and let $L(x)$ be the line through x parallel to L . Let $L(x) \cap H \equiv y$ and suppose $xy \not\subset S$. Since S is closed then there exists a point $u \in \text{int } xy$ and a closed solid sphere $U(u, r)$ with center u and radius r such that $S \cap U(u, r) = \emptyset$. Let $U(v, t)$ be a closed solid sphere with center $v \in xu$ and radius t with $t < r$. Define $K(v)$ as follows,

$$K(v) \equiv \text{conv}[U(u, r) \cup U(v, t)], v \in xu.$$

Since S is closed, and since $K(u) \cap S = \emptyset$, $K(x) \cap S \neq \emptyset$, there exists a point $w \in xu$ nearest to u such that $S \cap \text{bd } K(w) \neq \emptyset$, $S \cap \text{int } K(w) = \emptyset$. Choose a point z such that



$$z \in S \cap bd K(w) \neq \emptyset .$$

Clearly $D(z)$ contains no line $L(z, y_1)$ parallel to the line $L(x)$ with $y_1 \in M(z)$. This contradicts the statement following (1). Hence $xy \subset S$, and Theorem 1 has been proved.

In order to express the next theorem easily the following concepts are used.

DEFINITION 1. A set A has the parallel property $P(m)$ relative to a set B if every m or fewer points of A all lie on a family of parallel lines each of which intersects B .

A set A has the parallel property $P(A)$ relative to B if all the points of A lie on a family \mathcal{S} of parallel lines each of which intersects B .

THEOREM 2. Let B be a compact convex set in the Euclidean plane E_2 . Suppose that each closed connected set A in E_2 which is disjoint from B and which has property $P(3)$ relative to B also has the property $P(A)$ relative to B .

Then B is either a point, a closed line segment, a closed set bounded by a triangle or a closed set bounded by a parallelogram.

Proof. Since this theorem characterizes the sets B , the proof consists of two parts.

Let $\mathcal{A} = \{A\}$ denote the collection of all those closed connected sets A which are disjoint from B and which have property $P(3)$ relative to B . First, suppose B is either a point, a closed line segment, a closed triangular region or a closed region bounded by a parallelogram.

Case 1. Suppose $bd B$ is a parallelogram with consecutive vertices a_1, a_2, a_3, a_4 . The four lines determined by the four edges of B divide

the plane E_2 into nine parts. Let $V(a_i)$ ($i = 1, 2, 3, 4$) denote the unbounded open V -shaped region abutting B at a_i ($i = 1, 2, 3, 4$), and let $V(a_1, a_2)$ be the closed unbounded region abutting B along $a_1 a_2$, etc. Since $A \in \mathcal{A}$ is connected and disjoint from B , if $A \cap V(a_i) \neq \emptyset$, $A \cap V(a_{i+1}) \neq \emptyset$, ($i = 1, 2, 3, 4$; $a_5 = a_1$), the set A would not have the property $P(2)$ relative to B . Hence, it would not have property $P(3)$ relative to B . Therefore, we may relabel the vertices of B so that

$$(2) \quad A \subset V(a_1, a_2) \cup V(a_2) \cup V(a_2, a_3).$$

For $x \in A$, let $C(x)$ denote the union of all rays emanating from x which intersect B , and let $D(x)$ be that translate of $C(x)$ which sends x to the origin \emptyset of E_2 . Since B is compact and convex, and since $x \notin B$, the set $C(x)$, and hence $D(x)$, is a closed convex cone which is not all of E_2 . Let C be the unit circle with center at \emptyset so that $C \equiv [x : \|x\| = 1]$.

Define $M(x)$ as follows,

$$C \cap D(x) \equiv M(x), \quad x \in A.$$

Consider the collection of sets

$$\mathcal{M} \equiv \{M(x), x \in A\}.$$

Property $P(3)$ implies that there exists a semicircular arc C_1 of C such that every two members of \mathcal{M} have a non-empty connected intersection with C_1 . To see this observe that if in (2) we have $A \cap V(a_2) \neq \emptyset$, then for each point $x \in A \cap V(a_2)$ we have a connected intersection

$$M(x) \cap M(x_1) \cap M(x_2) \neq \emptyset$$

for every pair of points x_1, x_2 in A . If $A \cap V(a_2) = \emptyset$ then either $A \subset V(a_1, a_2)$ or $A \subset V(a_2, a_3)$, and the above italicized statement is also still true. (It is instructive to observe that condition $P(2)$ does not suffice to imply the above italicized sentence.) We may now apply Helly's theorem [2] (see Appendix) to the set \mathcal{M} to yield the existence of a point $u \in M(x)$ for all $M \in \mathcal{M}$. Let L be the line determined by \emptyset and u . For each $x \in A$, let $L(x)$ denote the line through x parallel to L . The above facts imply that $L(x) \cap B \neq \emptyset$, $x \in A$, by virtue of the definition of $C(x)$. Hence A has property $P(A)$ relative to B .

Case 2. Suppose B is a closed set bounded by a triangle with vertices a_1, a_2, a_3 . As argued in case 1, we may relabel the vertices so that

$$A \subset V(a_1) \cup V(a_1, a_2)$$

if $A \in \mathcal{A}$, where $V(a_1)$ is the open V -shaped region abutting B at a_1 , and where $V(a_1, a_2)$ similarly abuts B along a_1a_2 . The rest of the proof is exactly the same as Case 1.

Case 3. Suppose B is a closed segment a_1a_2 . If $A \in \mathcal{A}$, then A either lies on $L(a_1, a_2)$ or in one of the open half-spaces bounded by $L(a_1, a_2)$. The proof is either trivial or exactly the same as Case 1, or, for that matter, as Theorem 1.

Case 4. If B is a point, then A must lie on a line through B , and the conclusion is trivial.

This completes the first part of the proof.

(I) *Secondly, to complete the characterization, suppose B is a compact convex set which is neither a point, a line segment, a triangular region or a set bounded by a parallelogram.*

We will prove that for such a set B there exists a closed connected set A , disjoint from B , which has property $P(3)$ relative to B , and which does not have property $P(A)$ relative to B . In order to construct A we use the familiar concept of "exposed point."

DEFINITION 2. A point x in the boundary of a convex set $S \subset E_2$ is an exposed point of S if there exists a line L of support to S at x such that $S \cap L = x$.

To construct A , let x_1x_2 be a diameter of the set B described in the italicized statement (I). The points x_1, x_2 are exposed points of B since the line L_i through x_i ($i = 1, 2$) and perpendicular to x_1x_2 is such that $L_i \cap B = x_i$ ($i = 1, 2$). Let H be one of the open half-planes bounded by $L(x_1, x_2)$ such that $H \cap B \neq \emptyset$, since $B \not\subset L(x_1, x_2)$. It is a well-known elementary fact that B contains at least one exposed point in H . If B contains one and only one exposed point in each of the open half-planes H and $E_2 \sim \text{cl } H$, then B is a *quadrilateral*. (We have excluded the case of a parallelogram, here.) On the other hand, if B contains only one exposed point in H and none in $E_2 \sim \text{cl } H$ then B is bounded by a *triangle*, which is also excluded here. *For the moment*, suppose A contains at least two *exposed* points x_3 and x_4 in H . Without loss of generality, suppose x_1, x_2, x_4, x_3 occur on $\text{bd } B$ in that order as illustrated in Figure 2. (In Fig. 2, the dotted lines and curves, except for the points x_1, x_2, x_3, x_4 miss B .)

Let $L(x_i)$ ($i = 3, 4$) be two lines such that $B \cap L(x_i) \equiv x_i$. Observe that

$$\text{conv}(x_1 \cup x_3 \cup x_4 \cup x_2) \subset B.$$

There exist points x_{ij} ($i, j = 1, 2, 3, 4, i < j$) such that




Fig. 2.

$$x_{ij} \equiv L(x_i) \cap L(x_j) \quad (i < j, i, j = 1, 2, 3, 4, (i, j) \neq (1, 2)).$$

To construct the set A , illustrated in Figure 2, extend the segment x_3x_{13} to x_3y_{13} and x_3x_{24} to x_3y_{24} so that

$$\begin{aligned} L(x_2, y_{24}) \cap L(x_1, x_3) \cap H &\neq \emptyset \\ L(x_1, y_{13}) \cap L(x_2, x_4) \cap H &\neq \emptyset. \end{aligned}$$

Recall that $x_3 \in H, x_4 \in H$. Furthermore, replace a segment ab of $y_{13}x_{34}$ with midpoint x_3 by a semicircular arc C_3 with endpoints a and b and with $C_3 \cap B = \emptyset$. Introduce a corresponding arc C_4 at x_4 (see Fig. 2). The set A defined as follows

$$A \equiv (y_{13}a) \cup C_3 \cup (bx_{34}) \cup (x_{34}c) \cup C_4 \cup (dy_{24})$$

is illustrated in Figure 2. We may choose the arcs C_3 and C_4 sufficiently small so that A clearly has property $P(3)$. To see this observe first that $A \sim C_4$ has property $P(2)$. Hence, to see that A has property $P(3)$ one merely has to demonstrate that if $x \in C_4, y \in A, z \in A$, the triple $\{x, y, z\}$ has property $P(3)$ relative to B . However, clearly the set A does not have property $P(A)$ relative to B . This completes the proof when B in statement (I) has two exposed points in H . The only case remaining in this part of the proof is that in which B is a *quadrilateral* which is *not* a parallelogram. So to complete the proof suppose B is such a quadrilateral. In this case there must exist some two vertices of B , say x_1 and x_2 , such that the other two vertices x_3 and

x_4 of B are interior to a strip bounded by two parallel line $L(x_1)$ and $L(x_2)$ at x_1 and x_2 respectively such that x_3 and x_4 lie on the same side of $L(x_1, x_2)$. Hence, we have a situation which is essentially the same as in Figure 2 (perpendicularity was not essential), and the same construction can be carried out to yield a set A having property $P(3)$ relative to B but not $P(A)$. This completes the proof.

There exist further results related to property $P(m)$ and these will be presented in a subsequent paper. It should be mentioned that if in the hypothesis of Theorem 2 we replace $P(3)$ by $P(2)$ then B must be either a point or a line segment. Also it is easy to prove that if B is a compact strictly convex body in E_2 and if m is a prescribed integer, there exists a compact connected set A which is disjoint from B , which has property $P(m)$ relative to B but which does not have property $P(A)$ relative to B .

THEOREM 3. *Let S be a closed connected set in the Euclidean plane E_2 . Suppose there exist two points a and b in S such that the following holds. If x_1 and x_2 are points in S then there exist some two parallel lines, denoted by L_1 and L_2 , such that $L_i \cap S = x_i$ and such that $L_i \cap ab \neq \emptyset$ ($i = 1, 2$).*

Then there exists a family \mathcal{S} of parallel lines in E_2 such that each point of S is contained in a member of \mathcal{S} which intersects S in either a line segment or a point.

Proof. If $x \in S$, by hypothesis there exists a line $L(x)$ through x such that

$$(4) \quad S \cap L(x) = x, ab \cap L(x) \neq \emptyset.$$

For $x \in S$, let $C(x)$ denote the union of all possible lines $L(x)$ satisfying (4). We will prove first that $C(x)$ is a two-napped cone, each nappe of which is convex, although it need not be closed. To prove this, we consider two cases.

First, suppose $x \notin ab$. Suppose $L_1(x), L_2(x)$ are two lines in $C(x)$ through x . Choose an arbitrary line $L(x)$ through x such that $L(x)$ intersects ab between $ab \cap L_1(x)$ and $ab \cap L_2(x)$. We will show that $L(x) \cap S = x$, so that $L(x) \subset C(x)$. The proof is indirect. Suppose a point y exists such that $y \in S \cap L(x)$, $y \neq x$. By hypothesis, there exists a line $L(y)$ through y such that $S \cap L(y) = y$, $ab \cap L(y) \neq \emptyset$. (See Fig. 3. In this figure, the dotted lines, except for the points x and y , miss the set S .) Since $L(y) \cap \text{intv } ab \neq \emptyset$, $L(x) \cap \text{intv } ab \neq \emptyset$, it is a simple matter to verify that the deletion from $E_2 \sim S$ of an appropriate ray of $L(y)$ together with an appropriate ray from $L_1(x)$ or $L_2(x)$ separates the plane into two disjoint open parts, one of which




Fig. 3.

contains a and the other of which contains b . However, this violates the fact that S is *connected*. Hence, we have a contradiction so that $S \cap L(x) = x$. Hence, when $x \notin ab$, the set $C(x)$ is a two-napped cone, each nappe of which is convex.

Secondly, suppose $x \in S \cap ab$. The proof follows the same pattern as in the case $x \notin ab$, (see Fig. 3). The obvious details are made self-evident there.

To complete the proof, choose an origin \emptyset in E_2 so that $\emptyset \notin L(a, b)$. Let $D(x)$ be that translate of $C(x)$ so that x goes to \emptyset . Consider the collection \mathcal{M} , defined as follows

$$\mathcal{M} \equiv \{M(x) \equiv L(a, b) \cap \text{cl } D(x), x \in S\}.$$

We have shown that every two members of \mathcal{M} have at least one point in common. Furthermore, if $x \notin L(a, b)$, then $M(x)$ is a compact interval. If $x \in ab$ the set $M(x)$ may be *unbounded* (although closed and convex), however, this will cause no difficulty because if $S \subset ab$, then $\bigcap_{x \in S} M(x) \neq \emptyset$ follows trivially, and if $S \not\subset ab$ Helly's theorem [2] can be used to yield the existence of a point u such that

$$u \in \bigcap_{x \in S} M(x).$$

Let L be the line determined by the two points u and \emptyset , since $\emptyset \neq u$. Let \mathcal{S} denote the set of all lines in E_2 which are parallel to L and which intersect S . We will prove that \mathcal{S} is a family as described in Theorem 3, and the proof is indirect. To do this, for $x \in S$ let $L(x)$ denote that line through x which belongs to \mathcal{S} so that $L(x)$ and L

are parallel. Suppose a point $x \in S$ exists such that $S \cap L(x)$ is not connected. Since S is closed if $S \cap L(x)$ is *not connected* there exist points c and d in $L(x)$ such that $c \in S, d \in S, c \neq d, S \cap \text{intv } cd = \emptyset$. The segment cd is usually called a cross-cut of the complement of S . Since S is *connected*, there exists a component K of the complement of S such that the removal of $\text{intv } cd$ from K yields two disjoint parts of K , at least one of which is bounded (see Fig. 4) which we denote by K_1 .




Fig. 4.

There exist points $y \in \text{bd } K_1$ with $y \notin L(c, d)$, sufficiently close to $L(x)$ such that the $L(y) \in \mathcal{P}$ parallel to $L(x)$ intersects K_1 and also intersects $\text{bd } K$ in a nonconnected set. Since $\text{cl } C(y)$ is the closure of a nonempty two-napped cone, each nappe of which is convex, there exists a line $L_1(y) \subset C(y)$ through y , sufficiently close to $L(y)$ (in terms of angles), such that $L_1(y) \cap S \neq y$, a contradiction (see Fig. 4). Hence, we have proved that \mathcal{P} is a desired family, and the proof is complete.

The hypotheses of Theorem 3 do not imply that a family \mathcal{P} necessarily exists such that each $x \in S$ is contained in a member of \mathcal{P} which intersects S in just the point x . In fact the following is true.

There exists a compact set $S \subset E_2$ such that every finite collection of points x_1, x_2, \dots, x_n in S is contained in some collection of parallel lines L_1, L_2, \dots, L_n such that $L_i \cap S = x_i$ ($i = 1, \dots, n$), and yet no family \mathcal{P} of parallel lines exist such that each point of S is contained in a member of \mathcal{P} which intersects S in just one point.

We exhibit such a set S as follows.

EXAMPLE. Let (x_i, y_i) denote rectangular coordinates of a point p_i in E_2 . We define the sequence of points $\{p_i, i = 1, 2, \dots\}$ in E_2 as follows,

$$(5) \quad \begin{aligned} (x_{2n-1}, y_{2n-1}) &\equiv \left(\frac{1}{2n-1}, 0 \right), \\ (x_{2n}, y_{2n}) &\equiv \left(\frac{1}{2n-1}, \frac{1}{2n-1} \right), \end{aligned}$$

so that $(x_1, y_1) = (1, 0)$, $(x_2, y_2) = (1, 1)$, $(x_3, y_3) = (1/3, 0)$, etc. Define S as follows

$$S \equiv \text{cl} \bigcup_i p_i p_{i+1}$$

so that S is the increasing limit of a sequence of zig-zag polygonal paths. Beginning at $(1, 0)$, the odd segments are vertical, and the even segments have finite positive slope. Observe that

$$\lim_{n \rightarrow \infty} \frac{y_{2n-2} - y_{2n-1}}{x_{2n-2} - x_{2n-1}} = \lim_{n \rightarrow \infty} \frac{2n-1}{2} = \infty$$

so that the even segments have slopes approaching ∞ . It is a simple matter to verify this set S has the property described in the above italicized statement because of (5) and because $x_{2n} = y_{2n}$ ($n = 1, 2, \dots$).

For the concluding result we need the following concepts used by Horn and Valentine [4].

DEFINITION 3. The set B is a set of visibility for a set S in E_n if for each point $x \in S$ there exists some point $y \in B$ such that $xy \subset S$.

DEFINITION 4. A set $S \subset E_n$ is said to be an L_2 set if each pair of points in S can be joined by a polygonal arc consisting of at most two line segments.

Horn and Valentine [4] proved that a simply connected compact L_2 set in E_2 is expressible as the union of convex sets every two of which have a point in common. No simple characterization of non-simply connected compact L_2 sets has ever been given. The following theorem is a step in that direction.

THEOREM 4. Let S be a compact L_2 set in E_n (see Definition 3),
 (a) then each hyperplane in E_n has a translate which intersects S in a set of visibility for S ,

(b) also each $(n-2)$ -dimensional flat is contained in a hyperplane which intersects S in a set of visibility for S .

Proof. For each point $x \in S$, let $S(x)$ denote the set of all points y such that $xy \subset S$. Also define $C(x)$ to be

$$C(x) = \text{conv } S(x).$$

Since S is compact, the set $C(x)$ is compact. Since every *two* members of the collection $\{C(x), x \in S\}$ have a point in common, a theorem of Klee [6] implies that each hyperplane H' has a translate H which intersects every $C(x)$, $x \in S$. Since $S(x)$ is the union of rays having x in common, the fact $H \cap C(x) \neq 0$ implies $H \cap S(x) \neq 0$. Hence, for each point $x \in S$, there exists a point $y \in H \cap S$ such that $xy \subset S$. This establishes (a). In the same manner a theorem of Horn [3] implies that each $(n - 2)$ -dimensional flat is contained in a hyperplane which intersects every $C(x)$, $x \in S$, and the remainder of the proof of (b) is identical to that given for (a).

Appendix

THEOREM (Helly [2]). *Let \mathcal{F} be a family of compact convex sets in E_n containing at least $n + 1$ members. If every $n + 1$ members of \mathcal{F} have a point in common, then all of the members of \mathcal{F} have a point in common.*

THEOREM (Krasnosel'skii [5]). *Let S be a compact connected set in E_n . Suppose that for every $n + 1$ points $x_i \in S$ ($i = 1, \dots, n + 1$) there exists at least one point $y \in S$ such that $x_i y \subset S$ ($i = 1, \dots, n$). Then there exists a point $p \in S$ such that $xp \subset S$ for each point $x \in S$.*

THEOREM. *Let \mathcal{F} be a family of bounded closed convex sets in a Euclidean space E . Suppose \mathcal{F} contains at least n members. Suppose every n members of \mathcal{F} have a point in common.*

(Klee) *Then every flat of deficiency $n - 1$ has a translate which intersects every member of \mathcal{F} .*

(Horn) *Every flat of deficiency n is contained in a flat of deficiency $n - 1$ which intersects every member of \mathcal{F} .*

BIBLIOGRAPHY

1. L. Danzer, B. Grünbaum and V. Klee, *Helly's theorem and its relatives*, Symposium on Convexity, Proceedings of the Symposia in Pure Math. **17** Amer. Math. Soc. (1963).
2. E. Helly, *Über Mengen konvexer Körper mit gemeinschaftlichen Punkten*, Jber. Deutch Math. Verein **32** (1923), 175-176.
3. Alfred Horn, *Some generalizations of Helly's theorem on convex sets*, Bull. Amer. Math. Soc. **55** (1949), 923-929.
4. Alfred Horn and F. A. Valentine, *Some properties of L sets in the plane*, Duke Math. J. **16** (1949), 131-140.
5. M. A. Krasnoselskii, *Sur un critère pour qu'on domain soit étoilé*, Math. Sb. (61) **19** (1946), 309-310.
6. Victor Klee, *On certain intersection properties of convex sets*, Canad. J. Math. **3** (1951), 271-275.
7. F. A. Valentine, *Convex sets*, McGraw-Hill Book Co. (1964).

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California

R. M. BLUMENTHAL

University of Washington
Seattle, Washington 98105

*J. DUGUNDJI

University of Southern California
Los Angeles, California 90007

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Pacific Journal of Mathematics

Vol. 16, No. 1

November, 1966

Larry Armijo, <i>Minimization of functions having Lipschitz continuous first partial derivatives</i>	1
Edward Martin Bolger and William Leonard Harkness, <i>Some characterizations of exponential-type distributions</i>	5
James Russell Brown, <i>Approximation theorems for Markov operators</i>	13
Doyle Otis Cutler, <i>Quasi-isomorphism for infinite Abelian p-groups</i>	25
Charles M. Glennie, <i>Some identities valid in special Jordan algebras but not valid in all Jordan algebras</i>	47
Thomas William Hungerford, <i>A description of $\text{Mult}_i(A^1, \dots, A^n)$ by generators and relations</i>	61
James Henry Jordan, <i>The distribution of cubic and quintic non-residues</i>	77
Junius Colby Kegley, <i>Convexity with respect to Euler-Lagrange differential operators</i>	87
Tilla Weinstein, <i>On the determination of conformal imbedding</i>	113
Paul Jacob Koosis, <i>On the spectral analysis of bounded functions</i>	121
Jean-Pierre Kahane, <i>On the construction of certain bounded continuous functions</i>	129
V. V. Menon, <i>A theorem on partitions of mass-distribution</i>	133
Ronald C. Mullin, <i>The enumeration of Hamiltonian polygons in triangular maps</i>	139
Eugene Elliot Robkin and F. A. Valentine, <i>Families of parallels associated with sets</i>	147
Melvin Rosenfeld, <i>Commutative F-algebras</i>	159
A. Seidenberg, <i>Derivations and integral closure</i>	167
S. Verblunsky, <i>On the stability of the set of exponents of a Cauchy exponential series</i>	175
Herbert Walum, <i>Some averages of character sums</i>	189