

Pacific Journal of Mathematics

DERIVATIONS AND INTEGRAL CLOSURE

A. SEIDENBERG

DERIVATIONS AND INTEGRAL CLOSURE

A. SEIDENBERG

Let \mathcal{O} be an integral domain containing the rational numbers, Σ its quotient field, D a derivation of Σ , and \mathcal{O}' the ring of elements in Σ quasi-integral over \mathcal{O} . It is shown that if $D\mathcal{O} \subset \mathcal{O}$, then $D\mathcal{O}' \subset \mathcal{O}'$.

According to a lemma of Posner [4], which is also used by him in a subsequent paper [5], if \mathcal{O} is a finite integral domain over a ground field F of characteristic 0 and D is a derivation over F sending \mathcal{O} into itself, then D also sends the integral closure of \mathcal{O} into itself. The proof of this in [4] is wrong, but the statement itself is correct and a proof is here supplied. More generally it is proved that if \mathcal{O} is any integral domain containing the rational numbers and D is a derivation such that $D\mathcal{O} \subset \mathcal{O}$, then $D\mathcal{O}' \subset \mathcal{O}'$, where \mathcal{O}' is the ring of elements in the quotient field Σ of \mathcal{O} that are quasi-integral over \mathcal{O} . The theorem is not true for characteristic $p \neq 0$, but if one uses the Hasse-Schmidt differentiations instead of derivations, one gets the corresponding theorem for a completely arbitrary integral domain \mathcal{O} .

Let \mathcal{O} be an arbitrary integral domain containing the rational numbers, and let $\bar{\mathcal{O}}$ be the integral closure of \mathcal{O} . The question whether $D\mathcal{O} \subset \mathcal{O}$ implies $D\bar{\mathcal{O}} \subset \bar{\mathcal{O}}$ is related to the question whether the ring of formal power series $\bar{\mathcal{O}}[[t]]$ is integrally closed. Thus consider the statements:

A. *For every \mathcal{O} , $D\mathcal{O} \subset \mathcal{O}$ implies $D\bar{\mathcal{O}} \subset \bar{\mathcal{O}}$, and*

B. *For every \mathcal{O} , $\bar{\mathcal{O}}[[t]]$ is integrally closed.* We show that A and B are equivalent statements. (We also show: C. *If $\bar{\mathcal{O}}[[t]]$ is integrally closed, then $D\mathcal{O} \subset \mathcal{O}$ implies $D\bar{\mathcal{O}} \subset \bar{\mathcal{O}}$.) Now according to the last exercise in Nagata's book *Local Rings*, [3; p. 202, Ex. 5], B is a true statement, but we give a counter-example, which also leads to a counter-example for A.*

2. Criticism of Posner's proof. Posner purports to prove that if P is a place of the quotient field Σ of \mathcal{O} that has F as residue field and is finite on \mathcal{O} and if $g \in \Sigma$ is finite at P , then Dg is finite at P . This is not so, as the following example shows. Let $\mathcal{O} = F[X, Y]$ be polynomial ring in two indeterminates over F . Let $D = \partial/\partial X$. Let P_1 be the place of $F(X, Y)$ over $F(Y/X)$ obtained by mapping X into 0, let P_2 be the place of $F(Y/X)/F$ obtained by

Received February 13, 1964 and in revised form June 25, 1964. This work was supported in part by the Air Force Office of Scientific Research.

mapping Y/X into any element of F , and let P be the composite place. Then $X, Y, Y/X$ are finite at P , but $\partial(Y/X)/\partial X = -Y/X^2$ is not.¹

One reason that Posner's proof fails is that there are no parameters such as those of which he speaks, except in the case that the degree of transcendency of \mathcal{O}/F is 1. In that case, Posner's argument yields a proof.

3. A generalization. Let \mathcal{O} be an arbitrary domain, with quotient field Σ . An element $\alpha \in \Sigma$ is said to be quasi-integral over \mathcal{O} if all powers of α are contained in a finite \mathcal{O} -module contained in Σ , or, what comes to the same, if there is a $d \in \mathcal{O}, d \neq 0$, such that $d\alpha^\rho \in \mathcal{O}, \rho = 0, 1, \dots$; (see [2]). If \mathcal{O} is a Noetherian domain, then the concepts of integral dependence and quasi-integral dependence (for elements in Σ) become the same; but it is the concept of quasi-integral dependence, rather than that of integral dependence, which is adapted to our considerations. The elements in Σ that are quasi-integral over \mathcal{O} form a ring \mathcal{O}' , which in the case \mathcal{O} is Noetherian is the integral closure $\bar{\mathcal{O}}$ of \mathcal{O} . The base field F plays little role, and it will be sufficient to assume that \mathcal{O} contains the rational numbers.

THEOREM. *Let \mathcal{O} be an arbitrary integral domain containing the rational numbers, let \mathcal{O}' be the ring of elements in the quotient field Σ of \mathcal{O} quasi-integral over \mathcal{O} , and let D be a derivation of Σ . Then: if $D\mathcal{O} \subset \mathcal{O}$, then $D\mathcal{O}' \subset \mathcal{O}'$.*

Proof. Let $\Sigma[[t]]$ be the ring of formal power series in a letter t over Σ and let $\Sigma((t))$ be its quotient field. The mapping $\Sigma c_i t^i \rightarrow \Sigma(Dc_i)t^i, i \geq 0, c_i \in \Sigma$, is a derivation of $\Sigma[[t]]$ into itself and extends D ; it has a unique extension to $\Sigma((t))$, which will also be denoted D . Let E be the expression $1 + tD + (t^2/2!)D^2 + \dots (=e^{tD})$. Then $\alpha + tD\alpha + (t^2/2!)D^2\alpha + \dots$, to be denoted $E\alpha$, has a meaning for every $\alpha \in \Sigma[[t]]$, i.e., the partial sums converge in the topology defined by powers of (t) ; and the mapping $\alpha \rightarrow E\alpha$ is an isomorphism of $\Sigma[[t]]$ into itself, as one easily verifies.² Its unique extension to $\Sigma((t))$ will

¹ Far from all, or even infinitely many, valuation rings \mathfrak{B} centered at (X, Y) being sent into themselves by $D = \partial/\partial X$, there is one and only one. In fact, restricting oneself to valuation rings \mathfrak{B} centered at (X, Y) , if $D\mathfrak{B} \subset \mathfrak{B}$, then $X/Y \notin \mathfrak{B}$, since $D(X/Y) = 1/Y \notin \mathfrak{B}$. Hence $Y/X \in \mathfrak{B}$, and therefore $D(Y/X), D^2(Y/X)$, etc. are also in \mathfrak{B} . Since $D^{n-1}(Y/X) = c_n Y/X^n (c_n \in K), v(Y) \geq n v(X)$ for $n = 1, 2, \dots$, where v is the valuation corresponding to \mathfrak{B} . Thus \mathfrak{B} could not be other than the ring of the valuation in which $v(X)$ is infinitely small with respect to $v(Y)$; and for that ring one checks that $D\mathfrak{B} \subset \mathfrak{B}$.

² We only use that $\alpha \rightarrow E\alpha$ is a monomorphism, but it is actually onto $\Sigma[[t]]$ as one sees from the identity $e^{tD}(e^{-tD}\alpha) = \alpha$.

also be denoted E . Since $D\mathcal{O} \subset \mathcal{O}$, one has $D\mathcal{O}[[t]] \subset \mathcal{O}[[t]]$, and since \mathcal{O} contains the rationals, $E\mathcal{O}[[t]] \subset \mathcal{O}[[t]]$.

Let α be quasi-integral over \mathcal{O} , and let $d \in \mathcal{O}$ be such that $d\alpha^\rho \in \mathcal{O}$, $\rho = 0, 1, \dots$. Then $E(d\alpha^\rho) = Ed(E\alpha)^\rho \in \mathcal{O}[[t]]$, $\rho = 0, 1, \dots$. Hence $dEd(E\alpha - \alpha)^\rho \in \mathcal{O}[[t]]$, $\rho = 0, 1, \dots$; here we use that d and Ed are in $\mathcal{O}[[t]]$. The coefficient of t^ρ in $dEd(E\alpha - \alpha)^\rho$, i.e., the leading coefficient, is $d^2(D\alpha)^\rho$; and this coefficient, as well as all the others, are in \mathcal{O} . Hence $D\alpha$ is quasi-integral over \mathcal{O} .

COROLLARY. *If $d \in \mathcal{O}$ and $\alpha \in \Sigma$ are such that $d\alpha^i \in \mathcal{O}$, $i = 0, 1, \dots, \rho$, then $d^2(D\alpha)^i \in \mathcal{O}$, $i = 0, 1, \dots, \rho$.*

Let $\mathfrak{C} = \{c \mid c \in \mathcal{O}, c\mathcal{O}' \subset \mathcal{O}'\}$; then \mathfrak{C} is an ideal, which in the case \mathcal{O}' is the integral closure $\bar{\mathcal{O}}$ of \mathcal{O} is called the conductor of \mathcal{O} .

COROLLARY. *If $D\mathcal{O} \subset \mathcal{O}$, then $D\mathfrak{C} \subset \mathfrak{C}$. In other words, \mathfrak{C} is a differential ideal for any derivation (or any family of derivations) sending \mathcal{O} into itself.*

Proof. If $c \in \mathfrak{C}$ and $\alpha \in \mathcal{O}'$, then $(Dc)\alpha = D(c\alpha) - cD\alpha \in \mathcal{O}$, so that also $(Dc)\mathcal{O}' \subset \mathcal{O}'$.

The last corollary can sometimes be used to prove that a given integral domain \mathcal{O} is integrally closed (see [4]). We first restrict ourselves to a class of integral domains \mathcal{O} such that $\bar{\mathcal{O}} = \mathcal{O}'$, for example, the class of Noetherian domains. Then we restrict ourselves further to a class \mathcal{E} of domains \mathcal{O} such that \mathcal{O} has a conductor $\mathcal{O}': \bar{\mathcal{O}} \neq (0)$, or equivalently, such that $\bar{\mathcal{O}}$ is contained in a finite \mathcal{O} -module (contained in Σ), for example, the class of finite integral domains (see [7; p. 267]), or quotient rings thereof, or the class of complete local domains (see [3; p. 114]). (For examples of Noetherian domains not having this property, see [3; p. 205 ff]; for an example in characteristic 0, see [1]). Then we can state:

COROLLARY. *Let \mathcal{O} be an integral domain belonging to a class \mathcal{E} defined just above, let \mathcal{O} contain the rational numbers, and let $\{D\}$ be a (finite or infinite) family of derivations of \mathcal{O} into itself. Then, if \mathcal{O} is differentially simple under $\{D\}$ (i.e., has no differential ideal other than (0) or (1)), then \mathcal{O} is integrally closed.*

4. Extension of D to $\bar{\mathcal{O}}$. The above is a simplification of our original proof for a finite integral domain. The idea was that since E sends $\mathcal{O}[[t]]$ into itself, it also sends the integral closure of $\mathcal{O}[[t]]$

into itself. It was then sufficient to prove that $\bar{\mathcal{O}}[[t]]$ is integrally closed; in fact, we have the following theorem for any integral domain \mathcal{O} containing the rational numbers.

THEOREM C. *If $\bar{\mathcal{O}}[[t]]$ is integrally closed and $D\mathcal{O} \subset \mathcal{O}$, then $D\bar{\mathcal{O}} \subset \bar{\mathcal{O}}$. (Here $\bar{\mathcal{O}}$ is the integral closure of \mathcal{O} .)*

Proof. If $\alpha \in \Sigma$, $\alpha = c/d$, $c, d \in \mathcal{O}$, then $E\alpha = Ec/Ed$, so $E\alpha$ is in the quotient field of $\mathcal{O}[[t]]$. If α is integral over \mathcal{O} , then $E\alpha = \alpha + tD\alpha + \dots$ is integral over $\mathcal{O}[[t]]$, hence in $\bar{\mathcal{O}}[[t]]$, whence $D\alpha \in \bar{\mathcal{O}}$.

Our proof that $\bar{\mathcal{O}}[[t]]$ was integrally closed for \mathcal{O} a finite integral domain depended on the following observation, which holds for an arbitrary domain \mathcal{O} .

THEOREM. *If \mathcal{O} is completely integrally closed (i.e., if $\mathcal{O}' = \mathcal{O}$), then so is $\mathcal{O}[[t]]$. More generally, for any \mathcal{O} , $(\mathcal{O}[[t]])' \subset \mathcal{O}'[[t]]$.*

Proof. Let $\alpha(t)$ be quasi-integral over $\mathcal{O}[[t]]$. Then there is a $d \in \mathcal{O}[[t]]$, $d = d(t) \neq 0$, such that $d\alpha^\rho \in \mathcal{O}[[t]]$, $\rho = 0, 1, \dots$. Since $\text{ord } d + \rho \text{ ord } \alpha \geq 0$, $\rho = 0, 1, \dots$, one first observes that $\alpha \in \Sigma[[t]]$. Let $d = d_s t^s + d_{s+1} t^{s+1} + \dots$, $d_s \neq 0$, and let $\alpha = \alpha_r t^r + \alpha_{r+1} t^{r+1} + \dots$. Since the leading coefficient of $d\alpha^\rho$ is in \mathcal{O} , we have $d_s \alpha_r^\rho \in \mathcal{O}$, whence α_r is quasi-integral over \mathcal{O} . Now $\alpha - \alpha_r t^r$ is quasi-integral over $\mathcal{O}[[t]]$, whence α_{r+1} is quasi-integral over \mathcal{O} ; and in this way one sees that all the coefficients of α are quasi-integral over \mathcal{O} .

If \mathcal{O} is Noetherian, then so is $\mathcal{O}[[t]]$. Hence:

COROLLARY. *If \mathcal{O} is an integrally closed Noetherian domain, then so is $\mathcal{O}[[t]]$.*

This is Nagata's (47.6) in [3; p. 200].

Finally, if \mathcal{O} is a finite integral domain, then so is $\bar{\mathcal{O}}$, whence in this case $\bar{\mathcal{O}}[[t]]$ is integrally closed. Recalling that $\bar{\mathcal{O}}$ is a finite \mathcal{O} -module (see [7; p. 267]), one sees that $\bar{\mathcal{O}}[[t]]$ is even the integral closure of $\mathcal{O}[[t]]$ in accordance with the following:

THEOREM. *Let \mathcal{O} be an integral domain whose integral closure is Noetherian and is a finite \mathcal{O} -module. Then the integral closure of $\mathcal{O}[[t]]$ is $\bar{\mathcal{O}}[[t]]$.*

Proof. Let $\bar{\mathcal{O}} = \mathcal{O}w_1 + \dots + \mathcal{O}w_s$. Then

$$\mathcal{O}[[t]] = \mathcal{O}[[t]]w_1 + \cdots + \mathcal{O}[[t]]w_s,$$

whence $\bar{\mathcal{O}}[[t]]$ is a finite $\mathcal{O}[[t]]$ -module and thus integral over $\mathcal{O}[[t]]$. Let d be a common denominator of the w_i when written as quotients of elements in \mathcal{O} . Then $d\bar{\mathcal{O}}[[t]] \subset \bar{\mathcal{O}}[[t]]$, whence $\mathcal{O}[[t]]$ and $\bar{\mathcal{O}}[[t]]$ have the same quotient field. As we have already seen that $\bar{\mathcal{O}}[[t]]$ is integrally closed, the proof is complete.

Although not necessary for our considerations, we mention the following:

THEOREM. *If \mathcal{O} is a Noetherian domain, then $\bar{\mathcal{O}}[[t]]$ is integrally closed, where t abbreviates a set t_1, \dots, t_n of n distinct letters.*

Proof. $\bar{\mathcal{O}}$ is a Krull ring (see [3; p. 118]), hence from the definition [3; p. 115], $\bar{\mathcal{O}}_p$ is a Noetherian valuation ring for every minimal prime ideal p of $\bar{\mathcal{O}}$. Moreover $\bar{\mathcal{O}} = \bigcap \bar{\mathcal{O}}_p$, where the intersection is taken over the minimal prime ideals of $\bar{\mathcal{O}}$ (see [3; p. 116]). Since $\bar{\mathcal{O}}_p[[t]]$ is integrally closed, also $\bar{\mathcal{O}}[[t]] = \bigcap \bar{\mathcal{O}}_p[[t]]$ is integrally closed.

Now consider the statements A and B mentioned at the beginning. We say that A and B are equivalent. Recall that we are assuming that \mathcal{O} contains the rational numbers.

$B \Rightarrow A$. This follows at once from C, the first theorem of this section.

$A \Rightarrow B$. Let α be in the quotient field of $\bar{\mathcal{O}}[[t]]$ and integral over $\bar{\mathcal{O}}[[t]]$. Then $\alpha \in \Sigma[[t]]$, $\alpha = \alpha_0 + \alpha_1 t + \cdots$. From an equation of integral dependence for α on $\bar{\mathcal{O}}[[t]]$, by placing $t = 0$, one sees that $\alpha_0 \in \bar{\mathcal{O}}$. Now apply A to the ring $\bar{\mathcal{O}}[[t]]$ and the derivation $D = \partial/\partial t$. Then $\partial\alpha/\partial t, \partial^2\alpha/\partial t^2, \dots$ are integral over $\bar{\mathcal{O}}[[t]]$, whence all the coefficients of α are in $\bar{\mathcal{O}}$.

Now according to the last exercise in Nagata's *Local Rings*, B is a true statement; however, we will show that this is incorrect.

THEOREM. *If \mathcal{O} is an (integrally closed) integral domain containing a field and there is a nonunit $b \in \mathcal{O}$ such that $\bigcap (b^n) \neq (0)$, then $\mathcal{O}[[t]]$ is not integrally closed.*

Proof. Let p be the characteristic and $n > 1$, an integer such that $n \not\equiv 0(p)$. Then $b^n + b^{n-2}t$ has an n th root $\alpha = b[1 + (t/b^2)]^{1/n} = b[1 + c_1(t/b^2) + c_2(t^2/b^4) + \cdots]$ in $\Sigma[[t]]$, where c_1, c_2, \dots are in the prime field of Σ and $c_1 \neq 0$. If $a \in \bigcap (b^n)$ and $a \neq 0$, then $a\alpha \in \mathcal{O}[[t]]$, so

that α is in the quotient field of $\mathcal{O}[[t]]$. Now α is integral over $\mathcal{O}[[t]]$, but is not in $\mathcal{O}[[t]]$. Hence $\mathcal{O}[[t]]$ is not integrally closed.

THEOREM. *Let \mathfrak{B} be a (proper) valuation ring containing a field. Then $\mathfrak{B}[[t]]$ is integrally closed if and only if \mathfrak{B} is of rank 1, i.e., if and only if there is no chain $0 < p_1 < p_0 < \mathfrak{B}$ of prime ideals.*

Proof. If \mathfrak{B} is of rank 1, then it is well-known and can be checked at once, that \mathfrak{B} is completely integrally closed. Hence $\mathfrak{B}[[t]]$ is completely integrally closed, hence integrally closed.

On the other hand, if \mathfrak{B} is of rank > 1 and $0 < p_1 < p_0 < \mathfrak{B}$ is a chain of prime ideals in \mathfrak{B} and $b \in p_0 - p_1$, then $p_1 \subset \cap (b^p)$, whence $\mathfrak{B}[[t]]$ is not integrally closed.

To get a counter-example to Nagata's last exercise, one has but to take \mathcal{O} to be a valuation ring of rank > 1 that contains a field.³

To get an example of a ring \mathcal{O} and derivation D such that $D\mathcal{O} \subset \mathcal{O}$ but $D\bar{\mathcal{O}} \not\subset \bar{\mathcal{O}}$, let \mathfrak{B} be a valuation ring of rank 2 containing the rational numbers, let $\mathcal{O} = \mathfrak{B}[[t]]$ and $D = \partial/\partial t$. Let b be a nonunit in \mathfrak{B} such that $\cap (b^p) \neq (0)$, and let

$$\alpha = (b^2 + t)^{1/2} = b \left[1 + c_1 \frac{t}{b^2} + c_2 \frac{t^2}{b^4} + \cdots \right],$$

where c_1, c_2, \dots are rational numbers. Then α is integral over $\mathcal{O} = \mathfrak{B}[[t]]$ but $D\alpha$ is not.

Concerning the proof spoken of at the beginning of this section, the author is obliged to Professor Mumford for the remark in context that if D is a derivation, then e^D , formally at any rate, is an isomorphism. The introduction of the parameter t on the one hand prevents the computations from collapsing into meaninglessness, and on the other allows one to recover D from e^{tD} .

5. The case of characteristic $p \neq 0$. For $p \neq 0$, the theorem of § 3 is not true, even for curves. Thus consider the curve given by $Y^p - X^p - X^{p+1} = 0$. One checks that $Y^p - X^p - X^{p+1}$ is irreducible (over the ground field F). Let (x, y) be a generic point of the curve over F . Let D be a derivation of $F(y)/F$ with $Dy = 1$; since x is separable over $F(y)$, D can be extended uniquely to a derivation, still to be denoted D , of $F(y, x)$. One finds $-(p+1)x^p Dx = 0$, hence $Dx = 0$. Let $\mathcal{O} = F[x, y]$. Then $D\mathcal{O} \subset \mathcal{O}$. Now y/x is integral

³ In reference to the exercise, Nagata [3; p. 221] cites Sugaku, Vol. 9, No. 1 (1957), p. 61, which we have not been able to locate; and while he notes that the proof there is not complete, he remarks that "a supplement is expected to appear soon".

over \mathcal{O} , since $(y/x)^p = 1 + x$, but $D(y/x) = 1/x$ is not, as otherwise it would be integral over $F[x]$.

However, if one uses the Hasse-Schmidt differentiations [6] instead of derivations, one gets the corresponding theorem.⁴ Recall that a differentiation D of a field Σ into itself is a sequence $D = (\delta_0, \delta_1, \delta_2, \dots)$ of mappings of Σ into itself with $\delta_0 = 1$ and satisfying the properties:

$$\begin{aligned} \delta_i(x + y) &= \delta_i x + \delta_i y \\ \delta_i xy &= \sum_{i+j=v} \delta_i x \delta_j y . \end{aligned}$$

By $D\mathcal{O} \subset \mathcal{O}$ we now mean $\delta_i \mathcal{O} \subset \mathcal{O}$ for every i . Then

$$E = \delta_0 + t\delta_1 + t^2\delta_2 + \dots$$

still yields an isomorphism and can be used instead of our previous E to get the conclusion $D\mathcal{O}' \subset \mathcal{O}'$. (After obtaining $\delta_1 \mathcal{O}' \subset \mathcal{O}'$ as before, we argue that $d^3 Ed(E\alpha - \alpha - t\delta_1 \alpha)^\rho \in \mathcal{O}[[t]]$, $\rho = 0, 1, \dots$, whence $d^4(\delta_2 \alpha)^\rho \in \mathcal{O}$, $\rho = 0, 1, \dots$, and $\delta_2 \alpha$ is quasi-integral over \mathcal{O} , etc.) In the case of characteristic 0, the same argument shows one can drop the assumption that \mathcal{O} contains the rationals (i.e., if one uses differentiations instead of derivations).

The corollaries of the theorem of § 3 also have easily stated generalizations, with similar proofs.

REMARK. Since $(1 + (1 + 4t)^{1/2})/2 \in Z[[t]]$, the last two theorems of § 4 hold without the field condition.

REFERENCES

1. Y. Akizuki, *Einige Bemerkungen über primäre Integritätsbereich mit Teilerkettenansatz*, Proc. Phys.-Math. Soc. Japan, 3rd ser., **17** (1935), 327-336.
2. W. Krull, *Beiträge zur Arithmetik kommutativer Integritätsbereiche, II*, Math. Zeitschrift, **41** (1936), 665-679.
3. M. Nagata, *Local Rings*. New York, 1962.
4. E. C. Posner, *Integral closure of differential rings*, Pacific J. Math. **10** (1960), 1393-1396.
5. ———, *Integral closure of rings of solutions of linear differential equations*, Pacific J. Math. **12** (1962), 1417-1422.
6. F. K. Schmidt, *Noch eine Begründung der Theorie der höheren Differentialquotienten in einer algebraischen Funktionenkörper einer Unbestimmten. Zusatz bei der Korrektur*, J. für die reine u. angewandte Math. **177** (1937), 223-237.
7. O. Zariski and P. Samuel, *Commutative Algebra*, Vol. 1. New York, 1958.

HARVARD UNIVERSITY
AND
UNIVERSITY OF CALIFORNIA, BERKELEY

⁴ For some useful information on differentiations, see K. Okugawa, "Basic properties of differential fields of an arbitrary characteristic and the Picard-Vessiot theory", J. of Math. of Kyoto Univ., Vol. 2 (1963), pp. 295-322.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California

R. M. BLUMENTHAL

University of Washington
Seattle, Washington 98105

*J. DUGUNDJI

University of Southern California
Los Angeles, California 90007

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Pacific Journal of Mathematics

Vol. 16, No. 1

November, 1966

Larry Armijo, <i>Minimization of functions having Lipschitz continuous first partial derivatives</i>	1
Edward Martin Bolger and William Leonard Harkness, <i>Some characterizations of exponential-type distributions</i>	5
James Russell Brown, <i>Approximation theorems for Markov operators</i>	13
Doyle Otis Cutler, <i>Quasi-isomorphism for infinite Abelian p-groups</i>	25
Charles M. Glennie, <i>Some identities valid in special Jordan algebras but not valid in all Jordan algebras</i>	47
Thomas William Hungerford, <i>A description of $\text{Mult}_i(A^1, \dots, A^n)$ by generators and relations</i>	61
James Henry Jordan, <i>The distribution of cubic and quintic non-residues</i>	77
Junius Colby Kegley, <i>Convexity with respect to Euler-Lagrange differential operators</i>	87
Tilla Weinstein, <i>On the determination of conformal imbedding</i>	113
Paul Jacob Koosis, <i>On the spectral analysis of bounded functions</i>	121
Jean-Pierre Kahane, <i>On the construction of certain bounded continuous functions</i>	129
V. V. Menon, <i>A theorem on partitions of mass-distribution</i>	133
Ronald C. Mullin, <i>The enumeration of Hamiltonian polygons in triangular maps</i>	139
Eugene Elliot Robkin and F. A. Valentine, <i>Families of parallels associated with sets</i>	147
Melvin Rosenfeld, <i>Commutative F-algebras</i>	159
A. Seidenberg, <i>Derivations and integral closure</i>	167
S. Verblunsky, <i>On the stability of the set of exponents of a Cauchy exponential series</i>	175
Herbert Walum, <i>Some averages of character sums</i>	189