

Pacific Journal of Mathematics

**REFLECTION AND APPROXIMATION BY INTERPOLATION
ALONG THE BOUNDARY FOR ANALYTIC FUNCTIONS**

JAMES MCLEAN SLOSS

REFLECTION AND APPROXIMATION BY INTERPOLATION ALONG THE BOUNDARY FOR ANALYTIC FUNCTIONS

JAMES M. SLOSS

Let there be given a function $f(z)$ analytic in an open connected set, not necessarily simply connected, which is bounded by simple closed analytic curves such that the function is continuous on the closure of the region and such that the real part of the function satisfies boundary conditions that are analytic in a neighborhood of the boundary. We want to interpolate $f(z)$ along the boundaries and find conditions that make the interpolants converge maximally to $f(z)$ throughout the closure of the region. The boundary condition on the real part of $f(z)$ permits the analytic continuation of $f(z)$ across the boundary curves and ensures that we are interpolating at points interior to the region of analyticity. In our error estimates (Theorem 1) maximal convergence depends in an essential way on how far we can reflect $f(z)$ and this in turn depends on the boundary values of the real part of $f(z)$ as well as on the geometry of the given region and its analytic boundaries. In Theorems 2 and 3, a simply connected region is considered. Special points of interpolation are given, these depend only on the parametric representation of the boundary curves and not a conformal map. These points are the image points of the Chebyshev polynomials.

Finally an example is given for a multiply connected region.

As is well known [2] Runge's beautiful theorem shows us that there exist certain "equidistributed" points on the analytic curves such that if we interpolate at these points the interpolants converge to the function. However, the proof depends on knowing the conformal map in order to know what the interpolation points are. Here we shall give conditions that do not require knowledge of the conformal map but for convergence depend on how far we can reflect. Along with these, we shall give simple error estimates. Moreover, we shall show that possible interpolation points are the images on the boundary of roots of the Chebyshev polynomials.

The aspects of this paper which are novel are

- (i) the use of reflection
- (ii) interpolation at boundary points which are gotten directly from the parametric representation of the boundary and do not depend on a conformal map

(iii) the use of the images of the roots of the Chebyshev polynomials as possible interpolation points.

Notation. Let R be a connected set whose boundary is Γ . Let $\Gamma = \Gamma^1 \cup \Gamma^2 \cup \dots \cup \Gamma^s$ where the Γ^j are bounded analytic contours in the $z = x + iy$ plane given by $F^j(x, y) = 0$ with $(F_x^j)^2 + (F_y^j)^2 \neq 0$ along $\Gamma^j, j = 1, 2, \dots, s$, where the F^j are real-valued analytic functions of x and y . We assume further that the Γ^j are pairwise disjoint. Let Γ^1 contain in its interior $\Gamma^2, \Gamma^3, \dots, \Gamma^s$ and contain in its exterior the point at infinity. Let Γ^j contain in its interior $a_j, 2 \leq j \leq s$. As shown in [3] there are "reflection" functions $G_j(z)$ defined on a neighborhood $D^j \cup \Gamma^j \cup \hat{D}^j$ of Γ^j . Assume $G_j(z)$ single-valued on $D^j \cup \Gamma^j \cup \hat{D}^j$ [3] shows.

(1) $z = \overline{G_j(z)}$ is Γ^j .

(2) $G_j(z)$ is analytic on $D^j \cup \Gamma^j \cup \hat{D}^j$, where D^j is contained in the connected R and D^j is contained in the complement of $\Gamma^j \cup D^j$ for $j = 1, 2, \dots, s$.

(3) The transformation $\hat{z} = \overline{G(z)}$ is an involution; i.e. $\hat{\hat{z}} = z$.

(4) If z is in D^j then \hat{z} is in \hat{D}^j and if z is in \hat{D}^j then \hat{z} is in D^j .

(5) $\overline{G[D^j]} = \hat{D}^j$ and $\overline{G[\hat{D}^j]} = D^j$. We assume the boundary of \hat{D}^j , that is not Γ^j , is a contour C^j and $G_j(z)$ is continuous on $\hat{D}^j \cup C^j$.

THEOREM 1. (H 1) *Let $f(z)$ be an analytic single valued function on R whose boundary is Γ such that the real part of $f(z)$ solves the Dirichlet problem in R with real boundary values $B_j(z)$ on Γ^j where $B_j(z)$ are single-valued and continuous in $D^j \cup \Gamma^j \cup \hat{D}^j \cup C^j$ and analytic in $D^j \cup \Gamma^j \cup \hat{D}^j$. Let $f(z)$ be continuous and single-valued on $R \cup \Gamma$.*

(H. 2) *Let $z_{n_j^1}^j, z_{n_j^2}^j, \dots, z_{n_j^{n_j+1}}^j, n_j = 0, 1, 2, \dots$, be points of $\Gamma^j, j = 1, 2, \dots, s$. Let $p_{n_1}^j(z)$ be the polynomial in z of degree n_1 that agrees with $B_1(z)$ at $z_{n_1^1}^1, z_{n_1^2}^1, \dots, z_{n_1^{n_1+1}}^1$ and let $p_{n_j}^j(z) (2 \leq j \leq s)$ be the polynomial in $1/(z - a_j)$ that agrees with $B_j(z)$ for $z - a_j = z_{n_j^1}^j, z_{n_j^2}^j, \dots, z_{n_j^{n_j+1}}^j$ where a_j is a point inside c^j .*

(H. 3) *Let*

$$\delta_{n_1}^1 = \min_{t \text{ on } C^1} \prod_{k=1}^{n_1+1} |t - z_{n_1^k}^1|$$

$$\mu_{n_1}^1 = \max_{z \text{ on } \Gamma^1} \prod_{k=1}^{n_1+1} |z - z_{n_1^k}^1|$$

and

$$\delta_{n_j}^j = \min_{t \text{ on } C^j} \prod_{k=1}^{n_j+1} \left| \frac{t - a_j - z_{n_j^k}^j}{t - a_j} \right|, \quad 2 \leq j \leq s$$

$$\mu_{n_j}^j = \max_{z \text{ on } \Gamma^j} \prod_{k=1}^{n_j+1} \left| \frac{z - a_j - z_{n_j^k}^j}{z - a_j} \right|, \quad 2 \leq j \leq s.$$

(H. 4) Let $\mu_{n_j}^j/\delta_{n_j}^j \rightarrow 0$ as $n_j \rightarrow \infty, j = 1, 2, \dots, s$. Then for

$$\mu = \left(\frac{1}{n_1}, \frac{1}{n_2}, \dots, \frac{1}{n_s} \right) \text{ and } |\mu| = \max \left\{ \frac{1}{n_1}, \frac{1}{n_2}, \dots, \frac{1}{n_s} \right\}.$$

(C. 1) $R_\mu(z) = \sum_{j=1}^s p_{n_j}^j(z)$ converges uniformly to $f(z)$ in $R \cup \Gamma$ as $|\mu| \rightarrow 0$ and thus $\text{Re } R_\mu(z)$ converges uniformly to $u(x, y)$ in R and uniformly to $B_j(z)$ on Γ^j .

(C. 2) Moreover in $R \cup \Gamma$:

$$|f(z) - R_\mu(z)| \leq \frac{1}{2\pi} \sum_{j=1}^s \frac{L_j M_j}{\delta_j} \mu_{n_j}^j / \delta_{n_j}^j$$

where $L_j = \text{length of } C^j, M_j = \max_{t \text{ on } C^j} |f(t)|$ and $\delta_j = \inf_{z \text{ on } \Gamma^j} \min_{t \text{ on } C^j} |t - z|$.

Proof. In order to avoid notation that only confuses, we shall prove the theorem for the case $s = 2$.

We first analytically continue $f(z)$ into $R \cup \Gamma^1 \cup \hat{D}^1 \cup \Gamma^2 \cup \hat{D}^2$. Let $f_j^*(z) = \overline{f[G_j(z)]}$ for z in $\Gamma^j \cup \hat{D}^j$. $f_j^*(z)$ is defined and analytic for z in $\Gamma^j \cup \hat{D}^j$ since $\hat{z} = \overline{G_j(z)}$ is in D^j for z in \hat{D}^j and $G_j(z)$ is analytic for z in $\Gamma^j \cup \hat{D}^j$. But $f_j^*(z) = \overline{f(z)}$ for z on Γ^j , thus on Γ^j

$$f(z) + f_j^*(z) = 2B_j(z).$$

Thus $f(z) = 2B_j(z) - f_j^*(z)$ analytically continues $f(z)$ into $\Gamma^j \cup \hat{D}^j$ since $f(z)$ is continuous up to and on Γ^j . Moreover, $f(z)$ is continuous on $\Gamma^j \cup \hat{D}^j \cup C^j$ since $G_j(z)$ and $B_j(z)$ are. Thus $f_j^*(z) \equiv \overline{f_j[G(z)]}$ analytically continues $f^*(z)$ into $\Gamma^j \cup D^j$ since $f_j^*(z)$ is continuous up to and on Γ^j . Let $\alpha_{n+1}(z) = (z - z_{n1}^1)(z - z_{n2}^1) \dots (z - z_{nn+1}^1)$

$$\beta_{m+1}(z) = \left(\frac{1}{z - a_2} - \frac{1}{z_{m1}^2} \right) \left(\frac{1}{z - a_2} - \frac{1}{z_{m2}^2} \right) \dots \left(\frac{1}{z - a_2} - \frac{1}{z_{mm+1}^2} \right).$$

Then for z on Γ :

$$p_{nm}(z) = \frac{1}{2\pi i} \int_{\sigma^1} \frac{f(t)}{t - z} \frac{\alpha_{n+1}(t) - \alpha_{n+1}(z)}{\alpha_{n+1}(t)} dt - \frac{1}{2\pi i} \int_{\sigma^2} \frac{f(t)}{t - z} \frac{\beta_{m+1}(t) - \beta_{m+1}(z)}{\beta_{m+1}(t)} dt$$

where $p_{nm}(z)$ is a rational function of $z, p_{nm}(z) = p_n^1(z) + p_{m+1}^2(z)$ in which $p_n^1(z)$ is the polynomial in z of degree $\leq n$ got by interpolating $f(z)$ along Γ^1 at $z_{n1}^1, z_{n2}^1, \dots, z_{nn+1}^1$ and $p_{m+1}^2(z)$ is the polynomial of degree $m + 1$ in $1/(z - a_2)$ got by interpolating $f(z)$ along Γ^2 so that $p_{m+1}^2(a_2 + z_{mj}^2) = f(z_{mj}^2)$. To see the latter let $x = 1/(z - a_2)$ and $y =$

$1/(t - a_2)$ then $\beta_{m+1}(t) = b_{m+1}(y)$ where $b_{m+1}(y)$ is a monic polynomial in y of degree $\leq m + 1$, thus we have

$$\beta_{m+1}(t) - \beta_{m+1}(z) = b_{m+1}(x) - b_{m+1}(y) = (x - y) \sum_{i=1}^m a_i(x)y^i$$

where $a_i(x)$ are polynomials in x of degree $\leq m$. But

$$x - y = \frac{1}{z - a_2} - \frac{1}{t - a_2} = (t - z)xy$$

thus

$$\int_{\sigma^2} \frac{f(t)}{t - z} \frac{\beta_{m+1}(t) - \beta_{m+1}(z)}{\beta_{m+1}(t)} dt$$

is a polynomial of degree $\leq m + 1$ in $1/(z - a_2)$. The error for z on Γ is given by:

$$f(z) - p_{nm}(z) = \frac{1}{2\pi i} \int_{\sigma^1} \frac{f(t)}{t - z} \frac{\alpha_{n+1}(z)}{\alpha_{n+1}(t)} dt - \frac{1}{2\pi i} \int_{\sigma^2} \frac{f(t)}{t - z} \frac{\beta_{m+1}(z)}{\beta_{m+1}(t)} dt .$$

Note that:

$$|\alpha_{n+1}(z)| \leq \mu_n^1, \quad |\alpha_{n+1}(t)| \geq \delta_n^1 \text{ for } z \text{ on } \Gamma^1 \text{ and } t \text{ on } C^1 \text{ and:}$$

$$\frac{\frac{1}{z - a_2} - \frac{1}{z^2_{n1}}}{\frac{1}{t - a_2} - \frac{1}{z^2_{n1}}} = \frac{z - a_2 - z^2_{n1}}{t - a_2 - z^2_{n1}} \frac{t - a_2}{z - a_2}$$

and thus

$$\left| \frac{\beta_{m+1}(z)}{\beta_{m+1}(t)} \right| \leq \frac{\mu_m^2}{\delta_m^2} \text{ for } z \text{ on } \Gamma^2 \text{ and } t \text{ on } C^2.$$

From these it follows:

$$|f(z) - p_{nm}(z)| \leq \frac{1}{2\pi} \left\{ \frac{L_1 M_1}{\delta_1} \frac{\mu_n^1}{\delta_1} + \frac{L_2 M_2}{\delta_2} \frac{\mu_m^2}{\delta_m^2} \right\} \text{ for } z \text{ on } \Gamma,$$

where L_j is the length of C^j ,

$$M_j = \max_{t \text{ on } C^j} |f(t)|, \quad \text{and} \quad \delta_j = \inf_{z \text{ on } \Gamma^j} \min_{t \text{ on } C^j} |t - z|$$

which is the result.

We next consider the case when Γ is a single analytic contour and ($C^j = C$) we write Γ in parametric form as $z(\sigma) = x(\sigma) + iy(\sigma)$ where $-1 \leq \sigma \leq 1$. Let $|z(\sigma_2) - z(\sigma_1)| \leq A|\sigma_2 - \sigma_1|$, let Γ contain the origin and

THEOREM 2 (H. 1) *Let $f(z)$ be an analytic single-valued function on R whose boundary is Γ such that the real part of $f(z)$ solves the Dirichlet problem in R with real boundary values $B(z)$ on Γ where $B(z)$ is a single-valued analytic function on $D \cup \Gamma \cup \hat{D}$ continuous on $\Gamma \cup \hat{D} \cup C$. Let $f(z)$ be continuous and single-valued on $R \cup \Gamma$.*

(H. 2) *Let $z_j^n = z(\sigma_j^n)$ where*

$$\sigma_j^n = \cos [(2j - 1) \pi / (2n + 2)], \quad j = 1, 2, \dots, n + 1$$

(H. 3) $\delta = \inf_{z \text{ on } \Gamma} \min_{t \text{ on } C} |t - z|$

(H. 4) $A < 2\delta$.

Then

(C. 1) $p_n(z) = \sum_{j=1}^{n+1} f(z_j^n) \frac{\omega_{n+1}(z)}{\omega'_{n+1}(z_j^n)(z - z_j^n)}$, where

$\omega_{n+1}(z) = \prod_{k=1}^{n+1} (z - z_k^n)$ and prime denotes differentiation, converges uniformly to $f(z)$ on $R \cup \Gamma$ as $n \rightarrow \infty$

(C. 2) $|f(z) - p_n(z)| \leq \frac{LM}{\pi\delta} \left(\frac{A}{2\delta}\right)^{n+1}$

where M is a constant depending on f , L is length of Γ .

Proof. As in the proof of Theorem 1 we have for z on Γ

$$|f(z) - p_n(z)| \leq \frac{L}{2\pi} M \max_{z \text{ on } \Gamma} |\omega_{n+1}(z)| / \delta^{n+2}.$$

But

$$\begin{aligned} |\omega_{n+1}(z)| &= |(z - z_1^n)(z - z_2^n) \cdots (z - z_{n+1}^n)| \\ &\leq A^{n+1} |(\sigma - \sigma_1^n)(\sigma - \sigma_2^n) \cdots (\sigma - \sigma_{n+1}^n)| \end{aligned}$$

where the σ_j^n are the roots of the Chebyshev polynomial

$$T_{n+1}(\sigma) = \cos [(n + 1) \text{ arc } \cos \sigma]$$

of degree $n + 1$. Thus since $(\sigma - \sigma_1^n)(\sigma - \sigma_2^n) \cdots (\sigma - \sigma_{n+1}^n)$ is monic

$$|\omega_{n+1}(z)| \leq A^{n+1} T_{n+1}(\sigma) / 2^n.$$

Thus

$$\max_{z \text{ on } \Gamma} |\omega_{n+1}(z)| \leq A^{n+1} / 2^n \quad \text{and} \quad |f(z) - p_n(z)| \leq \frac{LM}{\delta\pi} \left(\frac{A}{2\delta}\right)^{n+1}.$$

Next let $\Gamma: z(s) = x(s) + iy(s)$ where s is arc length $0 \leq s \leq L$.

THEOREM 3. (H. 1) *Same as Theorem 2.*

(H. 2) *Same as Theorem 2 but $z_j^n = z(\sigma_j^n)$ where*

$$s_j^n = \frac{L}{2} \cos [(2j - 1)\pi/2(n + 1)] + \frac{L}{2}, \quad j = 1, 2, \dots, n + 1.$$

(H 3) $\delta = \inf_{z \text{ on } \Gamma} \min_{t \text{ on } \mathcal{O}} |t - z|.$

(H 4) $L < 4\delta.$

(C 1) *Same as Theorem 2.*

(C 2) $|f(z) - p_n(z)| \leq \frac{LM}{\pi\delta} \left(\frac{L}{4\delta}\right)^{n+1}$

where M is a constant depending on f .

Proof. As in the proof of theorem for z on Γ :

$$|f(z) - p_n(z)| \leq \frac{L}{2\pi} M \max |\omega_{n+1}(z)| / \delta^{n+2}.$$

But since $|z - z_j^n| \leq |s - s_j^n|$ where $z = z(s)$ and $z_j^n = z(s_j^n)$ and since $|(s - s_1^n)(s - s_2^n) \dots (s - s_{n+1}^n)| \leq L^{n+1}/2^{2n+1}$ see e.g. [1] we have

$$|f(z) - p_n(z)| \leq \frac{LM}{\pi\delta} \left(\frac{L}{4\delta}\right)^{n+1}.$$

EXAMPLE. We shall now apply the ideas of this paper to a particular geometrical configuration. Let

Γ^1 be a circle of radius 15 centered at the origin

Γ^2 be a circle of radius 1 centered at $(-13, 0)$

Γ^3 be an ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, \quad a = 1.075, \quad b = 1.$$




Fig. 1.

Let R be the interior of Γ^1 less Γ^2, Γ^3 and the interiors of Γ^2 and Γ^3 . Let $f(z)$ be analytic on R and continuous on $R \cup \Gamma^1 \cup \Gamma^2 \cup \Gamma^3$. Moreover let the real part of $f(z)$ satisfy boundary conditions $B_1(z)$ on Γ^1 , $B_2(z)$ on Γ^2 and $B_3(z)$ on Γ^3 where:

$$B_1(z) \text{ is analytic on } |z| = 15$$

$$B_2(z) \text{ is analytic on } |z + 13| = 1$$

and $B_3(z)$ is analytic in and on $\Gamma^3 - \{-.395 < x < .395, y = 0\}$ See figure.

For example we might have $Re f(z) = P_k(x, y)$ on Γ^k ($k = 1, 2, 3$) where $P_k(x, y)$ is a polynomial.

Then since:

$$G_1(z) = (15)^2 z^{-1}$$

$$G_2(z) = (z + 13)^{-1} - 13$$

$$(E. 1) \quad G_3(z) = \frac{z(\alpha^2 + b^2) \pm 2ab\sqrt{z^2 + b^2 - \alpha^2}}{\alpha^2 - b^2} \quad a > b \text{ see [3]}$$

and $z = \hat{z} = \overline{G_k(z)}$ on Γ_k we have on Γ_k

$$P_k\left(\frac{z + \bar{z}}{2}, \frac{z - \bar{z}}{2i}\right) = P_k\left[\frac{z + G_k(z)}{2}, \frac{z - G_k(z)}{2i}\right] \equiv B_k(z)$$

which are meromorphic functions that fulfill the requirements of $B_1(z), B_2(z)$ and $B_3(z)$ (in the case of $B_3(z)$ we make a cut between the foci $\pm\sqrt{\alpha^2 - b^2}$).

Let

$$r_k = 15 \exp\left(\frac{2\pi i}{n+1} k\right) \quad \text{and} \quad \alpha_{n+1}(z) = (z - r_1)(z - r_2) \cdots (z - r_{n+1})$$

and

$$p_n(z) = \sum_{k=1}^{n+1} f(r_k) \frac{\alpha_{n+1}(z)}{\alpha_{n+1}'(r_k)(z - r_k)}$$

where the prime signifies differentiation. $p_n(z)$ is clearly the polynomial of degree $\leq n$ that interpolates $f(z)$ at $z = r_k$ on $\Gamma^1, k = 1, \dots, n+1$. Next let

$$s_k = \exp\left(\frac{2\pi i}{m+1} k\right)$$

and

$$\beta_{m+1}(z) = \left(\frac{1}{z + 13} - \frac{1}{s_1}\right)\left(\frac{1}{z + 13} - \frac{1}{s_2}\right) \cdots \left(\frac{1}{z + 13} - \frac{1}{s_{m+1}}\right)$$

and

$$q_m(z) = \sum_{k=1}^{m+1} f(s_k - 13) \frac{B_{m+1}(z)}{B'_{m+1}(s_k - 13) \left(\frac{1}{s_k} - \frac{1}{z + 13} \right) (s_k)^2}$$

$q_m(z)$ is a polynomial of degree $\leq m$ in $1/(z + 13)$ such that $q_m(s_k - 13) = f(s_k - 13)$ where $s_k - 13$ is on Γ^2 , $k = 1, 2, \dots, m + 1$.

Finally let l be the length of the ellipse Γ^3 and

$$\sigma_k = \cos [(2k - 1)\pi/2(j + 1)], k = 1, 2, \dots, j + 1.$$

Then the ellipse Γ^3 can be written

$$z(\sigma) = x(\sigma) + iy(\sigma) = a \cos (2\pi\sigma/l) + ib \sin (2\pi\sigma/l), -l/2 \leq \sigma \leq l/2$$

σ is arc length parameter shifted. Let

$$t_k = z(\sigma_k l/2) \quad \text{and} \quad \kappa_{j+1}(z) = \left(\frac{1}{z} - \frac{1}{t_1} \right) \left(\frac{1}{z} - \frac{1}{t_2} \right) \dots \left(\frac{1}{z} - \frac{1}{t_{j+1}} \right)$$

and

$$r_j(z) = \sum_{k=1}^{j+1} f(t_k) \frac{\kappa_{j+1}(z)}{\kappa'_{j+1}(t_k) \left(\frac{1}{t_k} - \frac{1}{z} \right) (t_k)^2}$$

$r_j(z)$ is clearly the polynomial in $1/z$ of degree $\leq j$ such that $r_j(t_k) = f(t_k)$ $k = 1, 2, \dots, j + 1$ where t_k is on Γ^3 .

Then the assertion is

$$p_n(z) + q_m(z) + r_j(z)$$

converges uniformly to $f(z)$ on $R \cup \Gamma^1 \cup \Gamma^2 \cup \Gamma^3$ as

$$\frac{1}{n} + \frac{1}{m} + \frac{1}{j} \rightarrow 0.$$

For Γ^1 , we use Runge's theorem. Since $B_1(z)$ is analytic on Γ^1 , then $f(z)$ can be continued across Γ^1 , i.e. $f(z)$ is analytic for $15 \leq |z| \leq 15 + \epsilon$ where ϵ is some positive number. Thus in the notation of the theorem

$$\begin{aligned} \delta_n^1 &= \min_{|t|=15+\epsilon} \left| \prod_{k=1}^{n+1} (t - r_k) \right| = \min_{|t|=15+\epsilon} |t^{n+1} - 15^{n+1}| \\ &= \min_{|\tau|=1+\epsilon/15} 15^{n+1} |\tau^{n+1} - 1| \geq 15^{n+1} \min_{|\tau|=1+\epsilon/15} \{|\tau|^{n+1} - 1\} \\ &= 15^{n+1} \{[1 + \epsilon/15]^{n+1} - 1\} \end{aligned}$$

$$\mu_n^1 = \max_{|z|=15} \left| \prod_{k=1}^{n+1} (z - r_k) \right| = 15^{n+1} \max_{|\zeta|=1} |\zeta^{n+1} - 1| \leq 2 \cdot 15^{n+1}$$

and

(E 2) $u_n^1/\delta_n^1 \leq 2/\{[1 + \varepsilon/15]^{n+1} - 1\} \rightarrow 0$ as $n \rightarrow \infty$.

For Γ^2 , since $B_2(z)$ is analytic on Γ^2 , then $f(z)$ can be continued across Γ^2 , i.e. $f(z)$ is analytic for $1 - \varepsilon \leq |z + 13| \leq 1$ where ε is some positive number. Thus if $C^2 = \{z: |z + 13| = 1 - \varepsilon\}$ we have:

$$\begin{aligned} \max_{z \text{ on } \Gamma^2} |\beta_{m+1}(z)| &= \max_{z \text{ on } \Gamma^2} \left| \prod_{k=1}^{m+1} \left(\frac{1}{z + 13} - \frac{1}{s_k} \right) \right| = \max_{z \text{ on } \Gamma^2} \left| \left(\frac{1}{z + 13} \right)^{m+1} - 1 \right| \\ &= \max_{|\zeta|=1} \left| \left(\frac{1}{\zeta} \right)^{m+1} - 1 \right| \leq 2 \end{aligned}$$

and

$$\begin{aligned} \min_{t \text{ on } \sigma^2} |\beta_{m+1}(t)| &= \min_{t \text{ on } \sigma^2} \left| \left(\frac{1}{t + 13} \right)^{m+1} - 1 \right| \\ &= \min_{|\zeta|=1-\varepsilon} \left| \left(\frac{1}{\zeta} \right)^{m+1} - 1 \right| \geq \left(\frac{1}{1-\varepsilon} \right)^{m+1} - 1. \end{aligned}$$

From these we see that

$$(E. 3) \quad \frac{\max_{z \text{ on } \Gamma^2} |\beta_{m+1}(z)|}{\min_{t \text{ on } \sigma^2} |\beta_{m+1}(t)|} \leq \frac{2}{\left(\frac{1}{1-\varepsilon} \right)^{m+1} - 1} \rightarrow 0 \text{ as } m \rightarrow \infty,$$

For Γ^3 we note from the reflection function $G(z)$ given by (E. 1) that the interior of the ellipse Γ^3 minus the line $-c \leq x \leq c$, $c^2 = a^2 - b^2$, is reflected exterior to the given ellipse but interior to the ellipse e_2

$$\frac{\hat{x}^2}{\hat{a}^2} + \frac{\hat{y}^2}{\hat{b}^2} = 1$$

where $\hat{a} = (a^2 + b^2)/c$, $\hat{b} = 2ab/c$.

In the case of our ellipse we have $a = 1.075$, $b = 1$ and $c = .395$, $\hat{a} = 5.46$, $\hat{b} = 5.44$, thus e_2 is contained in Γ^1 , and does not intersect or contain points of Γ^2 and thus $f(z)$ can be extended to be analytic in $\Gamma^3 - \{z | -.395 < x < .395, y = 0\}$.

The length of the ellipse Γ^3 is given by:

$$l = 4a \int_0^{\pi/2} \sqrt{1 - k^2 \sin^2 \theta} d\theta$$

where

$$\begin{aligned} k &= c/a < 1. \text{ In our case } k = .368 \text{ and thus} \\ l &= 4a (1.516) \end{aligned}$$

using a table for elliptic integrals. Let c^3 be the rectangular contour

$$(-.395 - \varepsilon \leq x \leq .395 + \varepsilon, y = -\varepsilon), (x = .395 + \varepsilon, -\varepsilon \leq y \leq \varepsilon), \\ (-.395 - \varepsilon \leq x \leq .395 + \varepsilon, y = \varepsilon), (x = -.395 - \varepsilon, -\varepsilon \leq y \leq \varepsilon)$$

where $\varepsilon > 0$ is arbitrarily small. Then consider

$$\frac{1}{2\pi i} \int_{\sigma^3} \frac{f(t)}{t-z} \frac{\kappa_{j+1}(z)}{\kappa_{j+1}(t)} dt.$$

But

$$|\kappa_{j+1}(z)| = \left| \prod_{k=1}^{j+1} \left(\frac{1}{z} - \frac{1}{t_k} \right) \right| = \left| \prod_{k=1}^{j+1} \frac{t_k - z}{t_k z} \right| \\ \leq \prod_{k=1}^{j+1} \frac{\sigma - \sigma_k l/2}{|t_k| |z|} \leq \left(\frac{l}{2} \right)^{j+1} \prod_{k=1}^{j+1} \frac{|\theta - \sigma_k|}{|t_k|}. \quad \sigma = \text{arc length},$$

where $-1 \leq \theta \leq 1$ since $|z| \geq 1$ for z on Γ^3 . Also for t on C^3

$$|\kappa_{j+1}(t)| = \left| \prod_{k=1}^{j+1} \frac{t_k - t}{t_k t} \right| \geq (a - c - \eta\varepsilon)^{j+1} \prod_{k=1}^{j+1} \frac{1}{|t_k| |t|}$$

where η is some fixed constant. But since $|t| \leq c + \varepsilon/2$ for t on c^3 we see that

$$|\kappa_{j+1}(t)| \geq \left(\frac{a - c - \eta\varepsilon}{c + \varepsilon\sqrt{2}} \right)^{j+1} \prod_{k=1}^{j+1} \frac{1}{|t_k|}.$$

Combining the above results gives

$$A \equiv |\kappa_{j+1}(z)/\kappa_{j+1}(t)| \leq \left(\frac{c + \varepsilon\sqrt{2}}{a - c - \eta\varepsilon} \right)^{j+1} \left(\frac{l}{2} \right)^{j+1} 2^{-j} |T_{j+1}(\theta)|$$

where we have utilized the fact that the σ_k are the roots of the Chebyshev polynomial $T_{j+1}(\theta) = \text{cox} [(j+1) \text{ are } \cos \theta]$. Thus

$$A \leq 2 \left(\frac{c + \varepsilon\sqrt{2}}{a - c - \eta\varepsilon} \right)^{j+1} \left(\frac{l}{4} \right)^{j+1}.$$

But

$$\frac{l}{4a} = 1.516 < \frac{1}{a} \frac{a - c - \eta\varepsilon}{c + \varepsilon\sqrt{2}} = \frac{1}{1.075} \frac{.680 - \eta\varepsilon}{.395 + \sqrt{2}\varepsilon}, = 1.60 + g(\varepsilon)$$

where $g(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$. Thus for ε sufficiently small

$$(E. 4) \quad l/4 \leq \frac{a - c - \eta\varepsilon}{c + \varepsilon\sqrt{2}}.$$

Utilizing (E. 2), (E. 3) and (E. 4) we have from Theorem 1 that $p_n(z) + p_m(z) + r_j(z)$ converges uniformly to $f(z)$ in $R \cup \Gamma^1 \cup \Gamma^2 \cup \Gamma^3$ as $(1/n) + (1/m) + (1/j) \rightarrow 0$.

We remark finally that there would be no new difficulties if Γ had contained in addition $\Gamma^4 \cup \Gamma^5 \cup \Gamma^6$ where Γ^4 is the circle $|z - 10i| = 4$, Γ^5 the circle $|z + 10i| = 4$ and Γ^6 is the circle $|z - 12| = 2$.

REFERENCES

1. T. M. Apostol, *Calculus*, Vol. II, Blaisdell Pub. Co.
2. S. N. Mergelyan, *Uniform approximations to functions of a complex variable*, Trans. Ser. 1, Vol. 3, Series and Approximation, Amer. Math. Soc.
3. J. M. Sloss, *Reflection of Biharmonic Functions Across Analytic Boundary Conditions with Examples*, Pacific. J. Math. **13** (1963), 1401-1415.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California

R. M. BLUMENTHAL

University of Washington
Seattle, Washington 98105

*J. DUGUNDJI

University of Southern California
Los Angeles, California 90007

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

* Paul A. White, Acting Editor until J. Dugundji returns.

Gert Einar Torsten Almkvist, <i>Stability of linear differential equations with periodic coefficients in Hilbert space</i>	383
Richard Allen Askey and Stephen Wainger, <i>A transplantation theorem for ultraspherical coefficients</i>	393
Joseph Barback, <i>Two notes on regressive isols</i>	407
Allen Richard Bernstein and Abraham Robinson, <i>Solution of an invariant subspace problem of K. T. Smith and P. R. Halmos</i>	421
P. R. Halmos, <i>Invariant subspaces of polynomially compact operators</i>	433
Leon Bernstein, <i>New infinite classes of periodic Jacobi-Perron algorithms</i>	439
Richard Anthony Brualdi, <i>Permanent of the direct product of matrices</i>	471
W. Wistar (William) Comfort and Kenneth Allen Ross, <i>Pseudocompactness and uniform continuity in topological groups</i>	483
James Michael Gardner Fell, <i>Algebras and fiber bundles</i>	497
Alessandro Figà-Talamanca and Daniel Rider, <i>A theorem of Littlewood and lacunary series for compact groups</i>	505
David London, <i>Two inequalities in nonnegative symmetric matrices</i>	515
Norman Jay Pullman, <i>Infinite products of substochastic matrices</i>	537
James McLean Sloss, <i>Reflection and approximation by interpolation along the boundary for analytic functions</i>	545
Carl Weinbaum, <i>Visualizing the word problem, with an application to sixth groups</i>	557