

# Pacific Journal of Mathematics

**ON THE NONSINGULARITY OF COMPLEX MATRICES**

PAUL CAMION AND ALAN JEROME HOFFMAN

## ON THE NONSINGULARITY OF COMPLEX MATRICES

PAUL CAMION\* AND A. J. HOFFMAN\*\*

**Let  $A = (a_{ij})$  be a real square matrix of order  $n$  with nonnegative entries, and let  $M(A)$  be the class of all complex matrices  $B = (b_{ij})$  of order  $n$  such that, for all  $i, j$ ,  $|b_{ij}| = a_{ij}$ . If every matrix in  $M(A)$  is nonsingular, we say  $M(A)$  is regular, and it is the purpose of this note to investigate conditions under which  $M(A)$  is regular.**

Many sufficient conditions have been discovered (cf., for instance, [8] and [3], and their bibliographies), motivated by the fact that the negation of these conditions, applied to the matrix  $B - \lambda I$ , yields information about the location of the characteristic roots. We shall show that a mild generalization of the most famous conditions [2] is not only sufficient but also necessary. (The application of our result to characteristic roots will not be discussed here, but is contained in [5]. See also [7] and [9]).

If

$$(1.1) \quad a_{ii} > \sum_{i \neq j} a_{ij}, \quad i = 1, \dots, n,$$

then ([2])  $M(A)$  is regular. Clearly if  $P$  is a permutation matrix, and  $D$  a diagonal matrix with positive diagonal entries, such that  $PAD$  satisfies (1.1), then  $M(A)$  is regular. We shall show that, conversely, if  $M(A)$  is regular, there exist such matrices  $P$  and  $D$  so that (1.1) holds.

**2. Notation and lemmas.** If  $x = (x_1, \dots, x_n)$  is a vector,  $x^D$  is the diagonal matrix whose  $i$ th diagonal entry is  $x_i$ . If  $M = (m_{ij})$  is a matrix,  $M^v$  is the vector whose  $i$ th coordinate is  $m_{ii}$ . A vector  $x = (x_1, \dots, x_n)$  is positive if each  $x_j > 0$ ;  $x$  is semi-positive if  $x \neq 0$  and each  $x_j \geq 0$ . A diagonal matrix  $D$  is positive (semipositive) if  $D^v$  is positive (semi-positive). If  $A = (a_{ij})$  is a matrix with nonnegative entries, a particular entry  $a_{ij}$  is said to be dominant in its column if

$$a_{ij} > \sum_{k \neq i} a_{kj}.$$

**LEMMA 1.** *If  $e_1, \dots, e_n$  are nonnegative numbers such that the largest does not exceed the sum of the others, then there exist complex numbers  $z_i$  such that*

---

Received June 29, 1964. \*Euratom

\*\* The work of this author was supported in part by the Office of Naval Research under Contract No. Nonr 3775(00), NR 047040.

$$(2.1) \quad |z_i| = e_i \quad i = 1, \dots, n$$

and

$$(2.2) \quad \Sigma z_i = 0 .$$

*Proof.* It is geometrically obvious (and can easily be proved by induction) that the conditions on  $\{e_i\}$  imply there exists a (possibly degenerate) polygon in the complex plane whose successive sides have length  $e_1, e_2, \dots, e_n$ . Let the vertices  $x_1, \dots, x_n$  be so numbered that  $|x_i - x_{i+1}| = e_i, i = 1, \dots, n-1, |x_n - x_1| = e_n$ . Setting  $z_i = x_i - x_{i+1}$  obviously satisfies (2.1) and (2.2).

LEMMA 2. *Let  $M$  be a real matrix with  $m$  rows and  $n$  columns. Then*

$$(2.3) \quad Mx \leq 0, \quad x \text{ semi-positive}$$

*is inconsistent if and only if*

$$(2.4) \quad w'M > 0, \quad w \geq 0$$

*is consistent. Further, if (2.4) holds, we may assume there exists a  $w$  satisfying (2.4) with at most  $n$  coordinates of  $w$  positive.*

This lemma is well known in the theory of linear inequalities.

3. THEOREM. *Let  $A = (a_{ij})$  be a matrix of order  $n$  with each entry nonnegative. The following statements are equivalent:*

(3.1)  *$M(A)$  is regular:*

(3.2) *if  $D$  is any semi-positive diagonal matrix, then  $DA$  contains an entry dominant in its column;*

(3.3) *there exists a permutation matrix  $P$  and a positive diagonal matrix  $D$  such that  $PAD$  satisfies (1.1).*

*Proof.* (3.1)  $\Rightarrow$  (3.2). Assume (3.2) false for some semipositive  $D$  with  $D^v = (d_1, \dots, d_n)$ . Let  $(a_{1j}d_1, \dots, a_{nj}d_n)$  be any column vector of  $DA$ . The coordinates of this vector satisfy the hypotheses of Lemma 1, so there exist complex numbers  $z_1, \dots, z_n$  satisfying

$$(3.4) \quad \Sigma z_i = 0 ,$$

and

$$(3.5) \quad |z_i| = a_{ij}d_i, \quad i = 1, \dots, n .$$

Let

$$b_{ij} = a_{ij}z_i/|z_i| ,$$

with  $z_i/|z_i| = 1$ , if  $z_i = 0$ . Then (3.4) and (3.5) become

$$(3.6) \quad \sum d_i b_{ij} = 0 ,$$

and

$$(3.7) \quad |b_{ij}| = a_{ij} , \quad i, j = 1, \dots, n .$$

But (3.7) states  $B \in M(A)$ , and (3.6)—since not all  $d_i$  are 0—asserts a linear dependence among the rows of  $B$ . Thus  $B \in M(A)$  would be singular, violating (3.1).

(3.2)  $\Rightarrow$  (3.3). Let  $K$  be a matrix of order  $n$  with  $k_{ii} = 1, k_{ij} = -1$  for  $i \neq j$ , and let  $A_j$  be the  $j$ 'th column of  $A$ . Consider the system of  $n^2$  linear inequalities in the semi-positive vector  $x$

$$(3.7) \quad KA_j^p x \leq 0 , \quad j = 1, \dots, n .$$

Notice that (3.2) is identical with the statement that (3.7) is inconsistent. By Lemma 2, there exist  $n$  nonnegative vectors  $\mu^1, \dots, \mu^n$  such

$$(3.8) \quad \sum_j \mu^j KA_j^p > 0 .$$

Let  $\mu^j = (\mu_1^j, \dots, \mu_n^j)$ . By the last sentence of Lemma 2, we may assume at most  $n$  of the  $n^2$  numbers  $\{\mu_k^j\}$  are positive.

Since each row of each  $KA_j^p$  contains at most one positive entry, it follows from (3.8) that exactly  $n$  of the  $\{\mu_k^j\}$  are positive. We now show that, for each  $j$ , there is exactly one  $k$  such that  $\mu_k^j > 0$ . Assume otherwise, then for (say)  $j = j^*, \mu^{j^*} = 0$ . Let  $\tilde{A}$  be the matrix obtained from  $A$  by replacing  $A_{j^*}$  by 0. Then (3.8) would still hold with  $A_j$  replaced by 0, so (from the "only if" part of Lemma 2), for any semi-positive diagonal matrix  $E, E\tilde{A}$  contains an entry dominant in its column. Let  $y$  be a real nonzero vector orthogonal to the columns of  $\tilde{A}$ , let  $N = \{i \mid y_i \geq 0\}$ , and  $N'$  the complementary set of indices. Then, for each  $j$ .

$$(3.9) \quad \sum_{i \in N} y_i a_{ij} = \sum_{i \in N'} (-y_i) a_{ij} .$$

If  $E$  is the diagonal matrix with  $E^r = (|y_1|, \dots, |y_n|)$ , then  $E\tilde{A}$ , from (3.9), would contain no entry dominant in its column, a contradiction.

Let  $\sigma$  be the mapping sending  $j \rightarrow k$ , where  $\mu_k^j > 0$ . By (3.8),  $\sigma$  is a permutation of  $\{1, \dots, n\}$ , and

$$a_{i, \sigma^{-1}i} \mu_i^{\sigma^{-1}i} > \sum_{j \neq i} a_{i, \sigma^{-1}j} \mu_j^{\sigma^{-1}j} , \quad i = 1, \dots, n ,$$

which is (3.3).

(3.3)  $\Rightarrow$  (3.1) was noted in the introduction.

4. **Remarks.** (i) It is perhaps worth pointing out that the permutation in (3.3) is unique. For, without loss of generality, assume  $P$  and  $D$  both the identity matrix, so that (1.1) holds. Assume  $Q$  and  $E$  given so that  $QAE$  satisfies (1.1). If  $Q$  is not the identity permutation, then there must exist some cycle such that (say)

$$(4.1) \quad \begin{aligned} a_{q_1 q_2} e_{q_2} &> a_{q_1 q_1} e_{q_1} \\ &\cdot \quad \cdot \quad \cdot \\ a_{q_{r-1} q_r} e_{q_r} &> a_{q_{r-1} q_{r-1}} e_{q_{r-1}} \\ a_{q_r q_1} e_{q_1} &> a_{q_r q_r} e_{q_r} . \end{aligned}$$

Multiplying the inequalities (4.1) together, we obtain

$$a_{q_1 q_2} \cdots a_{q_r q_1} > a_{q_1 q_1} \cdots a_{q_r q_r} ,$$

which violates (1.1).

In fact, it is clear from the foregoing that the diagonal entries in the  $PAD$  of (3.3) will be that collection of  $n$  entries of  $A$ , one from each row and column, whose product is a maximum. Further, that collection is necessarily unique. Finding the collection amounts to solving the assignment problem of linear programming [1] where the "scores" are  $\{\log a_{ij}\}$ . In some cases this can be done easily ([4]), but not in general [6].

(ii) If we had confined our attention to real rather than complex matrices, our theorem does not apply, and the problem seems difficult. With somewhat stronger hypotheses than the real case of (3.1), the problem has been solved by Ky Fan [3].

#### REFERENCES

1. G. B. Dantzig, *Linear Programming and Extensions*, Princeton University Press, Princeton, 1963.
2. J. Desplanques, *Théorème d'Algèbre*, J. Math. Spec. (3) **1** (1887), 12-13.
3. Ky Fan, *Note on circular disks containing the eigenvalues of a matrix*, Duke Math J. **25** (1958), 441-445.
4. A. J. Hoffman, *On Simple Linear Programming Problems*, Proceeding of Symposia in Pure Mathematics, Vol. III, Convexity, 1963, American Mathematical Society, pp. 317-327.
5. B. W. Levinger and R. S. Varga, *Minimal Gerschgorin sets II*, to be published.
6. T. S. Motzkin, *The Assignment Problem*, Proceedings of Symposia in Applied Mathematics, Vol. VI, Numerical Analysis, 1956, American Mathematical Society, pp. 109-125.
7. Hans Schneider, *Regions of exclusion for the latent roots of a matrix*, Proc. American Math. Soc. **5** (1954), 320-322.
8. Olga Taussky, *A recurring theorem on determinants*, Amer. Math. Monthly **56** (1949), 672-676.
9. R. S. Varga, *Minimal Gerschgorin sets*, to appear in Pacific J. Math.

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON

Stanford University  
Stanford, California

R. M. BLUMENTHAL

University of Washington  
Seattle, Washington 98105

\*J. DUGUNDJI

University of Southern California  
Los Angeles, California 90007

RICHARD ARENS

University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. Yosida

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL ORDNANCE TEST STATION

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

\* Paul A. White, Acting Editor until J. Dugundji returns.

# Pacific Journal of Mathematics

Vol. 17, No. 2

February, 1966

Henry A. Antosiewicz, <i>Boundary value problems for nonlinear ordinary differential equations</i> .....	191
Bernard Werner Levinger and Richard Steven Varga, <i>Minimal Gerschgorin sets. II</i> .....	199
Paul Camion and Alan Jerome Hoffman, <i>On the nonsingularity of complex matrices</i> .....	211
J. Chidambaraswamy, <i>Divisibility properties of certain factorials</i> .....	215
J. Chidambaraswamy, <i>A problem complementary to a problem of Erdős</i> ....	227
John Dauns, <i>Chains of modules with completely reducible quotients</i> .....	235
Wallace E. Johnson, <i>Existence of half-trajectories in prescribed regions and asymptotic orbital stability</i> .....	243
Victor Klee, <i>Paths on polyhedra. II</i> .....	249
Edwin Haena Mookini, <i>Sufficient conditions for an optimal control problem in the calculus of variations</i> .....	263
Zane Clinton Motteler, <i>Existence theorems for certain quasi-linear elliptic equations</i> .....	279
David Lewis Outcalt, <i>Simple <math>n</math>-associative rings</i> .....	301
David Joseph Rodabaugh, <i>Some new results on simple algebras</i> .....	311
Oscar S. Rothaus, <i>Asymptotic properties of groups generation</i> .....	319
Ernest Edward Shult, <i>Nilpotence of the commutator subgroup in groups admitting fixed point free operator groups</i> .....	323
William Hall Sills, <i>On absolutely continuous functions and the well-bounded operator</i> .....	349
Joseph Gail Stampfli, <i>Which weighted shifts are subnormal</i> .....	367
Donald Reginald Traylor, <i>Metrizability and completeness in normal Moore spaces</i> .....	381