

Pacific Journal of Mathematics

**THE SUM OF TWO INDEPENDENT EXPONENTIAL-TYPE
RANDOM VARIABLES**

EDWARD MARTIN BOLGER

THE SUM OF TWO INDEPENDENT EXPONENTIAL-TYPE RANDOM VARIABLES

E. M. BOLGER

Let X_1, X_2 be nondegenerate, independent, exponential-type random variables (r.v.) with probability density functions, (p.d.f.) $f_1(x_1; \theta), f_2(x_2; \theta)$, (not necessarily with respect to the same measure), where $f_i(x_i; \theta) = \exp\{x_i p_i(\theta) + q_i(\theta)\}$ for $\theta \in (a, b)$ and $p_i(\theta)$ is an analytic function of θ (for $\operatorname{Re} \theta \in (a, b)$) with $p_i'(\theta)$ never equal to zero on (a, b) . If X_1, X_2 are neither both normal nor both Poisson type r.v.'s, then $X_1 + X_2$ is an exponential-type r.v. if and only if $p_1'(\theta) = p_2'(\theta)$.

2. Lemmas. It follows from Patil's result ([3]) that a r.v. X is of exponential type if and only if the cumulants, $\lambda_j(\theta)$, exist and satisfy

$$(1) \quad \lambda_j'(\theta) = p'(\theta)\lambda_{j+1}(\theta) \quad \text{for } j = 1, 2, 3, \dots$$

Lehmann ([2], p. 52) has shown that $q(\theta)$ and hence also $\lambda_j(\theta)$ are analytic functions of $p(\theta)$. Then $\lambda_j(\theta)$ is an analytic function of θ for $\operatorname{Re} \theta \in (a, b)$.

Let $\lambda_{j,i}(\theta)$ be the j^{th} cumulant of X_i and $\lambda_j(\theta)$ the j^{th} cumulant of Y . Then

$$(2) \quad \lambda_j(\theta) = \lambda_{j,1}(\theta) + \lambda_{j,2}(\theta)$$

$$(3) \quad \lambda_{j,i}'(\theta) = p_i'(\theta)\lambda_{j+1,i}(\theta) \quad \text{for } j = 1, 2, 3, \dots$$

Let $h_j(\theta) = \lambda_{j,1}(\theta)\lambda_{j,2}(\theta) - \lambda_{j,2}(\theta)\lambda_{j,1}(\theta)$ and $c(\theta) \equiv \lambda_{2,2}(\theta)/\lambda_{2,1}(\theta)$.

LEMMA 1. *If $h_3(\theta) \equiv 0$ and if $c'(\theta) \equiv 0$, then either X_1 and X_2 are both normal or $p_1'(\theta) \equiv p_2'(\theta)$.*

Proof. Since $h_3(\theta) \equiv 0$,

$$(4) \quad \lambda_{3,2}(\theta) = c(\theta)\lambda_{3,1}(\theta).$$

Since $c'(\theta) \equiv 0$,

$$(5) \quad \lambda_{2,2}'(\theta) = c(\theta)\lambda_{2,1}'(\theta).$$

From (3), (4) and (5) it follows that

$$p_2'(\theta)\lambda_{3,2}(\theta) = c(\theta)p_1'(\theta)\lambda_{3,1}(\theta) = p_1'(\theta)\lambda_{3,2}(\theta).$$

If $\lambda_{3,2}(\theta) \equiv 0$, then $\lambda_{3,1}(\theta) \equiv 0$ and X_1, X_2 are both normal. If there is a point θ_0 such that $\lambda_{3,2}(\theta) \neq 0$, then there is a neighborhood, $N(\theta_0)$, in which $\lambda_{3,2}(\theta) \neq 0$. For $\theta \in N(\theta_0)$, $p'_1(\theta) = p'_2(\theta)$. By analyticity, $p'_1(\theta) = p'_2(\theta)$ for $\theta \in (a, b)$.

LEMMA 2. *If $h_j(\theta) \equiv 0$ for $j > 2$ and if $c'(\theta) \neq 0$, then X_1 and X_2 are Poisson type r.v.'s.*

Proof. Since $h_j(\theta) \equiv 0$,

$$(6) \quad \lambda_{j,2}(\theta) = c(\theta)\lambda_{j,1}(\theta).$$

Differentiating (6) and using (3), we get

$$c(\theta)\lambda'_{j,1}(\theta) + c'(\theta)\lambda_{j,1}(\theta) = p'_2(\theta)\lambda_{j+1,2}(\theta).$$

Then,

$$(7) \quad c(\theta)p'_1(\theta)\lambda_{j+1,1}(\theta) + c'(\theta)\lambda_{j,1}(\theta) = p'_2(\theta)c(\theta)\lambda_{j+1,1}(\theta).$$

In particular,

$$(8) \quad c(\theta)p'_1(\theta)\lambda_{3,1}(\theta) + c'(\theta)\lambda_{2,1}(\theta) = p'_2(\theta)c(\theta)\lambda_{3,1}(\theta).$$

Multiplying (7) by $\lambda_{3,1}(\theta)$ and (8) by $\lambda_{j+1,1}(\theta)$, we find that

$$(9) \quad c'(\theta)[\lambda_{2,1}(\theta)\lambda_{j+1,1}(\theta) - \lambda_{3,1}(\theta)\lambda_{j,1}(\theta)] = 0 \quad \text{for } j \geq 2.$$

Since $c'(\theta) \neq 0$, there is a sub-interval M of (a, b) in which $c'(\theta) \neq 0$. For $\theta \in M$,

$$\lambda_{2,1}(\theta)\lambda_{j+1,1}(\theta) - \lambda_{3,1}(\theta)\lambda_{j,1}(\theta) = 0,$$

or

$$(10) \quad \lambda_{j+1,1}(\theta) = \frac{\lambda_{3,1}(\theta)}{\lambda_{2,1}(\theta)}\lambda_{j,1}(\theta).$$

By analyticity, (10) is true for all $\theta \in (a, b)$. Now let $a(\theta) = \lambda_{3,1}(\theta)/\lambda_{2,1}(\theta)$. Then, by (3),

$$\begin{aligned} p'_1(\theta)\lambda_{4,1}(\theta) &= \lambda'_{3,1}(\theta) = a'(\theta)\lambda_{2,1}(\theta) + a(\theta)\lambda'_{2,1}(\theta) \\ &= a'(\theta)\lambda_{2,1}(\theta) + a(\theta)p'_1(\theta)\lambda_{3,1}(\theta). \end{aligned}$$

Since $\lambda_{4,1}(\theta) = a(\theta)\lambda_{3,1}(\theta)$, it follows that

$$a'(\theta)\lambda_{2,1}(\theta) = 0.$$

So $a'(\theta) = 0$ and $a(\theta) = d$. Then (10) becomes

$$(11) \quad \lambda_{j+1,1}(\theta) = d\lambda_{j,1}(\theta) \quad \text{for } j \geq 2.$$

This implies

$$(12) \quad \lambda_{j,1}(\theta) = d^{j-2}\lambda_{2,1}(\theta) \quad \text{for } j \geq 2 .$$

By (6),

$$(13) \quad \lambda_{j,2}(\theta) = d^{j-2}c(\theta)\lambda_{2,1}(\theta) \quad \text{for } j \geq 2 .$$

Now,

$$\begin{aligned} p_1'(\theta) &= \lambda_{1,1}'(\theta)/\lambda_{2,1}(\theta) , \\ p_1'(\theta) &= \lambda_{2,1}'(\theta)/\lambda_{3,1}(\theta) = \lambda_{2,1}'(\theta)/d\lambda_{2,1}(\theta) . \end{aligned}$$

So

$$(14) \quad \lambda_{1,1}(\theta) = d^{-1}\lambda_{2,1}(\theta) + k_1 .$$

Similarly,

$$(15) \quad \lambda_{1,2}(\theta) = d^{-1}c(\theta)\lambda_{2,1}(\theta) + k_2 .$$

Using (12), (13), (14) and (15), we find that

$$\begin{aligned} \log M_1(t; \theta) &= k_1 t + d^{-2}\lambda_{2,1}(\theta)(e^{dt} - 1) \\ \log M_2(t; \theta) &= k_2 t + d^{-2}c(\theta)\lambda_{2,1}(\theta)(e^{dt} - 1) , \end{aligned}$$

where $M_i(t; \theta)$ is the moment generating function corresponding to $f_i(x_i; \theta)$.

This concludes the proof of Lemma 2.

3. The sum of two independent exponential-type random variables.

THEOREM 1. *If X_1, X_2 are neither both normal nor both Poisson type r.v.'s, then $X_1 + X_2$ is an exponential-type r.v. if and only if $p_1'(\theta) = p_2'(\theta)$.*

Proof. If $p_1'(\theta) = p_2'(\theta)$, then it follows from (2) and (3) that

$$\begin{aligned} \lambda_{j+1}(\theta) &= \lambda_{j+1,1}(\theta) + \lambda_{j+1,2}(\theta) \\ &= [p_1'(\theta)]^{-1}\lambda_{j,1}'(\theta) + [p_1'(\theta)]^{-1}\lambda_{j,2}'(\theta) \\ &= [p_1'(\theta)]^{-1}\lambda_j'(\theta) . \end{aligned}$$

Conversely, assume $X_1 + X_2$ is an exponential-type r.v.. Then, using (1), (2), and (3), we find that

$$(16) \quad p'(\theta)[\lambda_{j,1}(\theta) + \lambda_{j,2}(\theta)] = p_1'(\theta)\lambda_{j,1}(\theta) + p_2'(\theta)\lambda_{j,2}(\theta) .$$

In particular,

$$(17) \quad p'(\theta)[\lambda_{2,1}(\theta) + \lambda_{2,2}(\theta)] = p'_1(\theta)\lambda_{2,1}(\theta) + p'_2(\theta)\lambda_{2,2}(\theta).$$

Multiplying (16) by $\lambda_{2,1}(\theta)$ and (17) by $\lambda_{j,1}(\theta)$ and then subtracting, we get

$$(18) \quad [p'(\theta) - p'_2(\theta)]h_j(\theta) \equiv 0 \quad \text{for } j \geq 2.$$

Now, if for some $j_0 \geq 2$, $h_{j_0}(\theta) \not\equiv 0$, then there is a subinterval, M , of (a, b) in which $h_{j_0}(\theta) \neq 0$. Then, for $\theta \in M$, $p'_2(\theta) = p'(\theta)$. By analyticity, $p'_2(\theta) = p'(\theta)$ for all $\theta \in (a, b)$. Substitution in (16) yields $p'_1(\theta) = p'(\theta)$ for $\theta \in (a, b)$. If, on the other hand, $h_j(\theta) \equiv 0$, for $j \geq 2$, the result follows from Lemmas 1 and 2 since we assumed that X_1, X_2 are neither both normal nor both Poisson type r.v.'s.

It should be noted that Girshick and Savage [1] proved that if X_1 and X_2 are independent identically distributed r.v.'s such that their sum is of exponential-type, then X_1 and X_2 are also of exponential-type.

The following theorem gives necessary and sufficient conditions for the sum of two Poisson-type r.v.'s to be exponential-type.

THEOREM 2. *If $\log M_i(t; \theta) = C_i t + A_i(\theta)[l^{b_i t} - 1]$, then $X_1 + X_2$ is an exponential-type r.v. if and only if either $b_1 = b_2$ or $p'_1(\theta) = p'_2(\theta)$.*

Proof. If $X_1 + X_2$ is an exponential-type r.v., then, as in the proof of the preceding theorem,

$$[p'(\theta) - p'_2(\theta)]h_j(\theta) \equiv 0 \quad \text{for } j \geq 2.$$

Equivalently,

$$(19) \quad \begin{aligned} & [\lambda_{j,1}(\theta)\lambda_{2,2}(\theta) - \lambda_{j,2}(\theta)\lambda_{2,1}(\theta)] \\ & = p'_2(\theta)[p'(\theta)]^{-1}[\lambda_{j,1}(\theta)\lambda_{2,2}(\theta) - \lambda_{j,2}(\theta)\lambda_{2,1}(\theta)] \end{aligned} \quad \text{for } j \geq 2.$$

Since, for $j \geq 2$, $\lambda_{j,i}(\theta) = b_i^j A_i(\theta)$, (19) becomes

$$[b_1^j b_2^2 - b_2^j b_1^2] A_1(\theta) A_2(\theta) = p'_2(\theta) [p'(\theta)]^{-1} [b_1^j b_2^2 - b_2^j b_1^2] A_1(\theta) A_2(\theta).$$

But $A_1(\theta) A_2(\theta) > 0$, so that

$$[b_1^j b_2^2 - b_2^j b_1^2] = p'_2(\theta) [p'(\theta)]^{-1} [b_1^j b_2^2 - b_2^j b_1^2].$$

Now, if $b_1^j b_2^2 = b_2^j b_1^2$ for all $j \geq 2$, then $b_1^3 b_2^2 = b_2^3 b_1^2$, so that $b_1 = b_2$. On the other hand, if, for some j_0 , $b_1^{j_0} b_2^2 - b_2^{j_0} b_1^2 \neq 0$, then $p'_2(\theta) = p'(\theta)$ and it follows that $p'_1(\theta) = p'_2(\theta)$.

Conversely, if $p'_1(\theta) = p'_2(\theta)$, then $X_1 + X_2$ is an exponential-type r.v. since (1) is satisfied. If $b_1 = b_2$, let

$$p'(\theta) = [A_1'(\theta) + A_2'(\theta)]/b_1[A_1(\theta) + A_2(\theta)] .$$

It is easy to see that (1) is again satisfied.

The author wishes to thank William L. Harkness for his help in the preparation of this paper.

REFERENCES

1. M. Girshick and L. Savage, *Bayes and minimax estimate for quadratic loss functions*, Second Berkeley Symposium on Probability and Statistics, University of California Press, Berkeley, 1951, 67-68.
2. E. L. Lehmann, *Testing Statistical Hypotheses*, John Wiley, New York, 1959.
3. G. P. Patil, *A characterization of the exponential-type distribution*, *Biometrika* **50** (1963), 205-207.

Received August 17, 1964, and in revised form February 26, 1965.

BUCKNELL UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California

R. M. BLUMENTHAL

University of Washington
Seattle, Washington 98105

*J. DUGUNDJI

University of Southern California
Los Angeles, California 90007

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Pacific Journal of Mathematics

Vol. 18, No. 1

March, 1966

Edward Joseph Barbeau, <i>Semi-algebras that are lower semi-lattices</i>	1
Steven Fredrick Bauman, <i>The Klein group as an automorphism group without fixed point</i>	9
Homer Franklin Bechtell, Jr., <i>Frattini subgroups and Φ-central groups</i>	15
Edward Kenneth Blum, <i>A convergent gradient procedure in prehilbert spaces</i>	25
Edward Martin Bolger, <i>The sum of two independent exponential-type random variables</i>	31
David Wilson Bressler and A. P. Morse, <i>Images of measurable sets</i>	37
Dennison Robert Brown and J. G. LaTorre, <i>A characterization of uniquely divisible commutative semigroups</i>	57
Selwyn Ross Caradus, <i>Operators of Riesz type</i>	61
Jeffrey Davis and Isidore Isaac Hirschman, Jr., <i>Toeplitz forms and ultraspherical polynomials</i>	73
Lorraine L. Foster, <i>On the characteristic roots of the product of certain rational integral matrices of order two</i>	97
Alfred Gray and S. M. Shah, <i>Asymptotic values of a holomorphic function with respect to its maximum term</i>	111
Sidney (Denny) L. Gulick, <i>Commutativity and ideals in the biduals of topological algebras</i>	121
G. J. Kurowski, <i>Further results in the theory of monodiffrie functions</i>	139
Lawrence S. Levy, <i>Commutative rings whose homomorphic images are self-injective</i>	149
Calvin T. Long, <i>On real numbers having normality of order k</i>	155
Bertram Mond, <i>An inequality for operators in a Hilbert space</i>	161
John William Neuberger, <i>The lack of self-adjointness in three-point boundary value problems</i>	165
C. A. Persinger, <i>Subsets of n-books in E^3</i>	169
Oscar S. Rothaus and John Griggs Thompson, <i>A combinatorial problem in the symmetric group</i>	175
Rodolfo DeSapio, <i>Unknotting spheres via Smale</i>	179
James E. Shockley, <i>On the functional equation $F(mn)F((m, n)) = F(m)F(n)f((m, n))$</i>	185
Kenneth Edward Whipple, <i>Cauchy sequences in Moore spaces</i>	191