# Pacific Journal of Mathematics

# ON THE SPECTRAL RADIUS OF HERMITIAN ELEMENTS IN GROUP ALGEBRAS

ANDRZEJ HULANICKI

Vol. 18, No. 2

April 1966

### ON THE SPECTRAL RADIUS OF HERMITIAN ELEMENTS IN GROUP ALGEBRAS

#### A. HULANICKI

Let G be a discrete group and  $\mathfrak{A}$  the  $L_1$  algebra over the field of complex numbers of G. The aim of the paper is to consider some combinatorial conditions on the group G which imply symmetry of the algebra  $\mathfrak{A}$ . One result is as follows:

If a group F contains a subgroup G of finite index such that any element of G has finitely many conjugates, then the group algebra  $\mathfrak{A}$  of F is symmetric.

A Banach \*-algebra  $\mathfrak{A}$  is said to be symmetric if for every element  $x \in \mathfrak{A}$  the spectrum of the element  $x^*x$  is real and nonnegative. If the algebra  $\mathfrak{A}$  contains the unit element, this can be equivalently stated as

For every linear functional F on  $\mathfrak{A}$ 

(1)  $F(y + yx^*x) = 0$  for every  $y \in \mathfrak{A}$  implies F = 0.

It is well known and easy to prove that the group algebra of a finite or Abelian locally compact group is symmetric. If a discrete group is neither finite nor Abelian, all that is known about symmetry of its group algebra seems to be due to Bonic [2].

Bonic proved the following facts:

(i) The group algebra of the direct product of two groups one of which is Abelian, the other of which has a symmetric group algebra is symmetric.

(ii) The group algebra of a splitting extension of an Abelian group by the cyclic group of two elements is symmetric.

(iii) The group algebra of a free nonAbelian group is not symmetric.

(i) and (ii) together with the fact that the group algebras of finite and Abelian groups are symmetric establishes the scope of groups about which, using Bonic's results, one may assert symmetry. As a matter of fact, it is not more than the direct products of finitely many finite, Abelian and dihedral-type groups. The method applied by Bonic is based on (1), which in the case of group algebra of a discrete group is equivalent to the implication

(2) If f is bounded and 
$$f(t) + \sum_{s \in G} x^* x(s^{-1}t) f(s) \equiv 0$$
, then  $f \equiv 0$ .

It is not surprising that, if some of the elements of the group appear as complicated expressions of the generators, then to prove or disprove (2) may become difficult.

We propose another approach to this circle of problems.

Our starting point is a theorem of Raikov, cf. [9], [10]. Denote by  $\nu(x)$  the spectral radius of an element x in a Banach \*-algebra  $\mathfrak{A}$ , that is, let

$$u(x) = \lim_{n o \infty} ||x^n||^{1/n}$$
.

For any \*-representation  $x \to T_x$  of  $\mathfrak{A}$  we have

$$u(x^*x) \ge ||T_{x*x}||,$$

cf. e.g. [11]. Raikov's theorem says:

A Banach \*-algebra  $\mathfrak{A}$  is symmetric if, and only if,

$$(3) \qquad \qquad \boldsymbol{\nu}(x^*x) = \sup || \ T_{x^*x} ||$$

where the least upper bound on the right hand side is taken over all \*-representations of  $\mathfrak{A}$ .

Now let  $\mathfrak{A}$  be a group algebra  $\mathfrak{A} = \mathfrak{A}(G)$  of a discrete group G with the usual norm, multiplication, and involution

$$||x||=\sum\limits_{s\in G}|x(s)|$$
 ,  $x*y(s)=\sum\limits_{t\in G}x(t)y(t^{-1}s)$  ,  $x^{\sim}(s)=\overline{x(s^{-1})}$  ,

respectively. Consider the left regular representation T of  $\mathfrak{A}(G) x \to T_x$ , where the Hilbert space of T is  $L_2(G)$  equipped with norm

$$||x||_2 = (\sum\limits_{s \in G} |x(s)|^2)^{1/2} ext{ and } T_x f = x * f, f \in L_2(G)$$
 .

Denote

 $\lambda(x) = || T_x || .$ 

It may happen (cf. [7]) that the left regular representation weakly contains (cf. [3]) all \*-representation of  $\mathfrak{A}(G)$ , which is to say that  $\lambda(x^* * x) \geq ||T'_{x^* * x}||$  for any \*-representation T' of  $\mathfrak{A}(G)$ . Then (3) turns into the equality

(4) 
$$\lambda(x^{\sim} * x) = \nu(x^{\sim} * x) .$$

Our aim is to find sufficient conditions on a group G which imply (4) for any element x of its group algebra  $\mathfrak{A}(G)$ , and consequently the symmetry of  $\mathfrak{A}(G)$ . We formulate the conditions in the forthcoming §1 and we devote §2 to the proof that, in fact, they imply (4). Finally, in §3, we discuss the classes of groups for which our conditions are satisfied.

The author is most indebted to Professor B. H. Neumann for a conversation about these classes of groups as well as to Dr. Neil W. Rickert who suggested several corrections and simplifications.

1. Let G be a discrete group,  $\mathfrak{A} = \mathfrak{A}(G)$  its group algebra. For any function x on G we denote by N(x) the support of the function x, i.e.,

$$N(x) = \{s: x(s) \neq 0\}$$
.

Let  $A_1, \dots, A_n$  be a family of subset of G. By  $A_1 \dots A_n$  we denote the set of all products  $a_1 \dots a_n$ , where  $a_i \in A_i$ ,  $i = 1, \dots, n$ . We shall also use the abbreviated notation  $A^n$  for  $A \dots A$ . For a finite subset A of G we denote by |A| the number of the elements of A. Clearly,

$$(5) \qquad |A_1 \cdots A_n| \leq |A_1| \cdots |A_n|.$$

For any two nonnegative integers  $m \leq n$  we denote by t(m, n) a sequence

$$t(m, n) = \langle t_1, \cdots, t_n \rangle$$

of elements of G such that at most m of the  $t'_i s$ ,  $i = 1, \dots, n$ , are different from the unity of G. We say that a group G satisfies condition (C) if

(C) There exists a constant k such that for any finite set  $A \subset G$  there exists a constant C = C(A, c) such that for any sequence  $\langle t_1, \dots, t_n \rangle = t(m, n)$ 

$$|At_1At_2\cdots At_n| \leq Ck^mc^n$$
 for any  $c>1$  .

For m = 0 (or, which is the same, for bounded m) condition (C) turns into the following one:

(A - S) For any finite subset A of G

$$|A^n| = o(c^n)$$
 for any  $c > 1$  .

This condition has been considered by G. M. Adel'son-Vel'skii and Yu. A. Šreider in [1]. They have proved that (A - S) implies the existence of an invariant Banach mean value on G. Clearly, (A - S) implies the Følner conditions cf. [4].

2. First we formulate two simple lemmas.

LEMMA 1. If G is a (discrete) group, then for any  $x \in \mathfrak{A}(G) \subset L_2(G)$ , we have  $\lambda(x) \geq ||x||_2$ . In fact,

$$\lambda(x) = \sup_{||y||_2=1} ||x * y||_2 \ge ||x * \delta||_2 = ||x||_2$$
,

where  $\delta$ , the unit element of  $\mathfrak{A}(G)$ , is the function which takes value 1 at the unity of G and zero elsewhere.

LEMMA 2. If  $x_1, \dots, x_n \in \mathfrak{A}(G)$ , then (i)  $x_1 * \cdots * x_n(s) = \sum_{t_1,\dots,t_n \in G} x_1(t_1) \cdots x_n(t_n) \delta(t_n^{-1} \cdots t_1^{-1}s)$ (ii)  $N(x_1 * \cdots * x_n) \subseteq N(x_1) \cdots N(x_n)$ (iii)  $N(x^{\sim}) = (N(x))^{-1} = \{t^{-1} : t \in N(x)\}.$ 

The proof of (i) is obtained by an elementary induction. To verify (ii) we simply note that, by (i),  $s \in N(x_1 * \cdots * x_n)$  implies

$$t_{n-1}^{-1} \cdots t_1^{-1} s = t_n \in N(x_n)$$
 and  $t_i \in N(x_i), i = 1, \cdots, n$ .

Therefore  $s = t_1 \cdots t_n \in N(x_1) \cdots N(x_n)$ .

From now on we shall frequently use the abbreviated notation  $x^{**}$  for  $x*\cdots*x$ .

THEOREM 1. If a group G satisfies (A - S), then for every x such that N(x) is finite equality (4) holds.

*Proof.* Suppose N(x) is finite  $\neq \emptyset$ . Let  $A = (N(x))^{-1}N(x)$ . Then A is finite and, by Lemma 2 (ii)-(iii),  $N(x^{\sim} * x) \subset A$ . We have

$$egin{aligned} ||\, (x^{\sim}*x)^{*n}\,||\,&=\sum\limits_{s\,\in\,G}|\, (x^{\sim}*x)^{*n}(s)\,|\ &=\sum\limits_{s\,\in\,A^n}|\, (x^{\sim}*x)^{*n}(s)\,|\,&\leq |\,A^n\,|^{1/2}\,(\sum\limits_{s\,\in\,A^n}|\, (x^{\sim}*x)^{*n}(s)\,|^2)^{1/2}\ &=|\,A^n\,|^{1/2}\,||\, (x^{\sim}*x)^{*n}\,||_2\,\,. \end{aligned}$$

Hence, by Lemma 1, since  $T_{x\sim_{*}x}$  is a hermitian operator,

$$|| (x^{\sim} * x^{*})^{*n} || \leq |A^{n}|^{1/2} \lambda((x^{\sim} * x)^{*n}) = |A^{n}|^{1/2} (\lambda(x^{\sim} * x))^{*n}$$
 .

Consequently,

$$u(x^{\sim}*x) = \lim_{n o \infty} || (x^{\sim}*x)^{*n} ||^{1/n} \leq \lim_{n o \infty} \sup ||A^n|^{1/2n} \lambda(x^{\sim}*x) \; .$$

But, since G satisfies (A-S) and  $A \neq \emptyset$ ,

$$\lim_{n\to\infty}\sup|A^n|^{1/2n}=1,$$

whence  $\nu(x^* * x) \leq \lambda(x^* * x)$ , which completes the proof of Theorem 1.

COROLLARY 1. If a group G satisfies condition (A–S), then for any  $x \in \mathfrak{A}(G)$  such that N(x) is finite  $\operatorname{Sp}_{A(G)}(x^{\sim} * x) \geq 0$ .

*Proof.* By a standard argument (cf. e.g. [9 pp. 300-301]) we reduce our task to proving that for each  $x \in \mathfrak{A}(G)$  with finite support  $(\delta + x^{\sim} * x)^{-1}$  exists. Let  $\mathfrak{A}$  be the (commutative) algebra of hermitian elements generated (algebraically) by  $\delta$  and  $\delta + x^{\sim} * x = y_0$ . Then, by Lemma 2 (ii)-(iii), for any  $y \in A$ , N(y) is finite, whence, since G satisfies (A—S), by Theorem 1,  $\nu(y^{\sim} * y) = \lambda(y^{\sim} * y)$  and, consequently, since  $y^{\sim} = y, \nu(y) = \lambda(y)$ .

Therefore the completion  $\mathfrak{A}^{\nu}$  and  $\mathfrak{A}^{\lambda}$  of  $\mathfrak{A}$  in the norms  $\nu$  and  $\lambda$ respectively, are equal. But  $\mathfrak{A}^{\lambda}$  is isometrically isomorphic to a Banach \*-algebra with the unity of hermitian operators, so  $y_0^{-1}$  exists in  $\mathfrak{A}^{\lambda}$ and, consequently,  $y_0^{-1} \in \mathfrak{A}^{\nu}$ . This show that in the Gelfand representation of  $\mathfrak{A}^{(1)(1)}$ , the completion of  $\mathfrak{A}$  in the norm  $||\cdot||$ , we have  $\hat{y}_0(M) \neq 0$  for any maximal ideal M of  $\mathfrak{A}^{(1)(1)}$ , which, by Gelfand's theorem, proves that  $y_0^{-1} \in \mathfrak{A}^{(1)(1)} \subset \mathfrak{A}(G)$ , as required.

By virtue of Theorem 1, if G satisfies (A—S), then equality (4) holds for a dense (in the norm  $||\cdot||$ ) set of hermitian elements x of  $\mathfrak{A}(G)$ . Clearly the function  $\lambda$  is continuous on  $\mathfrak{A}(G)$ , so if we knew that  $\nu$  were continuous, equality (4) would be an immediate consequence of Theorem 1. However, it is known that, in general, for a noncommutative Banach \*-algebra the function need not be continuous. The example of S. Kakutani, as presented in [11 p. 282], shows that  $\nu$ is not continuous on the algebra of the bounded operators of a Hilbert space. In this case, however,  $\nu$  is continuous on hermitian elements. We do not know whether in the case of a nonsymmetric group algebra the function  $\nu$  is continuous on hermitian elements.

THEOREM 2. If a group G satisfies condition (C), then equality (4) holds for all elements  $x \in \mathfrak{A}(G)$ .

*Proof.* Suppose G satisfies (C) and let  $z \in \mathfrak{A}(G)$ . For any positive  $\varepsilon$  we write

where N(x) = A is finite and  $||y|| < \varepsilon$ . We are going to prove

(7) 
$$\nu(z^* * z) \leq k || y || + \lambda(x) ,$$

where k is the constant whose existence is postulated by condition (C). Since  $\lambda$  is continuous (with respect to the norm  $||\cdot||$ ), (7) implies

A. HULANICKI

$$\mathcal{V}(z^{\sim}*z) \leq \lambda(z^{\sim}*z)$$
,

as desired. To prove (7) we write

$$(z^{\sim}*z)^{*n} = (y + x)^{*n} = \sum_{m=0}^{n} \sum_{\sigma,\tau} y^{*p_1} * x^{*q_1} * \cdots * y^{*p_r} * x^{*q_r}$$
,

where the summation extends over the set C(m, n) of all sequences  $\sigma = \langle p_1, \dots, p_r \rangle$  and  $\tau = \langle q_1, \dots, q_r \rangle$  of nonnegative integers such that  $p_1 + \dots + p_r = m, q_1 + \dots + q_r = m - n$ . Hence

$$(8) \qquad ||(z^{\sim}*z)^{*n}|| \leq \sum_{m=0}^{n} \sum_{\sigma,\tau} ||y^{*p_{1}}*x^{*q_{1}}*\cdots *y^{*p_{r}}*x^{*q_{r}}||.$$

By Lemma 2 (i), we have

$$|| y^{*p_1} * x^{*q_1} * \cdots * y^{*p_r} * x^{*q_r} || = || y^{*p_1} * x^{*q_1} * \cdots * y^{*p_r} * x^{*q_r} * \delta ||$$
  
=  $\sum_{s \in G} \left| \sum_{lpha, eta} y(t_1) \cdots y(t_m) x(s_1) \cdots x(s_{n-m}) \delta(a_r^{-1} b_r^{-1} \cdots a_1^{-1} b_1^{-1} s) \right|,$ 

where the summation is all over the set of sequences

Hence

$$(9) \qquad || y^{*p_{1}} * x^{*q_{1}} * \cdots * y^{*p_{r}} * x^{*q_{r}} || \leq \sum_{s \in G} \sum_{\alpha} |y(t_{1})| \cdots |y(t_{m})|.$$

$$\left| \sum_{\beta} x(b_{1}^{-1}s_{1})x(s_{2}) \cdots x(s_{q_{1}})x(b_{2}^{-1}s_{q_{1}+1})x(s_{q_{1}+2}) \cdots x(s_{q_{1}+q_{2}}) \cdots x(b_{r}^{-1}s_{q_{1}+\dots+q_{r-1}+1})x(s_{q_{1}+\dots+q_{r-1}+2}) \cdots x(s_{n-m})\delta(s_{n-m}^{-1} \cdots s_{1}^{-1}s) \right|$$

$$= \sum_{\alpha} |y(t_{1})| \cdots |y(t_{m})| || x_{b_{1}} * x^{*(q_{1}-1)} * \cdots * x_{b_{r}} * x^{*(q_{r}-1)} ||,$$

where  $x_{b_i}(s) = x(b_i^{-1}s)$ ,  $i = 1, \dots, r$ . But, since N(x) = A,  $N(x_{b_i}) = b_i A$ ,  $i = 1, \dots, r$ . Hence, by Lemma 2(ii),

$$N(x_{b_1} * x^{*(q_1-1)} * \cdots * x_{b_r} * x^{*(q_r-1)}) \subset b_1 A A^{(q_1-1)} \cdots b_r A A^{(q_r-1)} = b_1 A^{q_1} \cdots b_r A^{q_r} = P.$$

Therefore, by Lemma 1 and Schwarz inequality,

(10)  
$$\begin{aligned} || x_{b_1} * x^{*(q_1-1)} * \cdots * x_{b_r} * x_{b_r} * x^{*(q_r-1)} || \\ &\leq |P|^{1/2} || x_{b_1} * x^{*(q_1-1)} * \cdots * x_{b_r} * x^{*(q_r-1)} ||_2 \\ &\leq |P|^{1/2} \lambda(x_{b_1}) (\lambda(x))^{q_1-1} \cdots \lambda(x_{b_r}) (\lambda(x))^{q_r-1} \\ &= |P|^{1/2} (\lambda(x))^{n-m} , \end{aligned}$$

the last equality being true by virtue of the easily verified fact that  $\lambda(x_b) = \lambda(x)$ .

On the other hand,

$$|P| \leq \sup_{b_1, \cdots, b_r \in \mathcal{G}} |b_1 A^{q_1} \cdots b_r A^{q_r}| \leq \sup |t_1 A t_2 A \cdots t_{n-m} A|$$

where the least upper bound on the right hand side is taken over all sequences  $t(m, n - m) = \langle t_1, \dots, t_{n-m} \rangle$  with at most m of the  $t'_i$ s different from the unity of G. Hence, since G satisfies (C),

 $|P|^{1/2} \leq C(A, c)k^mc^n$  for any c > 1 .

Consequently, by (10) and (9),

(11)  
$$\begin{aligned} & || y^{*p_1} * x^{*q_1} * \cdots y^{*p_r} * x^{*q_r} || \\ & \leq \sum_{\alpha} |y(t_1)| \cdots |y(t_m)| k^m c^n C(\lambda(x))^{n-m} \\ & = c^n C k^m || y ||^m (\lambda(x))^{n-m} . \end{aligned}$$

Therefore, by (8) and (11),

$$||_{\mathbf{L}}^{\mathbf{L}}(z^{\sim} * z)^{*n} || \leq f_{\mathcal{A}}(n, n) \sum_{m=0}^{n} {n \choose m} k^{m} || y ||^{m} (\lambda(x))^{n-m} = c^{n} C(k || y || + \lambda(x))^{n}$$
 .

Hence we have

$$egin{aligned} 
u(z^{\sim}*z) &= \lim_{n=\infty} || \, (z^{\sim}*z)^{*n} \, ||^{1/n} \ &\leq \lim_{n=\infty} \sup c C^{1/n}(k \mid \mid y \mid \mid + \lambda(x)) = \, (k \mid \mid y \mid \mid + \lambda(x)) c \; . \end{aligned}$$

This completes the proof of Theorem 2.

3. In this section we present some propositions which exhibit some of the classes of groups for which conditions (A-S) and (C) are satisfied. First we establish some notations:

Let G be a group and let  $a, b \in G$ . We write

$$b^{-1}ab=a^{b},\,a^{-1}b^{-1}ab=[a,\,b]\;,\;[[a_{1},\,\cdots,\,a_{n-1}],\,a_{n}]=[a_{1},\,\cdots,\,a_{n}]$$
 .

If A, B are subsets of G,

$$A^{\scriptscriptstyle B} = \{a^{\scriptscriptstyle b}: a \in A, b \in B\}.$$

REMARK. If A is a subset of G, then for  $a_1, \dots, a_n \in A^G$  and  $b_1, \dots, b_n \in G$  we have

 $a_1b_1\cdots a_nb_n=b_1\cdots b_na_1'\cdots a_n'$ , where  $a_1',\cdots,a_n'\in A^G$ .

(This follows immediately from the fact that  $ab = ba^{b}$  for any  $a, b \in G$ ).

PROPOSITION 1. Let A be a finite subset of a group G such that  $A = A^{a}$ . Then  $|A^{n}| \leq n^{|A|}$ .

*Proof.* If  $a_1, \dots, a_n \in A$ , then

(12) 
$$a_1 \cdots a_n = b_1^{n_1} \cdots b_k^{n_k}, \quad b_1, \cdots, b_k \in A$$

and, moreover,  $n_1 + \cdots + n_k \leq n$  and  $k \leq |A|$ . To prove (12) which is, in fact, a version of Dietzmann's Lemma (cf. [8 Vol. II, p. 154]), we suppose that k is minimal with respect to all representations of  $a_1 \cdots a_n$  of the form (12). Suppose k > |A|. Then there exists at least two elements  $b_i$  and  $b_j$ ,  $1 \leq i < j \leq k$  such that  $b_i = b_j$ . Then

$$b_1^{n_1}\cdots b_k^{n_k}=b_1^{n_1}\cdots b_{i-1}^{n_i-1}b_i^{n_i+n_j}(b_{i+1}^c)^{n_{i+1}}\cdots (b_{j-1}^c)^{n_{j-1}}b_{i+1}^{n_{j+1}}\cdots b_k^{n_k}$$
 ,

where  $c = b_{s}^{n_{j}}$ , which is a contradiction, since  $b_{s}^{c} \in A^{\sigma} = A$ . By (12), we have

$$|A^n| \leq ( ext{the number of the sequences } n_1, \cdots, n_k \ ext{with } \sum_i n_i \leq n ext{ and } k \leq |A|) \leq n^{|A|}$$

DEFINITION. A group G is called a FC-group if for any  $a \in G$  the set  $\{a\}^{a}$  is finite. (cf. e.g. [8 Vol. II, pp. 154 and 269]).

PROPOSITION 2. If G is a FC-group, then it satisfies (C) with constant k = 1.

In fact, let A be a finite subset of G. Since G is FC-group, the set  $B = A^{G}$  is finite. Hence

$$|t_1At_2A\cdots t_nA| \leq |A^{t_2t_3\cdots t_n}A^{t_3\cdots t_n}\cdots A^{t_n}| \leq |B^n| \leq n^{|B|} = o(c^n)$$

for any c > 1.

PROPOSITION 3. If F is a group which contains a subgroup G of finite index in F such that G is a FC-group, then F satisfies (C).

*Proof.* We show first that if M it a finite subset of G, then the set  $M^F$  is finite. Let  $M_0 = M^q$ . Since G is FC-group,  $M_0$  is finite. Denote by  $\phi$  the function which selects one element  $\phi(f)$  out of each of the cosets  $Gf, f \in F$ , and let  $\phi(G) = 1$  — the unit element of F. Let

$$\phi(F)=H=\{h_1,\,\cdots,\,h_k\}$$
 .

We have

$$M^{\scriptscriptstyle F}=M_{\scriptscriptstyle 1}=igcup_{\scriptscriptstyle f\,{f \in}\, F}M_{\scriptscriptstyle 0}^{\phi(f)}$$
 .

In fact, if  $s, f \in F$ , there exists a  $g \in G$  such that

$$\phi(f)s = g\phi(\phi(f)s) = g\phi(fs)$$

Hence

$$M_1^s = \bigcup_{f \in F} M_0^{\phi(f)s} = \bigcup_{f \in F} M_0^{g\phi(fs)} = \bigcup_{f \in F} M_0^{\phi(f)} = M_1$$
.

Thus, since  $\phi(F)$  is finite,  $M^F$  is finite.

Now, let

$$Q = \{h_i h_j (\phi(h_i h_j))^{-1}: i, j = 1, \dots, k\}$$
 .

Clearly, Q is finite and  $Q \subset G$ . By virtue of what we have proved above,  $Q^F = Q_1$  is finite. We now prove that for any sequence  $u_1, \dots, u_n$  with  $u_i \in \{1, \dots, k\}, i = 1, \dots, n$ , we have

$$h_{u_1}\cdots h_{u_n} \in Q^n H \subset Q_1^n H = HQ_1^n$$

Let 
$$f_0 = 1, f_i = h_{u_1} \cdots h_{u_i}, i = 1, \cdots, n$$
, and let  $a_i = \phi(f_i)$ . Then  
 $f_n a_n^{-1} = a_0 h_{u_1} a_1^{-1} a_1 h_{u_2} a_2^{-1} a_2 \cdots a_{n-1}^{-1} a_{n-1} h_{u_n} a_n^{-1}$ .

But for any  $i = 1, \dots, n$ ,

$$a_i = \phi(f_i) = \phi(f_{i-1}h_{u_i}) = \phi(\phi(f_{i-1}(h_{u_i}) = \phi(a_{i-1}h_{u_i}) .$$

Hence

$$f_n a_n^{-1} = a_0 h_{u_1}(\phi(a_0 h_{u_1}))^{-1} \cdots a_{n-1} h_{u_n}(\phi(a_{n-1} h_{u_n}))^{-1} \in Q^n$$

and, consequently,

$$h_{u_1}\cdots h_{u_n}=f_n\in Q^na_n\subset Q^nH$$
 ,

which completes the proof of (13).

Now let A be a finite subset of F. There exists a finite set  $M \subset G$  such that  $A \subset HM$ . Consider the finite set

$$B = HM^{F}$$
.

Clearly,  $A \subset B$ . For every two nonnegative integers  $m \leq n$ , let t(m, n) be as in Section 1. We have

(14) 
$$|t_1B\cdots t_nB| = |(B\cdots B)^{s_1}(B\cdots B)^{s_2}\cdots (B\cdots B)^{s_{m+1}}|,$$

where  $\langle t_1, \dots, t_n \rangle = t(m, n), t_{j_1}, \dots, t_{j_m}$  are the  $t'_{js}$  different from the unity of F, for every  $i = 1, \dots, m s_i = t_{j_i} \cdots t_{j_m}$  and  $s_{m+1} = 1$ .

Each element of the set on the right hand side of (14) is of the form

$$f = (h_1 g_1 \cdots h_{r_1} g_{r_1})^{s_1} \cdots (h_{r_m+1} g_{r_m+1} \cdots h_{r_{m+1}} g_{r_{m+1}})^{s_{m+1}}$$
,

where  $g_i \in M^F$ ,  $i = 1, \dots, n$ . Then

$$f=(h_{{}_{1}}\cdots h_{{}_{r_{1}}})^{s_{1}}\cdots (h_{{}_{r_{m}+1}}\cdots h_{{}_{r_{m}+1}})^{s_{m}+1}g_{{}_{1}}'\cdots g_{{}_{n}}'$$
 ,

where  $g'_i \in M^F$ ,  $i = 1, \dots, n$ . But, by (13), for any  $i = 1, \dots, m+1$ we have

$$h_{r_{i-1}} \cdots h_{r_i} = h'_i q_i$$
, where  $h'_i \in H$  and  $q_i \in Q^{(r_i - r_{i-1})}$ 

Consequently, since  $Q_1^F = Q_1$ ,

$$f = h_1'^{s_1} \cdots h_{m+1}'^{s_{m+1}} q_1' \cdots q_{m+1}' g_1' \cdots g_n'$$

and so, since

.

$$\sum\limits_{i=1}^{m+1} (r_i\!-\!r_{i-1}) = n$$
 , $f\!\in H^{s_1}\cdots H^{s_{m+1}}\!Q_1^nM_1^n$  .

Therefore, since  $|H^s| = |H|$  for any  $s \in F$ , and |H| = k,

$$|t_1B\cdots t_nB| \leq |H^{s_1}|\cdots |H^{s_{m+1}}||Q_1^n||M_1^n| \leq |H|^m |H| n^{(|Q_1|+|M_1|)},$$

whence, since  $|H| n^{(|Q_1|+|M_1|)} = o(c^n)$ , Proposition 3 follows.

COROLLARY 2. If a group G contains a FC-group as a subgroup of finite index, then equality (4) holds for any element x of the group algebra A(G) of G and, consequently, A(G) is symmetric.

PROPOSITION 4. The direct product of finitely many groups which satisfy (C) satisfies (C).

PROPOSITION 5. If G is a locally nilpotent group, then G satisfies (A-S).

This follows from the fact that if  $A = \{a_1, \dots, a_k\}$  is a finite subset of G, H is the subgroup generated by A, which, by assumption, is nilpotent of class, say, s, then any element of  $A^n \subset H$  is of the form

$$\prod_{i=1}^{k} a_{i}^{n(i)} \prod_{i_{1},i_{2}=1}^{k} [a_{i_{1}}, a_{i_{2}}]^{n(i_{1},i_{2})} \cdots \prod_{i_{1},\cdots,i_{s}=1}^{k} [a_{i_{1}}, \cdots, a_{i_{s}}]^{n(i_{1},\cdots,i_{s})}$$

where  $n(i_1, \dots, i_j)$  are nonnegative integers  $\leq f_j(n), j = 1, \dots, s$ , and  $f_j(n)$  is an integral valued polynomial. (The exact from of  $f_j(n)$  can be easily established by the use of the collecting process as described in [6] and [5] but it is irrelevant to us here.) Thus

$$|A^n| \leq (f_1(n))^{k^1} \cdots (f_s(n))^{k^s} = o(c^n)$$
 for any  $c > 1$  .

PROPOSITION 6. Condition (C) is not satisfied by nilpotent groups, in general.

PROPOSITION 7. Condition (A—S) is not satisfied by soluble groups, in general.

#### References

1. G. M. Adel 'son-Vel' skit and Yu. A. Sreider, *The Banach mean on groups*, Uspehi Mat. Nauk (N.S.) **12** (1957), 131-136.

2. Robert A. Bonic, Symmetry in group algebras of discrete groups, Pacific J. Math. 11 (1961), 73-94.

3. J. M. G. Fell, The dual spaces of C\*-algebras, Trans. Amer. Math. Soc. 94 (1960), 365-403.

4. E. Følner, On groups with full Banach mean value, Math. Scand., 3 (1955), 243-254.

5. Marschall Hall, Jr., The theory of groups, The Macmillan Company, 1959.

6. Philip Hall, A contribution to the theory of groups of prime power order, Proc. London Math. Soc. 36 (1933), 29-95.

7. A. Hulanicki, Groups whose regular representation weakly contains all unitary representations, Studia Math. 24 (1964), 37-59.

8. A. G. Kurosh, The theory of group, Chelsea Publishing Company, (1956).

9. M. A. Naimark, Normed rings, P. Noordhoff, Ltd., Groningen, (1960).

10. D. Raikov, On the theory of normed rings with involution, Dokklady Akad. Nauk SSSR 54 (1949), 387-390 (Russian).

11. Charles E. Rickart, General theory of Banach algebras, D. Van Nostrand Company, (1960).

Received January 10, 1965.

INSTITUTE OF MATHEMATICS, POLISH AKADEMY OF SCIENCES and

THE UNIVERSITY OF WASHINGTON

## PACIFIC JOURNAL OF MATHEMATICS

#### EDITORS

H. SAMELSON

Stanford University Stanford, California

#### R. M. BLUMENTHAL

University of Washington Seattle, Washington 98105

#### \*J. DUGUNDJI

University of Southern California Los Angeles, California 90007

#### RICHARD ARENS

University of California Los Angeles, California 90024

#### ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN F. WOLF

K. YOSIDA

#### SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA CALIFORNIA INSTITUTE OF TECHNOLOGY UNIVERSITY OF CALIFORNIA MONTANA STATE UNIVERSITY UNIVERSITY OF NEVADA NEW MEXICO STATE UNIVERSITY OREGON STATE UNIVERSITY UNIVERSITY OF OREGON OSAKA UNIVERSITY UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY UNIVERSITY OF TOKYO UNIVERSITY OF UTAH WASHINGTON STATE UNIVERSITY UNIVERSITY OF WASHINGTON

AMERICAN MATHEMATICAL SOCIETY CHEVRON RESEARCH CORPORATION TRW SYSTEMS NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

\* Paul A. White, Acting Editor until J. Dugundji returns.

# Pacific Journal of Mathematics Vol. 18, No. 2 April, 1966

Alexander V. Arhangelskii, On closed mappings, bicompact spaces, and a	
problem of P. Aleksandrov	201
A. K. Austin, A note on loops	209
Lawrence Peter Belluce and William A. Kirk, Fixed-point theorems for	
families of contraction mappings	213
Luther Elic Claborn, Every abelian group is a class group	219
Luther Elic Claborn, A note on the class group	223
Robert Stephen De Zur, Point-determining homomorphisms on	
multiplicative semi-groups of continuous functions	227
Raymond William Freese, A convexity property	237
Frederick Paul Greenleaf, Characterization of group algebras in terms of	
their translation operators	243
Andrzej Hulanicki, On the spectral radius of hermitian elements in group	077
algebras	211
Michael Bahir Maschler and Bezalel Peleg, A characterization, existence proof and dimension bounds for the kernel of a game	289
Yiannis (John) Nicolas Moschovakis, <i>Many-one degrees of the predicates</i>	
$H_a(x)$	329
G. O. Okikiolu, <i>nth order integral operators associated with Hilbert</i>	
transforms	343
C. E. Rickart, Analytic phenomena in general function algebras	361
K. N. Srivastava, On an entire function of an entire function defined by	
Dirichlet series	379
Paul Elvis Waltman, Oscillation criteria for third order nonlinear	
differential equations	385