

Pacific Journal of Mathematics

**ON DUAL SERIES RELATIONS INVOLVING LAGUERRE
POLYNOMIALS**

K. N. SRIVASTAVA

ON DUAL SERIES RELATIONS INVOLVING LAGUERRE POLYNOMIALS

K. N. SRIVASTAVA

In this paper, we shall consider the problem determining the sequence $\{An\}$, such that

$$\sum_{n=0}^{\infty} \{An/\Gamma(n + \alpha + 1)\} L_n^\alpha(x) = f_1(x), \quad 0 \leq x < y,$$

$$\sum_{n=0}^{\infty} \{An/\Gamma(n + \alpha + 1/2)\} L_n^\alpha(x) = f_2(x), \quad y < x \leq \infty, \alpha > -1/2,$$

where $L_n^\alpha(x)$ is a Laguerre polynomial, the functions $f_1(x)$ and $f_2(x)$ being prescribed. By expressing the sequence $\{An\}$ in terms of a sequence of integrals involving an unknown function $g(u)$ the problem is reduced to that of solving an Abel integral equation for $g(u)$.

In recent years, dual series relations involving Fourier-Bessel, Dini series, trigonometric series and series of Jacobi polynomials have been investigated by various workers [1, 2, 5 to 12]. Here we shall apply the method developed by Sneddon and Srivastav for obtaining a solution of the dual series relations involving Laguerre polynomials.

As pointed out by Sneddon and Srivastav [6], with a view to simplify the calculations, we split the problem posed by the pair of dual equations given above into two parts: Problem (a). Determine the constants $\{An\}$ satisfying the dual series relations

$$(1.1) \quad \sum_{n=0}^{\infty} \{An/\Gamma(n + \alpha + 1)\} L_n^\alpha(x) = f_1(x), \quad 0 \leq x < y,$$

$$(1.2) \quad \sum_{n=0}^{\infty} \{An/\Gamma(n + \alpha + 1/2)\} L_n^\alpha(x) = 0, \quad y < x \leq \infty, \alpha > -1/2.$$

Problem (b). Determine the constants $\{An\}$ satisfying the dual series relations

$$(1.3) \quad \sum_{n=0}^{\infty} \{An/\Gamma(n + \alpha + 1)\} L_n^\alpha(x) = 0, \quad 0 \leq x < y,$$

$$(1.4) \quad \sum_{n=0}^{\infty} \{An/\Gamma(n + \alpha + 1/2)\} L_n^\alpha(x) = f_2(x), \quad y < x \leq \infty, \alpha > -1/2.$$

The solution of the general problem is obviously obtained merely by adding the solutions of problem (a) and (b). We suppose that functions $f_1(x)$ and $f_2(x)$ satisfy the following conditions:

(i) $F_1(x) = x^\alpha f_1(x)$ is finite and continuously differentiable for $0 \leq x < y$,

(ii) $F_2(x) = \int_x^\infty e^{-x} f_2(x) dx$ is finite and continuously differentiable for $y < x \leq \infty$.

As we shall presently see the classes of functions $f_1(x)$ and $f_2(x)$ for which the problem under discussion is solvable, must satisfy the above conditions,

2. In this section we list some results for ready reference. By combining the results [3, p. 292 (2), (3)], we have

$$(2.1) \quad \int_0^\infty x^\alpha e^{-x} L_n^\alpha(x) L_m^\alpha(x) dx = (n+1)_\alpha \cdot \delta_{mn},$$

where δ_{mn} is a Kronecker delta. From [4, p. 193 (27), (28)] we have

$$(2.2) \quad \frac{d}{dn} \{x^\alpha L_n^\alpha(x)\} = (n+\alpha) x^{\alpha-1} L_n^{\alpha-1}(x),$$

$$(2.3) \quad \int_x^\infty e^{-y} L_n^\alpha(y) dy = e^{-x} L_n^{\alpha-1}(x).$$

We shall also require the following results which are easily derived from the more general results given in [3, p. 293 (5), p. 405 (20)]. For $\alpha > -1/2$

$$(2.4) \quad \int_x^\infty (y-x)^{-1/2} e^{-y} L_n^\alpha(y) dy = \Gamma(1/2) e^{-x} L_n^{\alpha-1/2}(x),$$

$$(2.5) \quad \int_0^x (x-y)^{-1/2} y^\alpha L_n^\alpha(y) dy = \frac{\Gamma(n+\alpha+1) \Gamma(1/2)}{\Gamma(n+\alpha+3/2)} x^{\alpha+1/2} L_n^{\alpha+1/2}(x).$$

We also note that if $f(x)$ is continuously differentiable then Abel integral equation

$$(2.6) \quad f(x) = \int_0^x \frac{\phi(y)}{(x-y)^{1/2}} dy$$

has a continuous solution given by the equation

$$(2.7) \quad \phi(y) = \frac{1}{\Pi} \frac{d}{dy} \int_0^y \frac{f(x)}{(y-x)^{1/2}} dx.$$

Furthermore, if $f(x)$ is continuously differentiable then the integral equation

$$(2.8) \quad f(x) = \int_x^\infty \frac{\phi(y)}{(y-x)^{1/2}} dy$$

has a continuous solution

$$(2.9) \quad \phi(y) = -\frac{1}{\Pi} \frac{d}{dy} \int_y^\infty \frac{f(x)}{(x-y)^{1/2}} dx .$$

This can be easily established by simple methods given in [13, p. 229]. The analysis given here is purely formal and no attempt is made to justify the interchange of various limiting processes.

3. Solution of the problem (a). Let us suppose that for $0 \leqq x < y$

$$(3.1) \quad \sum_{n=0}^\infty \{An/\Gamma(n + \alpha + 1/2)\} L_n^\alpha(x) = -e^x \frac{d}{dx} \int_x^y \frac{g_1(u)}{(u-x)^{1/2}} du .$$

Using the orthogonal property (2.1), it can be shown that

$$(3.2) \quad An = -\frac{\Gamma(n + \alpha + 1/2) \Gamma(1/2)}{\Gamma(n + \alpha + 1)} \int_0^y x^\alpha L_n^\alpha(x) \left(\frac{d}{dx} \int_x^y \frac{g_1(u)}{(u-x)^{1/2}} du \right) dx .$$

Since

$$(3.3) \quad -\frac{d}{dx} \int_x^y \frac{g_1(u)}{(u-x)^{1/2}} du = \frac{g_1(y)}{(y-x)^{1/2}} - \int_x^y \frac{d}{du} \left\{ \frac{g_1(u)}{(u-x)^{1/2}} \right\} du$$

we obtain with the help of (2.5), the equation

$$(3.4) \quad An = \Gamma(n + 1) \Gamma(1/2) \int_0^y g_1(u) u^{\alpha-1/2} L_n^{\alpha-1/2}(u) du , \quad n = 0, 1, 2, \dots$$

If in the equation (1.1), we substitute for the coefficients An from (3.4), on interchanging the order of summation and integration, we get

$$(3.5) \quad f_1(x) = \int_0^y g_1(u) u^{\alpha-1/2} K_1(u, x) du , \quad 0 \leqq x < y ,$$

where

$$(3.6) \quad K_1(u, x) = \sum_{n=0}^\infty \frac{\Gamma(n + 1) \Gamma(1/2)}{\Gamma(n + \alpha + 1)} L_n^{\alpha-1/2}(u) L_n^\alpha(x)$$

with the help of equations (2.1) and (2.4) it can be shown that

$$(3.7) \quad K_1(u, x) = e^u x^{-\alpha} (x-u)^{-1/2} H(x-u)$$

where $H(t)$ is Heaviside's unit function. (2.7) is easily proved. Let

$$K_1(u, x) = \sum_{n=0}^\infty a_n L_n^\alpha(x)$$

where the coefficients a_n are given by

$$\begin{aligned} a_n &= \frac{\Gamma(n+1)}{\Gamma(n+\alpha+1)} \int_0^\infty K_1(u, x) x^\alpha e^{-u} L_n^\alpha(x) \\ &= \frac{\Gamma(n+1)}{\Gamma(n+\alpha+1)} e^x \int_u^\infty e^{-x} (x-u)^{-1/2} L_n^\alpha(x) dx \\ &= \frac{\Gamma(n+1) \Gamma(1/2)}{\Gamma(n+\alpha+1)} L_n^{\alpha-1/2}(u). \end{aligned}$$

Thus the equation (3.5) is equivalent to

$$(3.8) \quad F_1(x) = x^\alpha f_1(x) = \int_0^x \frac{g_1(u) u^{\alpha-1/2} e^u}{(x-u)^{1/2}} du, \quad 0 \leq x < y.$$

This is Abel integral equation, since $F_1(x)$ is finite and continuously differentiable, its solution is given by

$$(3.9) \quad u^{\alpha-1/2} e^u g_1(u) = \frac{1}{H} \frac{d}{du} \int_0^u \frac{x^\alpha f_1(x)}{(u-x)^{1/2}} dx.$$

The coefficients An may now be calculated with the help of the relations (3.4) and (3.9).

4. Solution of the problem (b). We start with the assumption that for $y < x \leq \infty$

$$(4.1) \quad \sum_{n=0}^{\infty} \{An/\Gamma(n+\alpha+1)\} L_n^\alpha(x) = x^{-\alpha} \int_y^x \frac{g_2(u)}{(x-u)^{1/2}} du.$$

This is equivalent to assuming that

$$(4.2) \quad An = \Gamma(n+1) \Gamma(1/2) \int_y^\infty g_2(u) e^{-u} L_n^{\alpha-1/2}(u) du, \quad n = 0, 1, 2, \dots$$

If we multiply both sides of the equation (1.4) by $\exp(-x)$ and integrate with respect to x from x to ∞ , $y < x \leq \infty$, we obtain

$$(4.3) \quad F_2(x) = \int_x^\infty e^{-x} f_2(x) dx = \sum_{n=0}^{\infty} \{An/\Gamma(n+\alpha+1)\} e^{-x} L_n^{\alpha-1}(x).$$

Substituting the values of the coefficients from (4.2) in the equation (4.3) we find on interchanging the order of summation and integration that

$$(4.4) \quad e^x F_2(x) = \int_y^\infty g_2(u) e^{-u} K_2(u, x) du, \quad y < x \leq \infty,$$

where

$$(4.5) \quad K_2(u, x) = \sum_{n=0}^{\infty} \frac{\Gamma(n+1) \Gamma(1/2)}{\Gamma(n+\alpha+1/2)} L_n^{\alpha-1/2}(u) L_n^{\alpha-1}(x).$$

From the relations (2.1) and (2.5) it easily follows that

$$(4.6) \quad K_2(u, x) - e^x u^{-\alpha+1/2} (u-x)^{-1/2} H(u-x).$$

Consequently the equation (4.4) reduces to the integral equation

$$(4.7) \quad F_2(x) = \int_y^\infty \frac{g_2(u) u^{1/2-\alpha} e^{-u}}{(u-x)^{1/2}} du, \quad y < x \leq \infty.$$

Since $F_2(x)$ is finite and continuously differentiable, the solution of the above equation is given by

$$(4.8) \quad g_2(u) = -\frac{e^u u^{\alpha-1/2}}{\Pi} \frac{d}{du} \int_0^u \frac{F_2(x)}{(x-u)^{1/2}} dx.$$

The coefficients A_n are given by the relations (4.2) and (4.8).

REFERENCES

1. W. D. Collins, *On some dual series equations and their applications to electrostatic problem of spheroidal caps*, Proc. Camb. Phil. Soc. **57** (1961), 367-384.
2. J. C. Cooke and C. J. Tranter, *Dual Fourier-Bessel series*, Quart. J. Mech. **12** (1959), 379-385.
3. A. Erdelyi (Editor), *Tables of Integral Transforms*, Vol. 2, McGraw-Hill, 1954.
4. ———, *Higher Transcendental Functions*, Vol 2 McGraw-Hill, 1953.
5. B. Noble, *Some dual series equations involving Jacobi polynomials*, Proc. Camb. Phil. Soc. **59** (1963), 363-372.
6. I. N. Sneddon and R. P. Srivastav, *Dual series relations-I. Dual relations involving Fourier-Bessel series*, Proc. Roy. Soc. Edin. A **66** (1964), 150-160.
7. R. P. Srivastav, *Dual series relations-II. Dual relations involving Dini series*, Proc. Roy. Soc. Edin. A **66** (1964), 161-172.
8. ———, *Dual series relations-III. Dual relations involving trigonometric series*, Proc. Roy. Soc. Edin. A **66** (1964) 173-184.
9. ———, *Dual series relations-IV. Dual relations involving series of Jacobi polynomials*, Proc. Roy. Soc. Edin. A **66** (1964), 185-191.
10. C. J. Tranter, *Dual trigonometric series*, Proc. Glasgow Math. Assoc. **4** (1959), 49-57.
11. ———, *A further note on dual trigonometric series*, Proc. Glasgow Math. Assoc. **4** (1960), 198-200.
12. ———, *An improved method for dual trigonometrical series*, Proc. Glasgow Math. Assoc. **6** (1964), 136-140.
13. E. T. Whittaker and G. N. Watson, *A Course of Modern Analysis*, Cambridge Univ. Press, 1920.

Received June 29, 1965.

M. A. COLLEGE OF TECHNOLOGY, BHOPAL
INDIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California

J. P. JANS
University of Washington
Seattle, Washington 98105

J. DUGUNDJI
University of Southern California
Los Angeles, California 90007

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 19, No. 3

July, 1966

S. J. Bernau, <i>The spectral theorem for unbounded normal operators</i>	391
Lu-san Chen, <i>Asymptotic behavior of solutions of parabolic equations of higher order</i>	407
Lawrence William Conlon, <i>An application of the Bott suspension map to the topology of EIV</i>	411
Neal Eugene Foland and John M. Marr, <i>Sets with zero-dimensional kernels</i>	429
Stanley Phillip Franklin and R. H. Sorgenfrey, <i>Closed and image-closed relations</i>	433
William Jesse Gray, <i>A note on topological transformation groups with a fixed end point</i>	441
Myron Goldstein, <i>K- and L-kernels on an arbitrary Riemann surface</i>	449
George Joseph Kertz and Francis Regan, <i>The exponential analogue of a generalized Weierstrass series</i>	461
Walter Leighton, <i>On Liapunov functions with a single critical point</i>	467
Bernard Werner Levinger and Richard Steven Varga, <i>On a problem of O. Taussky</i>	473
Lowell Duane Loveland, <i>Tame subsets of spheres in E^3</i>	489
Erik Andrew Schreiner, <i>Modular pairs in orthomodular lattices</i>	519
K. N. Srivastava, <i>On dual series relations involving Laguerre polynomials</i>	529
Arthur Steger, <i>Diagonability of idempotent matrices</i>	535
Walter Strauss, <i>On continuity of functions with values in various Banach spaces</i>	543
Robert Vermes, <i>On the zeros of a linear combination of polynomials</i>	553
Elliot Carl Weinberg, <i>On the scarcity of lattice-ordered matrix rings</i>	561
Harold Widom, <i>Toeplitz operators on H_p</i>	573
Neal Zierler, <i>On the lattice of closed subspaces of Hilbert space</i>	583
Irving Leonard Glicksberg, <i>Correction to: "Maximal algebras and a theorem of Radó"</i>	587
John Spurgeon Bradley, <i>Correction to: "Adjoint quasi-differential operators of Euler type"</i>	587
William Branham Jones, <i>Erratum: "Duality and types of completeness in locally convex spaces"</i>	588
Stanley P. Gudder, <i>Erratum: "Uniqueness and existence properties of bounded observables"</i>	588