

Pacific Journal of Mathematics

**A REPRESENTATION THEOREM FOR ABELIAN GROUPS
WITH NO ELEMENTS OF INFINITE p -HEIGHT**

DELMAR L. BOYER AND ADOLF G. MADER

A REPRESENTATION THEOREM FOR ABELIAN GROUPS WITH NO ELEMENTS OF INFINITE P-HEIGHT

D. L. BOYER AND A. MADER

The purpose of this note is to give a generalization of the representation Theorems 33.1 and 33.2 of [2]. Let G be an arbitrary abelian group and $B = [\bigoplus_{\lambda \in \mathcal{A}} \langle x_\lambda \rangle] \oplus [\bigoplus_{i \geq 1} B_i]$ be a p -basic subgroup of G , cf. [3], where $\bigoplus_{\lambda \in \mathcal{A}} \langle x_\lambda \rangle$ is the torsion-free part. For all $\lambda \in \mathcal{A}$ let $(F_p^*)_\lambda$ be a copy of the group of p -adic integers, and let $(F_p)_\lambda$ denote the infinite cyclic group of finite p -adic integers in $(F_p^*)_\lambda$. Then G can be mapped homomorphically into the complete direct sum $[\bigoplus_{\lambda \in \mathcal{A}} (F_p^*)_\lambda] \oplus [\bigoplus_{i \geq 1} B_i]$ with kernel $p^\omega G$. Furthermore, the image of G is a p -pure subgroup which contains $[\bigoplus_{\lambda \in \mathcal{A}} (F_p)_\lambda] \oplus [\bigoplus_{i \geq 1} B_i]$ as a p -basic subgroup and is in turn contained in the p -adic completion of this subgroup (See Section 1 for definitions). This representation is completely analogous to the representation theorem for p -groups which is contained as a special case, and hopefully it is of similar use.

Definitions and facts concerning p -adic and n -adic topologies.
 In this article we list the definitions and facts concerning p -adic and n -adic topologies that are needed in this paper. For references see [2], [3], and [5].

DEFINITION 1.1. The p -adic topology for an abelian group is the topology with the subgroups $p^n G$, $n = 1, 2, \dots$ as a basis for the neighborhoods of 0.

DEFINITION 1.2. The n -adic topology for an abelian group G is the topology with the subgroups $n!G$, $n = 1, 2, \dots$ as a basis for the neighborhoods of 0.

DEFINITION 1.3. The completion of an abelian group in the p -adic (resp. n -adic) topology is its metric space completion with respect to the metric $d(x, y) = 10^{-m}$, where m is the largest integer such that $x - y \in p^m G$ (resp. $m!G$).

PROPOSITION 1.4. If H is a p -pure (resp. pure) subgroup of the abelian group G , then the p -adic (resp. n -adic) topology of the subgroup is the same as the induced p -adic (resp. n -adic) topology.

THEOREM 1.5. *If an abelian group is complete in the n -adic topology, then it is a direct summand of every abelian group that*

contains it as a pure subgroup.

PROPOSITION 1.6. A subgroup H of an abelian group G is dense in the p -adic (resp. n -adic) topology if and only if the quotient group G/H is p -divisible (respectively divisible).

2. The representation theorems. Let G be an abelian group, let B be a p -basic subgroup, cf. Fuchs [3], of G , and we write $B = \bigoplus_{n \geq 0} B_n$ and $B_0 = \bigoplus_{\lambda \in A} \langle x_\lambda \rangle$. As in [1 p. 325], for each $g \in G$ and each natural n , we can write

2.1. $g = b_0^{(n)} + b_1 + \cdots + b_n + b_n^* + p^n g_n$ where $b_0^{(n)} \in B_0$, $b_i \in B_i$ for $1 \leq i \leq n$, $b_n^* \in \bigoplus_{i > n} B_i$, and $g_n \in G$. It is proved in [1] p. 326 that the b_i , $i \geq 1$, are unique in any such representation, and that, given two such representations, one for n and one for m , we have

$$2.2 \quad b_0^{(n)} - b_0^{(m)} \in p^{\min(m,n)} G.$$

For each λ , let $(F_p^*)_\lambda$ be the group of p -adic integers, and let $(F_p)_\lambda$ be the infinite cyclic subgroup of finite p -adic integers. We introduce the notation $P_1 = \bigoplus_{\lambda \in A}^* (F_p^*)_\lambda$, and $P_2 = \bigoplus_{i \geq 1}^* B_i$. P_1 and P_2 are complete groups in the n -adic topology, and the n -adic topology coincides with the p -adic topology. $\bigoplus_{\lambda \in A} (F_p^*)_\lambda$ and $\bigoplus_{i \geq 1} B_i$ are pure subgroups of P_1 and P_2 , hence they possess completions in P_1 and P_2 for the coinciding n -adic and p -adic topologies. Let $C_1 = [\bigoplus_{\lambda \in A} (F_p^*)_\lambda]^*$ and $C_2 = [\bigoplus_{i \geq 1} B_i]^*$, where the $*$ indicates the completion. Notice that C_i is a direct summand of P_i , $i = 1, 2$.

We define a map $\sigma: G \rightarrow P_1 \oplus P_2$ as follows. Let g have the representation 2.1 for each n . Write $b_0^{(n)} = \sum_{\lambda \in A} m_\lambda^{(n)} x_\lambda$, and write $m_\lambda^{(n)}$ in its p -adic expansion

$$2.3 \quad m_\lambda^{(n)} = \sum_{k \geq 0} a_{\lambda,k}^{(n)} p^k, \quad 0 \leq a_{\lambda,k}^{(n)} \leq p-1.$$

It follows from 2.2 that $a_{\lambda,k}^{(n)}$ is independent of n for $k < n$. Now define

$$2.4 \quad g\sigma = (\cdots, \sum_{k \geq 0} a_{\lambda,k}^{(k+1)} p^k, \cdots; b_1, b_2, \cdots).$$

THEOREM 2.5. *The map σ is a homomorphism, and $\ker \sigma = p^\circ G$, the subgroup of elements of infinite p -height. The p -basic subgroup B of G is mapped onto the group $[\bigoplus_{\lambda \in A} (F_p)_\lambda] \oplus [\bigoplus_{i \geq 1} B_i]$ which is a p -basic subgroup of $C_1 \oplus C_2$.*

Proof. It is easy to see that σ is a homomorphism. Let $g \in p^\circ G$, and write g as in 2.1. Then by the p -purity of B , each of $b_0^{(n)}$, b_1, \dots, b_n, b_n^* is divisible by p^n in the summand of B to which it belongs. Hence $b_1 = \cdots = b_n = 0$. Since $b_0^{(n)}$ is divisible by p^n in

in B_0 , it follows that in $m_\lambda^{(n)} = \sum_{k \geq 0} a_{\lambda,k}^{(n)} p^k$ the coefficient $a_{\lambda,k}^{(n)} = 0$ for $k \leq n - 1$. Thus $g\sigma = 0$. Conversely, assume $g\sigma = 0$. Then in the representation 2.1, $b_1 = b_2 = \dots = b_n = 0$, and in the equation $m_\lambda^{(n)} = \sum_{k \geq 0} a_{\lambda,k}^{(n)} p^k$, $0 \leq a_{\lambda,k}^{(n)} \leq p - 1$, we have $a_{\lambda,k}^{(k+1)} = 0$ for each k . The uniqueness of the $a_{\lambda,k}^{(n)}$ for $k < n$ implies $a_{\lambda,k}^{(n)} = 0$ for $0 \leq k < n$, i.e. $m_\lambda^{(n)}$ is divisible by p^n . Thus $b_0^{(n)}$ is divisible in B_0 by p^n . The remainder of this part of the proof is exactly as in the proof of Theorem 3 in [1] pp. 326-7. It is obvious from 2.1 that B is mapped onto

$$[\bigoplus_{\lambda \in I} (F_p)_\lambda] \oplus [\bigoplus_{i \geq 1} B_i],$$

and it is easy to check that this is a p -basic subgroup of $C_1 \oplus C_2$

THEOREM 2.6. *$G\sigma$ is p -pure in $P_1 \oplus P_2$, and $(G\sigma)^* = C_1 \oplus C_2$, where $*$ indicates the completion in the p -adic topology.*

Proof. By 2.5 $B\sigma$ is a p -pure subgroup of $P_1 \oplus P_2$. Since $G\sigma/B\sigma$ is a p -divisible (hence p -pure) subgroup of $(P_1 \oplus P_2)/B\sigma$, it follows that $G\sigma$ is a p -pure subgroup of $P_1 \oplus P_2$. Since $G\sigma$ is p -pure in $P_1 \oplus P_2$ it possesses a p -adic completion in $P_1 \oplus P_2$. $B\sigma \leq G\sigma$ implies $C_1 \oplus C_2 = (B\sigma)^* \leq (G\sigma)^*$, and since $B\sigma$ is dense in $G\sigma$ in the p -adic topology, $G\sigma \leq (B\sigma)^* = C_1 \oplus C_2$, thus $(G\sigma)^* \leq C_1 \oplus C_2$.

COROLLARY 2.7. *Every abelian group G with no elements of infinite p -height may be considered to be a p -pure subgroup of some group $[\bigoplus_{\lambda \in I}^* (F_p)_\lambda] \oplus [\bigoplus_{i \geq 1}^* B_i]$ and containing $[\bigoplus_{\lambda \in I} (F_p)_\lambda] \oplus [\bigoplus_{i \geq 1} B_i]$ as a p -basic subgroup.*

If G is a p -group, then $P_1 = 0$, and $G\sigma \leq (C_2)_t$, the torsion subgroup of C_2 . Thus in this case our theorems are exactly the important and useful Theorems 33.1 and 33.2 of [2].

REFERENCES

1. D. Boyer, *On the theory of p -basic subgroups of abelian groups*, Topics in abelian groups, Scott, Foresman and Company, 1963.
2. L. Fuchs, *Abelian Groups*, Pergamon Press, 1960.
3. ——— *Notes on abelian groups II*, Acta Math. Acad. Sci. Hung. XI (1960), 117-125.
4. D. Harrison, *Infinite abelian groups and homological methods*, Ann. of Math. (2) **69** (1959), 366-391.
5. I. Kaplansky, *Infinite abelian groups*, University of Michigan Press, 1954.

Received May 14, 1965.

UNIVERSITY OF IDAHO

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California

J. P. JANS

University of Washington
Seattle, Washington 98105

J. DUGUNDJI

University of Southern California
Los Angeles, California 90007

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Pacific Journal of Mathematics

Vol. 20, No. 1

September, 1967

Leonard Daniel Baumert, <i>Extreme copositive quadratic forms. II</i>	1
Edward Lee Bethel, <i>A note on continuous collections of disjoint continua</i>	21
Delmar L. Boyer and Adolf G. Mader, <i>A representation theorem for abelian groups with no elements of infinite p-height</i>	31
Jean-Claude B. Derderian, <i>Residuated mappings</i>	35
Burton I. Fein, <i>Representations of direct products of finite groups</i>	45
John Brady Garnett, <i>A topological characterization of Gleason parts</i>	59
Herbert Meyer Kamowitz, <i>On operators whose spectrum lies on a circle or a line</i>	65
Ignacy I. Kotlarski, <i>On characterizing the gamma and the normal distribution</i>	69
Yu-Lee Lee, <i>Topologies with the same class of homeomorphisms</i>	77
Moshe Mangad, <i>Asymptotic expansions of Fourier transforms and discrete polyharmonic Green's functions</i>	85
Jürg Thomas Marti, <i>On integro-differential equations in Banach spaces</i>	99
Walter Philipp, <i>Some metrical theorems in number theory</i>	109
Maxwell Alexander Rosenlicht, <i>Another proof of a theorem on rational cross sections</i>	129
Kenneth Allen Ross and Karl Robert Stromberg, <i>Jessen's theorem on Riemann sums for locally compact groups</i>	135
Stephen Simons, <i>A theorem on lattice ordered groups, results of Ptak, Namioka and Banach, and a front-ended proof of Lebesgue's theorem</i>	149
Morton Lincoln Slater, <i>On the equation $\varphi(x) = \int_x^{x+1} K(\xi) f[\varphi(\xi)] d\xi$</i>	155
Arthur William John Stoddart, <i>Existence of optimal controls</i>	167
Burnett Roland Toskey, <i>A system of canonical forms for rings on a direct sum of two infinite cyclic groups</i>	179
Jerry Eugene Vaughan, <i>A modification of Morita's characterization of dimension</i>	189