

Pacific Journal of Mathematics

A TOPOLOGICAL CHARACTERIZATION OF GLEASON PARTS

JOHN BRADY GARNETT

A TOPOLOGICAL CHARACTERIZATION OF GLEASON PARTS

JOHN GARNETT

Let A be a function algebra on its maximal ideal space $M(A)$, and let P be a Gleason part of $M(A)$. It is easily seen that P is then a σ -compact completely regular space. We prove the converse: if K is completely regular and σ -compact, then there exists a function algebra whose maximal ideal space contains a part homeomorphic to K . Every bounded continuous function on that part is the restriction of a function in the given algebra. Consequently no subset of the part can have an analytic structure.

Suppose X is a compact Hausdorff space and A is a subalgebra of $C(X)$, the algebra of continuous complex valued functions on X . Assume A separates the points of X , contains the constant functions, and is uniformly closed. A is then called a function algebra on X . With the weak star topology, the maximal ideal space $M(A)$ of A is a compact Hausdorff space. We consider X as embedded in $M(A)$ and A as a function algebra on $M(A)$.

In [4] Gleason noted that an equivalence relation could be defined on $M(A)$ by setting $x \sim y$ when the functional norm $\|x - y\|_{A^*} < 2$. The equivalence classes for this relation are called the "parts" of $M(A)$. In certain cases parts have been used to impose an analytic structure on $M(A)$ (see for example [7]).

Let P be a part of some $M(A)$. Then clearly P is a completely regular space and fixing $p \in P$ we have

$$P = \bigcup_{n=1}^{\infty} \{q \in M(A) : \|p - q\| \leq 2 - 1/n\},$$

where each term in the union is weak star closed, and hence compact, so that P is σ -compact.

Some results in this paper have been announced in [3].

2. We begin with a theorem which will be our basic tool in constructing parts.

THEOREM 1. *Let A be a function algebra, S a hull-kernel closed subset of $M(A)$ and P a part of $M(A)$. Then there is a function algebra B such that $M(B)$ contains a part Q homeomorphic to $P \cap S$. Moreover, $B|_Q$ is isometrically isomorphic to $A|_{P \cap S}$.*

Let α be a positive irrational number, and denote by A_α the function algebra on the torus T^2 generated by the functions $z \rightarrow z_1^m z_2^n$ where $m + n\alpha \geq 0$. Let m^0 be the point in $M(A_\alpha)$ represented by Haar measure on T^2 (which is multiplicative on A_α). Then $m^0 \notin T^2$ and $\{m^0\}$ is a part of $M(A_\alpha)$ ([5] p. 316). If J is a proper closed subset of T^2 , then $A_\alpha \setminus J$ is dense in $C(J)$ ([8] pp. 69–70), so that when $x \in M(A_\alpha) \setminus J$ there is a function $f \in A_\alpha$ such that $|f(x)| > \max_{z \in J} |f(z)|$, as otherwise evaluation at x would induce a complex homomorphism of $C(J)$.

Proof of Theorem 1. Let $A_\alpha \otimes A$ be the function algebra on $M(A_\alpha) \times M(A)$ generated by the functions of the form $(x, y) \rightarrow f(x)g(y)$ where $f \in A_\alpha$ and $g \in A$. $M(A_\alpha \otimes A)$ is homeomorphic to $M(A_\alpha) \times M(A)$ in a natural fashion.

Set $J = \{z \in T^2 : \text{Real } z_1 \leq 0\}$ and

$$X = (J \times M(A)) \cup (M(A_\alpha) \times S).$$

X is a compact subset of $M(A_\alpha) \times M(A)$. Our algebra B is the uniform closure on X of $\{h \mid X : h \in A_\alpha \otimes A\}$. $M(B)$ is then the $A_\alpha \otimes A$ -hull of X ,

$$\{g \in M(A_\alpha \otimes A) : |g(q)| \leq \max_{p \in X} |g(p)| \text{ for all } g \in A_\alpha \otimes A\}.$$

If $(x^0, y^0) \in M(A_\alpha \otimes A) \setminus X$, then $x^0 \notin J$ and $y^0 \notin S$. As S is hull-kernel closed, there is a function g in A with $g(y^0) = 1$ and $g(S) = 0$. As $x^0 \notin J$, there is a function f in A_α with $f(x^0) = 1$ and $\max_{z \in J} |f(z)| < 1$. Replace f by a suitable power f^n so that $\max_{z \in J} |f^n(z)| < (1/||g||)$. Then $h(x, y) = f^n(x)g(y)$ is in $A_\alpha \otimes A$ and $h(x^0, y^0) = 1$ while $\max_{p \in X} |h(p)| < 1$. Hence $M(B) = X$.

Take $Q = \{m^0\} \times (P \cap S)$. Then Q is subset of X . For $s \in P \cap S$, let $p_s = (m^0, s) \in Q$. Let $(x^0, y^0) \in X \setminus Q$. If $x^0 \neq m^0$, then using functions of the variable $x \in M(A_\alpha)$ alone we see that $(x^0, y^0) \not\sim p_s$ for any $s \in P \cap S$. Similarly if $y^0 \notin P$, then $(x^0, y^0) \not\sim p_s$ for all such s . Finally if $y^0 \in S$, then by the choice of X , $x^0 \neq m^0$. Hence Q is a union of parts.

If $s \in P \cap S$ and $g \in B$, then

$$g(p_s) = \int_{T^2} g(x, s) d\lambda$$

where λ is normalized Haar measure on T^2 , because λ represents m^0 for A_α . Take s and t in $P \cap S$, $g \in B$ with $||g|| \leq 1$. Then

$$\begin{aligned} |g(p_s) - g(p_t)| &\leq \int_{T^2} |g(x, s) - g(x, t)| d\lambda \\ &\leq \int_{T^2 \setminus J} 2d\lambda + \int_J |g(x, s) - g(x, t)| d\lambda. \end{aligned}$$

Now if $x \in J$, then $\{x\} \times M(A) \subset X$ so that $y \rightarrow g(x, y)$ is in A with norm ≤ 1 . Therefore there is a constant $c < 2$ such that for each $x \in J$ $|g(x, s) - g(x, t)| < c$, because $s \sim t$. Hence

$$|g(p_s) - g(p_t)| \leq \int_{x^2, J} 2d\lambda + \int_J cd\lambda < 2$$

and $p_s \sim p_t$. Thus Q is a part.

It is obvious that Q is homeomorphic to $P \cap S$ and that $B|Q = A|P \cap S$, because the coordinate x is constant on Q .

As a corollary to Theorem 1 we now prove a special case of our main result because in this case the proof is much simpler.

D denotes the closed unit disc in the complex plane and D^0 its interior. A_0 is the algebra of all functions continuous on D and analytic on D^0 . If K is a locally compact Hausdorff space, then $K^* = K \cup \{\infty\}$ is its one point compactification.

COROLLARY. *Let K be a locally compact σ -compact Hausdorff space. Then there exists a function algebra B such that $M(B)$ contains a part Q homeomorphic to K . Moreover $B|Q$ is isometrically isomorphic to $C(K^*)|K$.*

Proof. Let $A = \{f \in C(K^* \times D) : f| \{x\} \times D \in A_0 \text{ for each } x \in K^* \text{ and } f|K^* \times \{0\} \text{ is constant}\}$. Then $M(A) = K^* \times D / \approx$ where \approx identifies $K^* \times \{0\}$ to a point, and $P = \{(x, z) \in M(A) : |z| < 1\}$ is a part in $M(A)$, as P is a union of discs with the centers identified.

Since K is σ -compact, $\{\infty\}$ is a G_δ set in K^* . Hence there is a continuous function $h: K^* \rightarrow [1/2, 1]$ such that $h^{-1}(1) = \{\infty\}$. Let $S \subset M(A)$ be the graph of h , $S = \{(x, h(x)) : x \in K^*\}$. Then the function $g(x, z) = (h(x) - z/3h(x) - z)$ is in A and vanishes exactly on S , so that S is hull-kernal closed. And clearly $S \cap P$ is homeomorphic to K . Finally if $f \in C(K^*)$, then $f'(x, z) = zf(x)/h(x)$ is in A and $f'(x, h(x)) = f(x)$ when $x \in K$. Thus $A|S \cap P \cong C(K^*)|K$. The conclusion of the corollary now follows directly from Theorem 1.

3. Before proving our main theorem we construct the algebra to be used in place of the disc algebra A_0 . Let I be an index set, and let Y_I be the product of discs, $Y_I = \prod(D : i \in I)$. Denote by A_I the subalgebra of $C(Y_I)$ generated by the coordinate functions $z_i, i \in I$ where $z_i(p) = p_i$. Then $M(A_I) = Y_I$, for if $\varphi \in M(A_I)$, then $|\varphi(z_i)| \leq i$ so that φ is evaluation at $\lambda \in Y_I$ where $\lambda_i = \varphi(z_i)$. Let θ be the "origin" in $Y_I, z_i(0) = 0$ for all $i \in I$, and let P_0 be the part of $M(A_I)$ containing θ . We now need a well known fact which is proved using elementary conformal mappings of the disc D . If $(g_n)_{n=1}^\infty$ is a sequence in A with $\|g_n\| \leq 1$ and $g_n(x) \rightarrow 1$, then $x \sim y$ implies $g_n(y) \rightarrow 1$.

LEMMA. Let $p \in M(A_I)$. Then $p \in P_0$ if and only if there exists $a < 1$ such that $|z_i(p)| \leq a$ for all $i \in I$.

Proof. If no such number exists, then there is a sequence $(g_n)_{n=1}^\infty$ of coordinate functions z_i such that $g_n(p) \rightarrow 1$ while $g_n(\theta) = 0$. Hence $p \not\sim \theta$, by the above remark. If such an a exists, then let $\rho: D \rightarrow Y_I$ by $(\rho(t))_i = t/a \cdot p_i$. Then $\rho(0) = \theta$, $\rho(a) = p$ and for $f \in A_I$, $f \circ \rho \in A_0$ with $\|f \circ \rho\| \leq \|f\|$. Then as $\theta \sim a$ for A_0 we have $\theta \sim p$.

THEOREM 2. Let K be a σ -compact completely regular space. Then there is an algebra B and a part $Q \subset M(B)$ such that Q is homeomorphic to K and $B|Q \cong C^b(K)$, the algebra of bounded continuous functions on K .

Proof. Let βK be the Stone-Ćech compactification of K . Take⁽¹⁾ $I = \beta K \setminus K$ and set $A = \{f \in C(\beta K \times Y_I) : f|_{\{x\} \times Y_I} \in A_I \text{ for all } x \in \beta K \text{ and } f|_{\beta K \times \{\theta\}} \text{ is constant}\}$. Then $M(A) = \beta K \times Y_I / \approx$ where \approx identifies $\beta K \times \{\theta\}$ to a point, and $P = \{(x, z) \in M(A) : z \in P_0\}$ is a part of $M(A)$.

Write $K = \bigcup_{n=1}^\infty K_n$, where $K_n \subset K_{n+1}$ and each K_n is compact. Then for each $t \in \beta K \setminus K$ there exists a continuous function $h_t: \beta K \rightarrow [1/2, 1]$ with $h_t(t) = 1$ and $h_t(x) \leq 1 - 2^{-n}$ when $x \in K_n$. Let $\rho: \beta K \rightarrow M(A)$ be defined by $\rho(x) = (x, H(x))$ where $(H(x))_t = h_t(x)$ for each $t \in \beta K \setminus K$. Then ρ is a homeomorphism of βK onto $S = \rho(\beta K)$ and $\rho(K) = S \cap P$ by the above lemma. S is hull kernel closed in $M(A)$ because $S = \bigcap \{g_t^{-1}(0) : t \in \beta K \setminus K\}$ where $g_t(x, z) = (h_t(x) - z_t)/3h_t(x) - z_t$. And $A|S \cap P \cong C^b(K)$, because if $f \in C^b(K)$ and \tilde{f} is its unique extension to βK , then for any $t \in \beta K \setminus K$, $f'(x, z) = z_t \tilde{f}(x)/h_t(x)$ is in A and $f' = \tilde{f} \circ \rho^{-1}$ on S . The conclusion now follows from Theorem 1.

We remark that with these arguments one can get some restriction algebras $B|Q$ different from $C^b(K)$. For example, if K is compact and A_I is an algebra with $M(A_I) = K$, then there is an algebra B with part Q homeomorphic to K and $B|Q = A_I$.

4. Acknowledgments. Professor Eva Kallin Pohlmann³ simplified the example given in [3], enabling us to improve our arguments significantly. We thank Professor Lewis Robertson for a helpful conversation. Finally, we are deeply indebted to Professor Irving Glicksberg for introducing us to function algebras and for guiding our research throughout.

¹ If K is compact, let I be a singleton, and proceed as in the corollary.

² She observed that $X = \{(z, w) : |z| \leq 1, w = \pm 1/2\} \cup \{(z, w) : |z| = 1, \text{Im } z \geq 0, |w| \leq 1\}$ is a polynomially convex subset of \mathbb{C}^2 containing a part consisting of two discs.

REFERENCES

1. R. Arens, and I. M. Singer, *Generalized analytic functions*, Trans. Amer. Math. Soc. **81** (1956), 379-395.
2. E. Bishop, *Representing measures for points in a uniform algebra*, Bull. Amer. Math. Soc. **70** (1964), 121-122.
3. J. Garnett, *Disconnected Gleason parts*, Bull. Amer. Math. Soc. **72** (1966).
4. A. Gleason, *Function algebras*, Seminars on analytic functions, Vol. 2, Inst. for Advanced Study, Princeton, N. J., 1957.
5. K. Hoffman, *Analytic functions and logmodular Banach algebras*, Acta Math. **108** (1962), 271-317.
6. M. A. Naimark, *Normed rings*, P. Noordhoff, Groningen, 1959.
7. J. Wermer, *Dirichlet algebras*, Duke Math. J. **27** (1960), 373-382.
8. ———, *Banach algebras and analytic functions*, Advances in Math., Academic Press, New York, 1961.

Received January 12, 1966. The author holds a National Science Foundation Graduate Fellowship. This research is a part of his doctoral dissertation to be submitted to the University of Washington.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California

J. P. JANS

University of Washington
Seattle, Washington 98105

J. DUGUNDJI

University of Southern California
Los Angeles, California 90007

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Leonard Daniel Baumert, <i>Extreme copositive quadratic forms. II</i>	1
Edward Lee Bethel, <i>A note on continuous collections of disjoint continua</i>	21
Delmar L. Boyer and Adolf G. Mader, <i>A representation theorem for abelian groups with no elements of infinite p-height</i>	31
Jean-Claude B. Derderian, <i>Residuated mappings</i>	35
Burton I. Fein, <i>Representations of direct products of finite groups</i>	45
John Brady Garnett, <i>A topological characterization of Gleason parts</i>	59
Herbert Meyer Kamowitz, <i>On operators whose spectrum lies on a circle or a line</i>	65
Ignacy I. Kotlarski, <i>On characterizing the gamma and the normal distribution</i>	69
Yu-Lee Lee, <i>Topologies with the same class of homeomorphisms</i>	77
Moshe Mangad, <i>Asymptotic expansions of Fourier transforms and discrete polyharmonic Green's functions</i>	85
Jürg Thomas Marti, <i>On integro-differential equations in Banach spaces</i>	99
Walter Philipp, <i>Some metrical theorems in number theory</i>	109
Maxwell Alexander Rosenlicht, <i>Another proof of a theorem on rational cross sections</i>	129
Kenneth Allen Ross and Karl Robert Stromberg, <i>Jessen's theorem on Riemann sums for locally compact groups</i>	135
Stephen Simons, <i>A theorem on lattice ordered groups, results of Ptak, Namioka and Banach, and a front-ended proof of Lebesgue's theorem</i>	149
Morton Lincoln Slater, <i>On the equation $\varphi(x) = \int_x^{x+1} K(\xi) f[\varphi(\xi)] d\xi$</i>	155
Arthur William John Stoddart, <i>Existence of optimal controls</i>	167
Burnett Roland Toskey, <i>A system of canonical forms for rings on a direct sum of two infinite cyclic groups</i>	179
Jerry Eugene Vaughan, <i>A modification of Morita's characterization of dimension</i>	189