

Pacific Journal of Mathematics

**ON OPERATORS WHOSE SPECTRUM LIES ON A CIRCLE OR
A LINE**

HERBERT MEYER KAMOWITZ

ON OPERATORS WHOSE SPECTRUM LIES ON A CIRCLE OR A LINE

HERBERT KAMOWITZ

The purpose of this note is to prove the following theorem.

THEOREM. Let N be a bounded linear operator on Hilbert space H satisfying

$$(1) \quad \|NT - TN\| = \|N^*T - TN^*\|$$

for all bounded linear operators T . Then N is (obviously) normal and the spectrum of N lies on a circle or straight line.

Here N^* denotes the adjoint of the operator N .

It is clear that if S is a unitary or self-adjoint operator and α and β are complex numbers, then $N = \alpha I + \beta S$ satisfies (1). The theorem asserts that the converse is also true.

After noting that in dimensions two and three the theorem is trivially true, we proceed to the first of two parts of the proof.

I. Dimension four. Let H be four dimensional Euclidean space. Since a normal operator N is unitarily equivalent to a diagonal matrix it is no restriction to assume that N has the form

$$\begin{pmatrix} \lambda_1 & 0 & 0 & 0 \\ 0 & \lambda_2 & 0 & 0 \\ 0 & 0 & \lambda_3 & 0 \\ 0 & 0 & 0 & \lambda_4 \end{pmatrix}$$

where the λ_i 's are the eigenvalues of N . Consider the matrix

$$T = \begin{pmatrix} 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & i \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Then

$$NT - TN = \begin{pmatrix} 0 & 0 & (\lambda_1 - \lambda_3) & (\lambda_1 - \lambda_4) \\ 0 & 0 & (\lambda_2 - \lambda_3) & i(\lambda_2 - \lambda_4) \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

and $\|NT - TN\|^2$ is the largest eigenvalue of the matrix

$$A = \begin{pmatrix} |\lambda_1 - \lambda_3|^2 + |\lambda_1 - \lambda_4|^2 & (\lambda_1 - \lambda_3)\overline{(\lambda_2 - \lambda_3)} & 0 & 0 \\ & -i(\lambda_1 - \lambda_4)\overline{(\lambda_2 - \lambda_4)} & 0 & 0 \\ \overline{(\lambda_1 - \lambda_3)}(\lambda_2 - \lambda_3) & & |\lambda_2 - \lambda_3|^2 + |\lambda_2 - \lambda_4|^2 & 0 & 0 \\ + i\overline{(\lambda_1 - \lambda_4)}(\lambda_2 - \lambda_4) & & & & \\ 0 & & 0 & & 0 & 0 \\ 0 & & 0 & & 0 & 0 \end{pmatrix}.$$

On the other hand, $\|N^*T - TN^*\|^2$ is the largest eigenvalue of a matrix B which has the same form as A but with λ_i replaced by $\bar{\lambda}_i$, $i = 1, 2, 3, 4$.

Let $P(X)$ and $P^*(X)$ be the characteristic polynomials of A and B , respectively. Then from the form of the matrices A and B one sees that the polynomials $P(X)$ and $P^*(X)$ differ only in the coefficient of X^2 . Therefore $\|NT - TN\| = \|N^*T - TN^*\|$ if, and only if, $P(X)$ and $P^*(X)$ are equal and a routine computation shows that this holds if, and only if,

$$\begin{aligned} & [(\lambda_1 - \lambda_3)(\lambda_2 - \lambda_3) + i\overline{(\lambda_1 - \lambda_4)}(\lambda_2 - \lambda_4)] \\ & \quad \times [(\lambda_1 - \lambda_3)\overline{(\lambda_2 - \lambda_3)} - i(\lambda_1 - \lambda_4)\overline{(\lambda_2 - \lambda_4)}] \\ & = [(\lambda_1 - \lambda_3)\overline{(\lambda_2 - \lambda_3)} + i\overline{(\lambda_1 - \lambda_4)}(\lambda_2 - \lambda_4)] \\ & \quad \times [(\lambda_1 - \lambda_3)(\lambda_2 - \lambda_3) - i\overline{(\lambda_1 - \lambda_4)}(\lambda_2 - \lambda_4)] \end{aligned}$$

which reduces to the condition that $\overline{(\lambda_1 - \lambda_4)}(\lambda_2 - \lambda_3)(\lambda_1 - \lambda_3)(\lambda_2 - \lambda_4)$ be real. The latter holds if, and only if, $\lambda_1, \lambda_2, \lambda_3, \lambda_4$ lie on a line or circle, which proves the theorem in dimension four.

Let (a_1, a_2, a_3, a_4) be a vector in four dimensional H . Since

$$\begin{aligned} & (NT - TN)(a_1, a_2, a_3, a_4) \\ & = ((\lambda_1 - \lambda_3)a_3 + (\lambda_1 - \lambda_4)a_4, (\lambda_2 - \lambda_3)a_3 + i(\lambda_2 - \lambda_4)a_4, 0, 0), \end{aligned}$$

we remark that we have just shown that

$$(2) \quad \sup_{\sum |a_i|^2 = 1} [|(\lambda_1 - \lambda_3)a_3 + (\lambda_1 - \lambda_4)a_4|^2 + |(\lambda_2 - \lambda_3)a_3 + i(\lambda_2 - \lambda_4)a_4|^2]$$

is the largest root of $P(X) = 0$, and also that

$$(3) \quad \sup_{\sum |a_i|^2 = 1} [|\overline{(\lambda_1 - \lambda_3)}a_3 + \overline{(\lambda_1 - \lambda_4)}a_4|^2 + |(\lambda_2 - \lambda_3)a_3 + i\overline{(\lambda_2 - \lambda_4)}a_4|^2]$$

is the largest root of $P^*(X) = 0$.

II. Higher dimensions. Now let H be a finite or infinite dimensional Hilbert space of dimension > 4 . Let N be a normal operator satisfying $\|NT - TN\| = \|N^*T - TN^*\|$ for all T . Suppose N has spectrum S and spectral representation $N = \int_S \lambda dE_\lambda$. Let

$\lambda_1, \lambda_2, \lambda_3, \lambda_4$ be distinct points of S . (If S has no more than three points then the theorem is trivially true.) Choose an $\varepsilon > 0$ but smaller than the minimum of the distances $|\lambda_i - \lambda_j|$, $i \neq j$; let $S_k = S \cap \{\lambda \mid |\lambda - \lambda_k| < \varepsilon\}$, $k = 1, 2, 3, 4$. Let $E_k = \int_{S_k} dE_\lambda$ and choose $y \in H$ with $\|E_k y\| = 1$ for $k = 1, 2, 3, 4$ and let $x_k = E_k y$. Then $(x_i, x_j) = 0$ if $i \neq j$ and each x_k is an approximate eigenvector in the sense that

$$\begin{aligned} \|Nx_k - \lambda_k x_k\| &= \left\| \int_{S_k} \lambda dE_\lambda x_k - \lambda_k \int_{S_k} dE_\lambda x_k \right\| \\ &\leq \max_{\lambda \in S_k} |\lambda - \lambda_k| < \varepsilon. \end{aligned}$$

We can therefore write $Nx_k = \lambda_k x_k + \varepsilon u_k$ where $\|u_k\| \leq 1$. Similarly $N^* x_k = \bar{\lambda}_k x_k + \varepsilon u'_k$, $\|u'_k\| \leq 1$.

Define an operator T on H by

$$Tx = [(E_3 x, x_3) + (E_4 x, x_4)]x_1 + [(E_3 x, x_3) + i(E_4 x, x_4)]x_2.$$

Now for $x \in H$,

$$\begin{aligned} NTx &= [(E_3 x, x_3) + (E_4 x, x_4)]\lambda_1 x_1 \\ &\quad + [(E_3 x, x_3) + i(E_4 x, x_4)]\lambda_2 x_2 + \delta_1(x), \end{aligned}$$

where $\|\delta_1(x)\| \leq 4\varepsilon \|x\|$. Also

$$\begin{aligned} T(Nx) &= [(E_3 Nx, x_3) + (E_4 Nx, x_4)]x_1 \\ &\quad + [(E_3 Nx, x_3) + i(E_4 Nx, x_4)]x_2. \end{aligned}$$

Since $NE_k = E_k N$, $k = 1, 2, 3, 4$, it follows that

$$\begin{aligned} TNx &= [(E_3 x, \bar{\lambda}_3 x_3) + (E_4 x, \bar{\lambda}_4 x_4)]x_1 \\ &\quad + [(E_3 x, \bar{\lambda}_3 x_3) + i(E_4 x, \bar{\lambda}_4 x_4)]x_2 + \delta_2(x), \end{aligned}$$

where $\|\delta_2(x)\| \leq 4\varepsilon \|x\|$.

Thus

$$\begin{aligned} \|(NT - TN)x\| &= \|[(\lambda_1 - \lambda_3)(E_3 x, x_3) + (\lambda_1 - \lambda_4)(E_4 x, x_4)]x_1 \\ &\quad + [(\lambda_2 - \lambda_3)(E_3 x, x_3) + i(\lambda_2 - \lambda_4)(E_4 x, x_4)]x_2 + \gamma_1(x)\|, \end{aligned}$$

where $\|\gamma_1(x)\| \leq 8\varepsilon \|x\|$.

Similarly

$$\begin{aligned} \|(N^*T - TN^*)x\| &= \|[(\bar{\lambda}_1 - \bar{\lambda}_3)(E_3 x, x_3) + (\bar{\lambda}_1 - \bar{\lambda}_4)(E_4 x, x_4)]x_1 \\ &\quad + [(\bar{\lambda}_2 - \bar{\lambda}_3)(E_3 x, x_3) + i(\bar{\lambda}_2 - \bar{\lambda}_4)(E_4 x, x_4)]x_2 + \gamma_2(x)\|, \end{aligned}$$

where $\|\gamma_2(x)\| \leq 8\varepsilon \|x\|$.

Since x_1 and x_2 are orthogonal,

$$\begin{aligned} & \|NT - TN\|^2 \\ &= \sup_{\|x\|=1} [|(\lambda_1 - \lambda_3)(E_3x, x_3) + (\lambda_1 - \lambda_4)(E_4x, x_4)|^2 \\ &\quad + |(\lambda_2 - \lambda_3)(E_3x, x_3) + i(\lambda_2 - \lambda_4)(E_4x, x_4)|^2] + \delta \end{aligned}$$

and

$$\begin{aligned} & \|N^*T - TN^*\|^2 \\ &= \sup_{\|x\|=1} [|(\overline{\lambda_1 - \lambda_3})(E_3x, x_3) + (\overline{\lambda_1 - \lambda_4})(E_4x, x_4)|^2 \\ &\quad + |(\overline{\lambda_2 - \lambda_3})(E_3x, x_3) + i(\overline{\lambda_2 - \lambda_4})(E_4x, x_4)|^2] + \delta', \end{aligned}$$

where $\delta, \delta' \leq 8\varepsilon$.

Thus if $\|NT - TN\| = \|N^*T - TN^*\|$, we obtain by (2) and (3) that the largest zeros of $P(X)$ and $P^*(X)$ differ by at most 16ε . Therefore all the coefficients of $P(X)$ and $P^*(X)$ are close to each other and in particular the imaginary part of

$$(\overline{\lambda_1 - \lambda_4})(\overline{\lambda_2 - \lambda_3})(\lambda_1 - \lambda_3)(\lambda_2 - \lambda_4)$$

is bounded by a function of ε which goes to zero with ε . Letting $\varepsilon \rightarrow 0$ we again find that $(\overline{\lambda_1 - \lambda_4})(\overline{\lambda_2 - \lambda_3})(\lambda_1 - \lambda_3)(\lambda_2 - \lambda_4)$ must be real and so $\lambda_1, \lambda_2, \lambda_3$ and λ_4 lie on a line or a circle. Since the λ_i 's were arbitrary, the proof is concluded.

Received April 27, 1965.

AVCO CORPORATION
WILMINGTON, MASSACHUSETTS

UNIVERSITY OF MASSACHUSETTS
BOSTON, MASSACHUSETTS

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California

J. P. JANS

University of Washington
Seattle, Washington 98105

J. DUGUNDJI

University of Southern California
Los Angeles, California 90007

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Pacific Journal of Mathematics

Vol. 20, No. 1

September, 1967

Leonard Daniel Baumert, <i>Extreme copositive quadratic forms. II</i>	1
Edward Lee Bethel, <i>A note on continuous collections of disjoint continua</i>	21
Delmar L. Boyer and Adolf G. Mader, <i>A representation theorem for abelian groups with no elements of infinite p-height</i>	31
Jean-Claude B. Derderian, <i>Residuated mappings</i>	35
Burton I. Fein, <i>Representations of direct products of finite groups</i>	45
John Brady Garnett, <i>A topological characterization of Gleason parts</i>	59
Herbert Meyer Kamowitz, <i>On operators whose spectrum lies on a circle or a line</i>	65
Ignacy I. Kotlarski, <i>On characterizing the gamma and the normal distribution</i>	69
Yu-Lee Lee, <i>Topologies with the same class of homeomorphisms</i>	77
Moshe Mangad, <i>Asymptotic expansions of Fourier transforms and discrete polyharmonic Green's functions</i>	85
Jürg Thomas Marti, <i>On integro-differential equations in Banach spaces</i>	99
Walter Philipp, <i>Some metrical theorems in number theory</i>	109
Maxwell Alexander Rosenlicht, <i>Another proof of a theorem on rational cross sections</i>	129
Kenneth Allen Ross and Karl Robert Stromberg, <i>Jessen's theorem on Riemann sums for locally compact groups</i>	135
Stephen Simons, <i>A theorem on lattice ordered groups, results of Ptak, Namioka and Banach, and a front-ended proof of Lebesgue's theorem</i>	149
Morton Lincoln Slater, <i>On the equation $\varphi(x) = \int_x^{x+1} K(\xi) f[\varphi(\xi)] d\xi$</i>	155
Arthur William John Stoddart, <i>Existence of optimal controls</i>	167
Burnett Roland Toskey, <i>A system of canonical forms for rings on a direct sum of two infinite cyclic groups</i>	179
Jerry Eugene Vaughan, <i>A modification of Morita's characterization of dimension</i>	189