

Pacific Journal of Mathematics

ON CHARACTERIZING THE GAMMA AND THE NORMAL DISTRIBUTION

IGNACY I. KOTLARSKI

ON CHARACTERIZING THE GAMMA AND THE NORMAL DISTRIBUTION

IGNACY KOTLARSKI

We will characterize the gamma distribution by the nature of the joint distribution of the two quotients $X_1/X_3, X_2/X_3$ for three identically gamma distributed random variables.

It is well known that if two independent identically distributed random variables X_1, X_2 have the gamma distribution given by the density

$$(1) \quad f(x) = \begin{cases} 0 & \text{for } x \leq 0 \\ \frac{a^p}{\Gamma(p)} x^{p-1} e^{-ax} & \text{for } x > 0 \end{cases} \quad \begin{matrix} (a > 0) \\ (p > 0) \end{matrix}$$

then their quotient

$$(2) \quad Y = X_1/X_2$$

has the beta distribution of the second kind given by the density

$$(3) \quad g(y) = \begin{cases} 0 & \text{for } y \leq 0 \\ \frac{1}{B(p, p)} \cdot \frac{y^{p-1}}{(1+y)^{2p}} & \text{for } y > 0. \end{cases}$$

However, this property does not characterize the gamma distribution uniquely. There exist pairs of independent positive identically distributed random variables X_1, X_2 whose common distribution function $F(x)$ differs from the one given by the density (1), but where the quotients (2) are distributed according to the density (3). Some such distribution functions $F(x)$ are given by the following densities

$$(4) \quad \begin{aligned} f_1(x) &= \begin{cases} 0 & \text{for } x \leq 0 \\ \frac{a^p}{\Gamma(p)} x^{p-1} e^{-a/x} & \text{for } x > 0 \end{cases} \\ f_2(x) &= \begin{cases} 0 & \text{for } x \leq 0 \\ \frac{2\Gamma\left(\frac{2p+1}{2}\right)}{\Gamma\left(\frac{p}{2}\right)\Gamma\left(\frac{p+1}{2}\right)} \cdot \frac{x^{p-1}}{(1+x^2)^{p+1/2}} & \text{for } x > 0 \end{cases} \end{aligned}$$

$$f_3(x) = \begin{cases} 0 & \text{for } x \leq 0 \\ \frac{2\Gamma\left(\frac{2p+1}{2}\right)}{\Gamma\left(\frac{p}{2}\right)\Gamma\left(\frac{p+1}{2}\right)} \cdot \frac{x^p}{(1+x^2)^{p+1/2}} & \text{for } x > 0. \end{cases}$$

Now let X_1, X_2, X_3 , be three independent positive identically distributed random variables whose common distribution function is $F(x)$. It is interesting that although the distribution of one quotient (2) does not characterize $F(x)$ uniquely, the joint distribution of two such quotients

$$(5) \quad Y_1 = X_1/X_3, \quad Y_2 = X_2/X_3$$

characterize (by some assumptions) $F(x)$ uniquely, up to a change of the scale.

The condition of identical distributions of X_1, X_2, X_3 may be omitted. The distribution functions $F_k(x)$ of the independent positive random variables $X_k (k = 1, 2, 3)$ are uniquely characterized (by some assumptions) by the joint distribution of the pair of quotients (5), up to a change of the scale.

The normal distribution is also characterized in the same way. There are also some generalizations of these problems.

1. Some lemmas.

LEMMA 1. *Let X_1, X_2, X_3 be three independent real random variables, and let*

$$(6) \quad Z_1 = X_1 - X_3, \quad Z_2 = X_2 - X_3.$$

If the characteristic function of the pair (Z_1, Z_2) does not vanish, then the distribution of (Z_1, Z_2) determines the distributions of X_1, X_2, X_3 up to a change of the location.

Proof. Denote $\varphi(t_1, t_2)$ the characteristic function of the pair (Z_1, Z_2) , and $\varphi_k(t)$ the characteristic functions of $X_k (k = 1, 2, 3)$. Then there is

$$(7) \quad \begin{aligned} \varphi(t_1, t_2) &= E \exp [i(t_1 Z_1 + t_2 Z_2)] \\ &= E \exp [i\{t_1(X_1 - X_3) + t_2(X_2 - X_3)\}] \\ &= E \exp [i\{t_1 X_1 + t_2 X_2 + (-t_1 - t_2)X_3\}] \\ &= \varphi_1(t_1)\varphi_2(t_2)\varphi_3(-t_1 - t_2). \end{aligned}$$

The condition of nonvanishing of $\varphi(t_1, t_2)$ is equivalent to nonvanishing of any of the functions $\varphi_k(t) (k = 1, 2, 3)$.

Let U_1, U_2, U_3 be another three independent real random variables, having characteristic functions $\psi_k(t) = E e^{it\psi_k} (k = 1, 2, 3)$ and satisfying the assumptions of Lemma 1. Let $V_1 = U_1 - U_3, V_2 = U_2 - U_3$ and $\psi(t_1, t_2) = E \exp [i(t_1 V_1 + t_2 V_2)]$. Making the same considerations as for formula (7) it is easy to see that there is

$$(8) \quad \psi(t_1, t_2) = \psi_1(t_1)\psi_2(t_2)\psi_3(-t_1 - t_2).$$

Let the pairs (Z_1, Z_2) and (V_1, V_2) have the same distribution. Then their characteristic functions are equal. Using formulae (7) and (8) one can obtain the following equation

$$(9) \quad \psi_1(t_1)\psi_2(t_2)\psi_3(-t_1 - t_2) = \varphi_1(t_1)\varphi_2(t_2)\varphi_3(-t_1 - t_2) \\ (-\infty < t_1 < +\infty, -\infty < t_2 < +\infty).$$

Put

$$(10) \quad \psi_k(t) = \varphi_k(t) \cdot p_k(t) \quad (k = 1, 2, 3).$$

Putting (10) into the equation (9) one obtains the following equation

$$(11) \quad p_1(t_1)p_2(t_2)p_3(-t_1 - t_2) = 1 \left(\begin{array}{l} -\infty < t_1 < +\infty \\ -\infty < t_2 < +\infty \end{array} \right)$$

in which $p_k(t)$ are unknown functions; they are complex functions, continuous on the whole line $-\infty < t < +\infty$, satisfying the condition

$$(12) \quad p_k(0) = 1 \quad (k = 1, 2, 3).$$

In order to solve the equation (11) let us put $t_1 = t, t_2 = 0$ and, later $t_2 = t, t_1 = 0$. Then, using (12),

$$(13) \quad p_1(t) \cdot p_3(-t) = 1, \quad p_2(t) \cdot p_3(-t) = 1.$$

Putting $p_1(t)$ and $p_2(t)$ given by (13) into the equation (11) and changing signs, we obtain

$$(14) \quad p_3(t_1 + t_2) = p_3(t_1) \cdot p_3(t_2).$$

The only continuous function which satisfies the equation (14) and condition (12) is the exponential function

$$(15) \quad p_3(t) = e^{ct} \quad (-\infty < t < +\infty)$$

where c is a complex number. Putting (15) into (13) it is easy to see that

$$(16) \quad p_1(t) = p_2(t) = p_3(t) = e^{ct}.$$

Putting (16) into (10) there is

$$(17) \quad \psi_k(t) = e^{ct} \varphi_k(t) \quad (k = 1, 2, 3).$$

Since the known property of characteristic functions $\varphi(-t) = \overline{\varphi(t)}$ the formulae (17) become

$$(18) \quad \psi_k(t) = e^{ibt} \varphi_k(t) \quad (k = 1, 2, 3)$$

where b is a real constant. This means that the distributions of X_k are the same as of U_k up to a change of the location.

Remark 1. The assumption of nonvanishing of the joint characteristic function of the differences (6) may be replaced by the assumption that all X_k have analytic characteristic functions. All considerations are valid in such a case for t complex, being inside a circle $|t| < t_0$ ($t_0 > 0$), where the characteristic functions do not vanish. Because of the analyticity of the characteristic functions the formulae (18) may be spread on the whole real line.

Remark 2. If the assumption of nonvanishing of the characteristic functions is omitted then the theorem becomes false. In order to show this the following example is given.

Exemplé. Let (X_1, X_2, X_3) and (U_1, U_2, U_3) be two three's of independent real random variables having their characteristic functions $\varphi_k(t)$ and $\psi_k(t)$ respectively. Let be

$$(19) \quad \begin{cases} \varphi_1(t) = \varphi_2(t) = \psi_1(t) = \psi_2(t) = \begin{cases} 0 & \text{for } |t| > 1 \\ 1 - |t| & \text{for } |t| \leq 1 \end{cases} \\ \varphi_3(t) = \begin{cases} 0 & \text{for } |t| > 2 \\ 1 - |t|/2 & \text{for } |t| \leq 2 \end{cases} \\ \psi_3(t) = \varphi_3(t) \text{ for } |t| \leq 2, \psi_3(t+4) = \psi_3(t). \end{cases}$$

It is easy to see that in such a case the equation (9) holds though $\psi_3(t)$ is not the same as $\varphi_3(t)$. Hence there exist trios (X_1, X_2, X_3) and (U_1, U_2, U_3) of independent real random variables whose distributions are not the same, but where the distributions of the corresponding pairs of differences $(X_1 - X_3, X_2 - X_3)$, $(U_1 - U_3, U_2 - U_3)$ are the same.

Remark 3. Lemma 1 remains true if the three independent real random variables (X_1, X_2, X_3) are replaced by n such random variables (X_1, X_2, \dots, X_n) ($n \geq 3$), and the pair of differences $(X_1 - X_3, X_2 - X_3)$ by $n - 1$ differences $(X_1 - X_n, X_2 - X_n, \dots, X_{n-1} - X_n)$.

Remark 4. The differences in formula (6) may be replaced by sums.

Remark 5. Lemma 1 remains true if the real random variables X_k are replaced by n -dimensional real random vectors.

LEMMA 2. *Let X_1, X_2, X_3 be three independent positive random variables, and let (Y_1, Y_2) be their quotients given by formula (5). If the joint characteristic function of the pair $(\ln Y_1, \ln Y_2)$ does not vanish, then the distribution of (Y_1, Y_2) determines the distributions of X_1, X_2, X_3 up to change of the scale.*

Proof. The proof of Lemma 2 is obvious because $\ln X_k$ ($k = 1, 2, 3$) satisfy the assumptions of Lemma 1.

REMARK 6. The positive random variables X_k in Lemma 2 may be replaced by symmetrical about the origin real random variables satisfying the condition $P(X_k = 0) = 0$.

2. **Characterizing the gamma distribution.** The problem of characterizing the gamma distribution of two independent random variables X_1, X_2 by the distribution of their quotient was first posed by J. G. Mauldon [9], he showed that there is no such characterization. Further investigations on this problem were made by I. Kotlarski [1]. A full treatment of the problem has been made by R. G. Laha [6] (on this subject see also E. Lukacs, R. G. Laha [7], p. 59). The authors of [1] and [6] searched for the properties of the set of distribution functions $F(x)$ for which the quotient (2) is distributed according to (3), where X_1, X_2 are independent positive random variables identically distributed according to $F(x)$.

THEOREM 1. *Let X_1, X_2, X_3 be three independent positive random variables, let (Y_1, Y_2) be the pair of the quotients given by (5). The necessary and sufficient condition for X_k to be gamma distributed with parameters p_k and a (a - common, $k = 1, 2, 3$) is that the joint distribution of (Y_1, Y_2) is the bivariate beta distribution of the second kind given by the density*

$$(20) \quad g(y_1, y_2) = \begin{cases} \frac{\Gamma(p_1 + p_2 + p_3)}{\Gamma(p_1)\Gamma(p_2)\Gamma(p_3)} \cdot \frac{y_1^{p_1-1}y_2^{p_2-1}}{(1 + y_1 + y_2)^{p_1+p_2+p_3}} & \text{for } y_1 > 0 \\ & y_2 > 0 \\ 0 & \text{elsewhere} \end{cases}$$

Proof. The characteristic function of $\ln X_k$ where X_k is gamma

distributed with parameters p_k , a , is given by

$$(21) \quad \varphi_k(t) = E e^{it \ln X_k} = a^{-it} \frac{\Gamma(p_k + it)}{\Gamma(p_k)}.$$

The characteristic function of the pair $(\ln Y_1, \ln Y_2)$ is

$$(22) \quad \begin{aligned} \varphi(t_1, t_2) &= \varphi_1(t_1) \varphi_2(t_2) \varphi_3(-t_1 - t_2) \\ &= \frac{\Gamma(p_1 + it_1)}{\Gamma(p_1)} \cdot \frac{\Gamma(p_2 + it_2)}{\Gamma(p_2)} \cdot \frac{\Gamma(p_3 - it_1 - it_2)}{\Gamma(p_3)} \end{aligned}$$

it is easy to see that the characteristic function of $(\ln Y_1, \ln Y_2)$ where (Y_1, Y_2) is distributed according to the density (20) is also given by the right side of the formula (22), This ends the proof.

THEOREM 2. *Let X_1, X_2, X_3 be three independent positive random variables, let (U_1, U_2) be given by formulae*

$$(23) \quad U_1 = \frac{X_1}{X_1 + X_2}, \quad U_2 = \frac{X_1 + X_2}{X_1 + X_2 + X_3}$$

The necessary and sufficient condition for X_k to be gamma distributed with parameters p_k and a (a - common, $k = 1, 2, 3$) is that U_1, U_2 are independent beta distributed random variables, U_1 with parameters (p_1, p_2) , and U_2 with parameters $(p_1 + p_2, p_3)$.

Proof. The necessary condition is obvious. In order to prove the sufficient condition let us put

$$(24) \quad U_1 = \frac{Y_1}{Y_1 + Y_2}, \quad U_2 = \frac{Y_1 + Y_2}{1 + Y_1 + Y_2}$$

where Y_1 and Y_2 are given by (5). It is easy to see that if U_1, U_2 are independent, U_1 being distributed according to the beta law of the first kind with the density

$$(25) \quad g_1(u) = \begin{cases} \frac{1}{B(p_1, p_2)} u^{p_1-1} (1-u)^{p_2-1} & \text{for } 0 < u < 1 \\ 0 & \text{elsewhere} \end{cases}$$

and U_2 has a similar distribution with parameters $(p_1 + p_2, p_3)$, then (Y_1, Y_2) is distributed according to the density (20). Using Theorem 1 it is easy to see that X_k are gamma distributed with parameters (p_k, a) , (a - common, $k = 1, 2, 3$). This ends the proof.

3. Characterizing the normal distribution. The problem of characterizing the normal distribution of two independent symmetric about the origin random variables by the distribution of their quotient

has been considered by several authors. J. G. Mauldon [9] showed that there is no such characterization. Further investigation on this problem were made by R. G. Laha [3, 4, 5], I. Kotlarski [2], G. P. Steck [10].

Though there is no characterization of two independent symmetrical about the origin random variables by the distribution of their quotient, there is a characterization of three such random variables by the joint distribution of two their quotients. In this Section two theorems on characterizing the normal distribution in such a way are presented.

THEOREM 3. *Let X_1, X_2, X_3 be three independent real symmetrical about the origin random variables satisfying the condition $P(X_k = 0) = 0$, ($k = 1, 2, 3$). Let (Y_1, Y_2) be the pair of quotients given by formula (5). The necessary and sufficient condition for X_k to be normal distributed with a common standard deviation σ is that the joint distribution of (Y_1, Y_2) is the bivariate Cauchy distribution given by the density*

$$(26) \quad g(y_1, y_2) = \frac{1}{2\pi} \cdot \frac{1}{(1 + y_1^2 + y_2^2)^{3/2}} \quad \left(\begin{array}{l} -\infty < y_1 < +\infty \\ -\infty < y_2 < +\infty \end{array} \right)$$

Proof. The characteristic function of $\ln |X_k|$ where X_k is normal distributed with zero mean and standard deviation σ is

$$(27) \quad \varphi_k(t) = E \exp it \ln |X_k| = (\sigma\sqrt{2})^{it} \frac{\Gamma((1+it)/2)}{\sqrt{\pi}}$$

The characteristic function of the pair $(\ln |Y_1|, \ln |Y_2|)$ is

$$(28) \quad \begin{aligned} \varphi(t_1, t_2) &= \varphi_1(t_1)\varphi_2(t_2)\varphi_3(-t_1 - t_2) \\ &= \frac{1}{\pi^{3/2}} \Gamma\left(\frac{1+it_1}{2}\right) \Gamma\left(\frac{1+it_2}{2}\right) \Gamma\left(\frac{1-i(t_1+t_2)}{2}\right). \end{aligned}$$

It is easy to see that the characteristic function of $(\ln |Y_1|, \ln |Y_2|)$ where (Y_1, Y_2) is distributed according to the density (26) is also given by the right side of the formula (28). This ends the proof.

THEOREM 4. *Let X_1, X_2, X_3 be three independent real symmetrical about the origin random variables satisfying the condition $P(X_k = 0) = 0$ ($k = 1, 2, 3$). Denote*

$$(29) \quad V_1 = \frac{X_1}{\sqrt{X_1^2 + X_2^2}}, \quad V_2 = \frac{\sqrt{X_1^2 + X_2^2}}{\sqrt{X_1^2 + X_2^2 + X_3^2}}$$

The necessary and sufficient condition for X_k to be normal distri-

buted with a common standard deviation is that V_1 and V_2 are independent distributed according to the densities

$$(30) \quad h_1(v) = \begin{cases} \frac{1}{\pi} \cdot \frac{1}{\sqrt{1-v^2}} & \text{for } |v| < 1 \\ 0 & \text{elsewhere} \end{cases}$$

$$h_2(v) = \begin{cases} \frac{v}{\sqrt{1-v^2}} & \text{for } 0 < v < 1 \\ 0 & \text{elsewhere.} \end{cases}$$

Proof. The necessary condition is obvious. In order to prove the sufficient condition let us put

$$(31) \quad V_1 = \frac{Y_1}{\sqrt{Y_1^2 + Y_2^2}}, \quad V_2 = \frac{\sqrt{Y_1^2 + Y_2^2}}{\sqrt{1 + Y_1^2 + Y_2^2}}$$

where Y_1, Y_2 are given by formula (5). It is easy to see that if (V_1, V_2) are independently distributed according to (30), then (Y_1, Y_2) is distributed according to the density (26). Using Theorem 3 it is easy to see that all X_k are normal distributed with a common standard deviation. This ends the proof.

REFERENCES

1. I. Kotlarski, *On pairs of independent random variables whose quotients follow some known distribution*, Colloquium Mathematicum, Vol. IX, 1962, Fasc. 1, 151-162.
2. ———, *On random variables whose quotient follows the Cauchy law*, Colloquium Mathematicum, Vol. VII, 1960, Fasc. 2, 273-284.
3. R. G. Laha, *An example of a non-normal distribution where the quotient follows the Cauchy law*, Proceedings of the National Academy of Sciences **44** (1958), 222-223.
4. ———, *On a class of distribution functions where the quotient follows the Cauchy law*, Transactions of the American Mathematical Society **93** (1959), 205-215.
5. ———, *On the laws of Cauchy and Gauss*, Ann. Math. Statist. **30** (1959) 1165-1174.
6. ———, *On a problem connected with beta and gamma distributions*, Transactions of the American Mathematical Society, Vol. 113, No. 2, November 1964, 287-298.
7. E. Lukacs and R. G. Laha, *Application of Characteristic Functions*, London, 1964.
8. A. J. Markushevitch, *Teoria analititsheskich funkcij*, Moskwa, 1950, (*The Theory of Analytical Functions*, Moscow, 1950, Russian).
9. J. G. Mauldon, *Characterizing properties of statistical distributions*, Quarterly Journal of Mathematics, Oxford Series, 2.7, 1956, 155-160.
10. G. P. Steck, *A uniqueness property not enjoyed by the normal distribution*, Ann. Math. Statist. **29** (1958) 604-606.

Received February 26, 1965.

POLITECHNIKA WARSZAWSKA, POLAND

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California

J. P. JANS

University of Washington
Seattle, Washington 98105

J. DUGUNDJI

University of Southern California
Los Angeles, California 90007

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Leonard Daniel Baumert, <i>Extreme copositive quadratic forms. II</i>	1
Edward Lee Bethel, <i>A note on continuous collections of disjoint continua</i>	21
Delmar L. Boyer and Adolf G. Mader, <i>A representation theorem for abelian groups with no elements of infinite p-height</i>	31
Jean-Claude B. Derderian, <i>Residuated mappings</i>	35
Burton I. Fein, <i>Representations of direct products of finite groups</i>	45
John Brady Garnett, <i>A topological characterization of Gleason parts</i>	59
Herbert Meyer Kamowitz, <i>On operators whose spectrum lies on a circle or a line</i>	65
Ignacy I. Kotlarski, <i>On characterizing the gamma and the normal distribution</i>	69
Yu-Lee Lee, <i>Topologies with the same class of homeomorphisms</i>	77
Moshe Mangad, <i>Asymptotic expansions of Fourier transforms and discrete polyharmonic Green's functions</i>	85
Jürg Thomas Marti, <i>On integro-differential equations in Banach spaces</i>	99
Walter Philipp, <i>Some metrical theorems in number theory</i>	109
Maxwell Alexander Rosenlicht, <i>Another proof of a theorem on rational cross sections</i>	129
Kenneth Allen Ross and Karl Robert Stromberg, <i>Jessen's theorem on Riemann sums for locally compact groups</i>	135
Stephen Simons, <i>A theorem on lattice ordered groups, results of Ptak, Namioka and Banach, and a front-ended proof of Lebesgue's theorem</i>	149
Morton Lincoln Slater, <i>On the equation $\varphi(x) = \int_x^{x+1} K(\xi) f[\varphi(\xi)] d\xi$</i>	155
Arthur William John Stoddart, <i>Existence of optimal controls</i>	167
Burnett Roland Toskey, <i>A system of canonical forms for rings on a direct sum of two infinite cyclic groups</i>	179
Jerry Eugene Vaughan, <i>A modification of Morita's characterization of dimension</i>	189