

# Pacific Journal of Mathematics

**ON ASYMPTOTIC ESTIMATES FOR KERNELS OF  
CONVOLUTION TRANSFORMS**

ZEEV DITZIAN

## ON ASYMPTOTIC ESTIMATES FOR KERNELS OF CONVOLUTION TRANSFORMS

ZEEV DITZIAN

**In this paper we shall try to answer two open questions posed by Dauns and Widder in their paper "Convolution transforms whose inversion functions have complex roots" (Pacific Journal of Mathematics, 1965, Volume 15(2), pp. 427-442) on page 441.**

We shall be interested in the function  $G_{2m}(t)$  defined by

$$(1.1) \quad G_{2m}(t) = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} \frac{e^{st} ds}{E_{2m}(s)}$$

where

$$(1.2) \quad E_{2m}(s) = \prod_{k=m+1}^{\infty} \left(1 - \frac{s^2}{a_k^2}\right)$$

where  $\{a_k\}$  is a sequence of complex numbers such that

$$|\arg a_k| < \frac{\pi}{4} - \eta$$

for a fixed  $\eta$ ,  $0 < \eta < \pi/4$ ,

$$\sum_k |a_k|^{-2} < \infty, \quad 0 < \operatorname{Re} a_i \leq \operatorname{Re} a_{i+1} \text{ for all } i$$

and

$$(1.3) \quad \lim_{n \rightarrow \infty} |a_{n+1}|^2 \sum_{k=m+1}^{\infty} |a_k|^{-2} = \infty.$$

If a sequence  $\{a_k\}$  satisfies all the above assumptions, we shall denote it by  $\{a_k\} \in \text{class } C$ . We obtain condition  $B$ , defined in [1, p. 436], if we replace (1.3) by (1.4)

$$(1.4) \quad \lim_{n \rightarrow \infty} |a_{n+1}|^{4/3} \sum_{n+1}^{\infty} |a_k|^{-2} = \infty.$$

If we take  $a_k = k^\lambda 1/2 < \lambda < \infty$  then  $\{a_k\} \in \text{class } C$ , but of these sequences only those for which  $1/2 < \lambda < 3/2$  satisfy condition  $B$ .

We define as in [1]

$$(1.5) \quad V_m = \sum_{k=m+1}^{\infty} a_k^{-2} \quad \text{and} \quad S_m = \sum_{k=m+1}^{\infty} |a_k|^{-2}$$

and whenever  $\{a_k\} \in \text{class } C$  we prove

$$(1.6) \quad \lim_{m \rightarrow \infty} \int_{-\infty}^{\infty} |tG'_{2m}(t)| dt = (\cos^2 \varphi_m - \sin^2 \varphi_m)^{-3/2}$$

where

$$\varphi_m = \frac{1}{2} \arg V_m \left( -\frac{\pi}{2} < \arg V_m < \frac{\pi}{2} \right)$$

which answers the question posed in remark (3) [1, p. 441].

We shall also prove under the restriction  $\{a_k\} \in \text{class } C$  Corollary 4.3 and an analogous theorem to Theorem 4.1.

As a by product we shall have

$$(1.7) \quad \lim_{m \rightarrow \infty} S_m^{1/2} \frac{d^n}{dt^n} G_{2m}(S_m^{1/2}t) = \frac{1}{\sqrt{4\pi}} \left( \frac{S_m}{V_m} \right)^{1/2} \frac{d^n}{dt^n} \exp \left( -t^2 \frac{S_m}{4V_m} \right)$$

which is more than necessary for proving other results and is an interesting estimate of  $G_{2m}^{(n)}(t)$  by itself.

**2. Some lemmas.** In the author's thesis [2] and in a paper in collaboration with A. Jakimovski [3; Lemma 2.1.] the following lemma was proved:

**LEMMA 2.1.** *Suppose  $\sum_{k=1}^{\infty} |a_k|^{-2} < \infty$  then the assumptions*

$$(2.1) \quad \sum_{k=m+1}^{\infty} |a_k|^{-(2+\alpha)} = o \left( \left( \sum_{k=m+1}^{\infty} |a_k|^{-2} \right)^{1+(\alpha/2)} \right) \quad m \rightarrow \infty$$

*for some fixed  $\alpha > 0$  and*

$$(2.2) \quad \lim_{m \rightarrow \infty} \left( \max_{k > m} |a_k|^{-2} \right) \left( \sum_{k=m+1}^{\infty} |a_k|^{-2} \right)^{-1} = 0$$

*are equivalent, and therefore the assumptions (2.1) for all positive  $\alpha$  are equivalent.*

*Proof.* Let us assume (2.1) for some  $\alpha > 0$ . If (2.3) is not valid then a subsequence  $\{m(r)\}$  of  $m + 1, m + 2, \dots$  exists such that for some  $\beta > 0$

$$\left( \max_{k \geq m(r)+1} |a_k|^{-2} \right) S_{m(r)}^{-1} \geq \beta > 0$$

for all  $r \geq 1$ . Therefore

$$\sum_{k=m(r)+1}^{\infty} |a_k|^{-2-\alpha} \geq \left( \max_{k \geq m(r)+1} |a_k|^{-2} \right)^{1+(\alpha/2)} \geq \beta^{1+(\alpha/2)} S_{m(r)}^{1+(\alpha/2)}$$

which contradicts (2.1).

Assuming (2.2) then

$$\begin{aligned} \sum_{k=m+1}^{\infty} |a_k|^{-2-\alpha} &= \sum_{k=m+1}^{\infty} |a_k|^{-2} \leq \left( \max_{k \geq m+1} |a_k|^{-\alpha} \right) S_m \\ &= \left( \left( \max_{k \geq m+1} |a_k|^{-2} \right) S_m^{-1} \right)^{\alpha/2} S_m^{1+(\alpha/2)} \\ &= o(S_m^{1+(\alpha/2)}) \quad (m \rightarrow \infty). \end{aligned}$$

The following two lemmas are easy to verify.

**LEMMA 2.2.** *If  $\{a_k\} \in \text{class } C$  then  $\{a_k\}$  satisfies assumption (2.2). If  $|\arg a_k| < \pi/4 - \eta$ ,  $0 < \operatorname{Re} a_i \leq \operatorname{Re} a_{i+1}$  and  $\{a_k\}$  satisfies assumption (2.2) then  $\{a_k\} \in \text{class } C$ .*

**LEMMA 2.3.** *If  $|\arg a_k| < (\pi/4) - \eta$  and  $\sum |a_k|^{-2} < \infty$ , then*

$$(2.3) \quad \cos\left(\frac{\pi}{2} - 2\eta\right) S_n \leq |V_n| \leq S_n.$$

We define now  $F_m(z)$  by

$$(2.4) \quad F_m(z) = E_m(z \cdot S_m^{-1/2}) = \sum_{k=m+1}^{\infty} \left( 1 - \frac{z^2}{a_k^2 S_m} \right).$$

**LEMMA 2.4.** *Suppose  $\{a_k\} \in \text{class } C$  then there exist constants  $k(p) > 0$  independent of  $m$  so that for all real  $y$*

$$(2.6) \quad |F_m(iy)| > 1 + k(p)y^{2p} \quad \text{for } m > m_0(p).$$

*Proof.* Define  $a_k = |a_k| e^{i\beta_k}$ ,  $-(\pi/4) + \eta < \beta_k < (\pi/4) - \eta$

$$\begin{aligned} |F_m(iy)| &= \left| \prod_{k=m+1}^{\infty} \left( 1 - \frac{(iy)^2}{a_k^2 S_m} \right) \right| \\ &\geq \prod_{k=m+1}^{\infty} \left( 1 + \frac{y^2}{|a_k|^2 S_m} \cos 2\beta_k \right) \\ &\geq \prod_{k=m+1}^{\infty} \left( 1 + \frac{y^2 \cos\left(\frac{\pi}{2} - 2\eta\right)}{|a_k|^2 S_m} \right) \\ &= 1 + \sum_{p=1}^{\infty} \frac{y^{2p} \cos^p\left(\frac{\pi}{2} - 2\eta\right)}{S_m^p p!} \sum_{\substack{k(i) > m \\ i \neq j \\ k(i) \neq k(j)}} |a_{k(1)} \cdots a_{k(p)}|^{-2}. \end{aligned}$$

Since we have  $\lim_{m \rightarrow \infty} \max_{k > m} |a_k|^{-2} S_m^{-1} = 0$  we can find  $m_0(p)$  so that for  $m > m_0(p)$   $\max_{k > m} |a_k|^{-2} < (1/2p) S_m$ . Therefore we have

$$\begin{aligned} & \sum_{\substack{k(i) > m \\ k(i) \neq k(j), i \neq j}} |a_{k(1)} \cdots a_{k(p)}|^{-2} \\ &= \sum_{\substack{k(i) > m \\ k(i) \neq k(j), i \neq j}} \left( S_m - \sum_{i=1}^{p-1} |a_{k(i)}|^{-2} \right) |a_{k(1)} \cdots a_{k(p-1)}|^{-2} \\ &\geq \frac{1}{2} S_m \sum_{\substack{k(i) > m \\ k(i) \neq k(j), j \neq i}} |a_{k(1)} \cdots a_{k(p-1)}|^{-2} \geq \left(\frac{1}{2}\right)^p S_m^p. \end{aligned}$$

Hence

$$|F_m(iy)| \geq 1 + y^{2p} \frac{\cos^p\left(\frac{\pi}{2} - 2\eta\right) S_m^p}{S_m^p p! 2^p} = 1 + k(p)y^{2p}.$$

3. The asymptotic estimates for  $G_{2m}^{(k)}(t)$ .

**THEOREM 3.1.** *Let  $\{a_k\} \in$  class  $C$ ; then for all  $n = 0, 1, \dots$*

$$(3.1) \quad \lim_{m \rightarrow \infty} S_m^{1/2} \frac{d^n}{dt^n} G_{2m}(S_m^{1/2}t) = \frac{1}{\sqrt{4\pi}} \left(\frac{S_m}{V_m}\right)^{1/2} \frac{d^n}{dt^n} \exp\left(-t^2 \frac{S_m}{4V_m}\right)$$

*uniformly in  $-\infty < t < \infty$  (we choose  $\arg V_m^{1/2} = (1/2) \arg V_m$ ).*

*Proof.* Following the proof of the special case  $n = 0$  and  $\arg a_k = 0$  by Hirschman—Widder [4; pp. 140-1] we have

$$S_m^{1/2} G_{2m}(S_m^{1/2}t) = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} \frac{e^{zt} dz}{F_{2m}(s)} = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{e^{iyt} dy}{F_{2m}(iy)}.$$

By an estimate of [5; p. 246] we have for  $|z| < R$  and

$$\begin{aligned} R \cdot [|a_k| S_m^{1/2}]^{-1} &\leq \frac{1}{2} \\ \left| \log \left\{ \left(1 - \frac{z^2}{a_k^2 S_m}\right) \exp(z^2/a_k^2 S_m) \right\} \right| &\leq 4R^3 \frac{1}{|a_k|^3 S_m^{3/2}}. \end{aligned}$$

Recalling that  $\sum_{k=m+1}^{\infty} 1/a_k^2 S_m = V_m/S_m$  and since by Lemma 2.1

$$\sum_{k=m+1}^{\infty} \frac{1}{|a_k|^3 S_m^{3/2}} = o(1) \quad m \rightarrow \infty,$$

we have for  $|z| < R$  and  $m > m_0(R)$

$$\left| F_m(z) - \exp\left(-\frac{V_m z^2}{S_m}\right) \right| < \varepsilon_1.$$

Since by Lemma 2.4  $R > R_0(\varepsilon_2, \eta)$  implies

$$\int_R^{\infty} \frac{|y|^n dy}{|F_{2m}(iy)|} < \varepsilon_2 \quad \text{and} \quad \int_{-\infty}^{-R} \frac{|y|^n dy}{|F_{2m}(iy)|} < \varepsilon_2$$

we have

$$\begin{aligned}
 & S_m^{1/2} \frac{d^n}{dt^n} G_{2m}(S_m^{1/2}t) \\
 &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{(iy)^n e^{iyt}}{F_m(iy)} dy \\
 &= \frac{1}{2\pi} \int_{-\infty}^{\infty} (iy)^n \exp\left(-\frac{V_m}{S_m} y^2 + iyt\right) dy + o(1) \\
 &= \frac{d^n}{dt^n} \left\{ \exp\left(-\frac{t^2 S_m}{4V_m}\right) \right. \\
 &\quad \times \left. \frac{1}{2\pi} \int_{-\infty}^{\infty} \exp[-(V_m^{1/2} S_m^{-1/2} y - it S_m^{1/2} (4V_m)^{-1/2})^2] dy \right\} + o(1) \\
 &= \left(\frac{S_m}{V_m}\right)^{1/2} \frac{d^n}{dt^n} \left\{ \exp\left(-\frac{t^2 S_m}{4V_m}\right) \frac{1}{2\pi} \int_r e^{-z^2} dz \right\} + o(1) \\
 &= \frac{1}{\sqrt{4\pi}} \left(\frac{S_m}{V_m}\right)^{1/2} \frac{d^n}{dt^n} \exp\left(\frac{-t^2 S_m}{4V_m}\right) + o(1) \quad (m \uparrow \infty)
 \end{aligned}$$

using the residue theorem, the fact that  $e^{-z^2}$  is entire and that

$$|\arg V_m^{1/2}| < \frac{\pi}{4} \quad \text{for all } m .$$

As a corollary we derive

**THEOREM 3.2.** *If  $\{a_k\}$  satisfies assumption C then*

$$(3.2) \quad \int_{-\infty}^{\infty} |G_{2m}(t)| dt = (\cos^2 \varphi_m - \sin^2 \varphi_m)^{-1/2} + o(1) \quad m \rightarrow \infty$$

and

$$(3.3) \quad \int_{-\infty}^{\infty} |tG'_{2m}(t)| dt = (\cos^2 \varphi_m - \sin^2 \varphi_m)^{-3/2} + o(1) \quad m \rightarrow \infty$$

where  $2\varphi_m = \arg V_m$ .

*Proof.* Since by Lemma 2.4 of [1].

$$(3.4) \quad |G_{2m}(t)| < MS_m^{-1/2} \exp(-KS_m^{-1/2} |t|)$$

we have

$$\begin{aligned}
 \int_{-\infty}^{\infty} |G_{2m}(t)| dt &= \int_{-\infty}^{\infty} |S_m^{1/2} G_{2m}(S_m^{1/2}t)| dt = \int_{-R}^R |S_m^{1/2} G_{2m}(S_m^{1/2}t)| dt \\
 &+ o(1) \quad (R \uparrow \infty) .
 \end{aligned}$$

This combined with (3.1) and a simple integration yield (3.2).

To prove (3.3) we use Lemma 3.2 case A (since for  $\{a_k\} \in$  class C  $S_m \geq 4r_{m+1}^{-2} \equiv 4|a_{m+1}|^{-2}$  for  $m > m_0$ ) which is

$$(3.5) \quad |G'_{2m}(t)| \leq M_1 S_m^{-1} \exp(-K_1 S_m^{-1/2} |t|).$$

Therefore we have

$$\int_R^\infty |S_m t G'_{2m}(S_m^{1/2} |t|)| dt \leq M_1 \frac{1}{(K_1)^2} e^{-K_1 R} = o(1) \quad R \rightarrow \infty.$$

This implies

$$\begin{aligned} \int_{-\infty}^\infty |t G'_{2m}(t)| dt &= \int_{-\infty}^\infty |S_m t G'_{2m}(S_m^{1/2} t)| dt \\ &= \frac{1}{2\sqrt{4\pi}} \left| \frac{S_m}{V_m} \right|^{3/2} \int_{-\infty}^\infty t^2 \exp\left(-\frac{t^2}{4} S_m \operatorname{Re} \frac{1}{V_m}\right) dt + o(1) \\ &= \left( \frac{|V_m|^{-1}}{\operatorname{Re}(V_m^{-1})} \right)^{3/2} + o(1) = (\cos^2 \varphi_m - \sin^2 \varphi_m)^{-3/2} + o(1) \quad (m \uparrow \infty). \end{aligned}$$

4. **Remarks.** I. For the theorems and the lemmas proved in this paper  $0 < \operatorname{Re} a_i \leq \operatorname{Re} a_{i+1}$  is not essential and the condition (2.2) can replace it and (1.3).

II. Theorem 3.1 which replaces Theorem 4.1 yields for the case  $n = 0$  only the following

$$(4.6) \quad G_{2m}(t) = (4\pi V_m)^{-1/2} \exp(-t^2/4V_m) + o(S_m^{-1/2}) \quad m \rightarrow \infty$$

but if one follows the proof of Theorem 4.1 of [1] and Lemma 4.2 of [1] almost literally one obtains for  $\{a_k\} \in \text{class } C$

$$(4.7) \quad G_{2m}(t) = (4\pi V_m)^{-1/2} \exp(-t^2/4V_m) + o(|a_{m+1}|^{-2} S_m^{-3/2}) \quad m \rightarrow \infty$$

which is somewhat more general.

### REFERENCES

1. J. Dauns and D. V. Widder, *Convolution transforms whose inversion function have complex roots*, Pacific J. Math. (2) **15** (1965).
2. Z. Ditzian, *Inversion formulae and various properties of integral transforms* (Thesis, 1965, Hebrew University).
3. Z. Ditzian and A. Jakimovski, *Inversion and Jump formulae for convolution transform* (to appear).
4. I. I. Hirschman and D. V. Widder, *The Convolution Transform*, Princeton Press, 1955.
5. E. C. Titchmarsh, *The Theory of Functions*, Oxford University Press, 1932.

Received May 23, 1966.

WASHINGTON UNIVERSITY  
 ST. LOUIS, MISSOURI AND  
 MICHIGAN STATE UNIVERSITY  
 EAST LANSING, MICHIGAN

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON  
Stanford University  
Stanford, California

J. P. JANS  
University of Washington  
Seattle, Washington 98105

J. DUGUNDJI  
University of Southern California  
Los Angeles, California 90007

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL ORDNANCE TEST STATION

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

# Pacific Journal of Mathematics

Vol. 21, No. 2

December, 1967

Arne P. Baartz, <i>The measure algebra of a locally compact semigroup</i> . . . . .	199
Robert F. Brown, <i>On maps with identical fixed point sets</i> . . . . .	215
C. Buttin, <i>Existence of a homotopy operator for Spencer's sequence in the analytic case</i> . . . . .	219
Henry Werner Davis, <i>An elementary proof that Haar measurable almost periodic functions are continuous</i> . . . . .	241
Zeev Ditzian, <i>On asymptotic estimates for kernels of convolution transforms</i> . . . . .	249
Robert E. Edwards, <i>Boundedness principles and Fourier theory</i> . . . . .	255
John A. Hildebrant, <i>On compact unithetic semigroups</i> . . . . .	265
Marinus A. Kaashoek and David Clark Lay, <i>On operators whose Fredholm set is the complex plane</i> . . . . .	275
Sadao Kató, <i>Canonical domains in several complex variables</i> . . . . .	279
David Clifford Kay, <i>The ptolemaic inequality in Hilbert geometries</i> . . . . .	293
Joseph D. E. Konhauser, <i>Biorthogonal polynomials suggested by the Laguerre polynomials</i> . . . . .	303
Kevin Mor McCrimmon, <i>Macdonald's theorem with inverses</i> . . . . .	315
Harry Eldon Pickett, <i>Homomorphisms and subalgebras of multialgebras</i> . . . . .	327
Richard Dennis Sinkhorn and Paul Joseph Knopp, <i>Concerning nonnegative matrices and doubly stochastic matrices</i> . . . . .	343
Erling Stormer, <i>On anti-automorphisms of von Neumann algebras</i> . . . . .	349
Miyuki Yamada, <i>Regular semi-groups whose idempotents satisfy permutation identities</i> . . . . .	371