

Pacific Journal of Mathematics

THE EXTENSION OF BILINEAR FUNCTIONALS

THOMAS LEE HAYDEN

THE EXTENSION OF BILINEAR FUNCTIONALS

T. L. HAYDEN

Using the relationship between bilinear functionals and linear operators we obtain some theorems on the extension of bilinear functionals. To extend bilinear functionals in Hilbert Spaces a special constructional process is given which is a generalization of the usual inner product. This allows the construction of bilinear functionals with special properties. In particular it allows a generalization of the Lax-Milgram Theorem. We also extend the Lax-Milgram Theorem to reflexive Banach Spaces.

In order to fix the terminology, let U and V be Banach Spaces (real or complex), then by a bilinear functional we mean a function F from $U \times V$ to the complex (or real) numbers such that $F(u, v)$ is linear on U for each fixed $v \in V$ and vice versa. The norm of a bounded bilinear functional F , denoted by $\|F\|$, is defined as:

$$\|F\| = \inf \{K > 0: |F(u, v)| \leq K\|u\|\|v\| \text{ for all } u \in U, v \in V\}.$$

Hence a bounded bilinear functional is jointly continuous on $U \times V$ in the product topology and we note that:

$$\|F\| = \sup_{\substack{\|x\| \leq 1 \\ \|y\| \leq 1}} |F(x, y)| = \sup_{\substack{\|x\|=1 \\ \|y\|=1}} |F(x, y)| \leq \sup_{\|x\| \|y\|=1} |F(x, y)| \leq \|F\|.$$

If S and T are subspaces of U and V respectively and B_0 is a bounded bilinear functional on $S \times T$, then we call B an extension of B_0 to $U \times V$ if B is a bounded bilinear functional on $U \times V$ such that $B_0(s, t) = B(s, t)$ on $S \times T$ and $\|B_0\| = \|B\|$. If such an extension exists we shall say that B_0 can be extended to $U \times V$. Furthermore, if each bounded bilinear functional on $S \times T$ can be extended to $U \times V$ we will say that S and T have the bilinear extension property.

2. Some extension theorems.

THEOREM 1. *Suppose U, V, W are normed linear spaces and S and T are subsets of U and V respectively. A necessary and sufficient condition to extend a bounded bilinear operator B_0 from $S \times T$ into W to a bounded bilinear operator B from $(\text{Span } S) \times (\text{Span } T)$ into W is that there exists a constant c such that*

$$\left\| \sum_{k,j} \alpha_k \beta_j B_0(x_k, y_j) \right\| \leq c \left\| \sum_k \alpha_k x_k \right\| \left\| \sum_j \beta_j y_j \right\|.$$

For each finite subset $x_1, \dots, x_n \in S$ and $y_1, \dots, y_m \in T$ and every choice of scalars α_k and β_j .

The proof is similar to the proof of the linear case which is found for example in [3, p. 127]. That one can extend bounded bilinear functionals to the closure or completion of a space has a proof similar to the linear case. For one such version for bilinear functionals see [1, p. 105]. Unless it is specifically mentioned we will not assume that the subspaces in question in each theorem are necessarily closed. Hence in each such theorem one should first extend to the closure and then proceed with the proof.

Our next two theorems show the relation of extending bilinear functionals and the extension of linear operators. One of the best references to the problem of extending linear operators is Nachbin [6]. (See also [5]). References and additional information to the following facts may be found in [6].

We shall say that a Banach Space U has property E (the extension property of type ∞ in the terminology of [6]) if for any Banach Space V , any closed subspace S of V , and any continuous linear operator F_0 from S into U , there exists a continuous linear extension F of F_0 to V with values in U and the same norm. The classical Hahn-Banach theorem says U has property E if its dimension is one. A Banach Space U has property E if and only if for any Banach Space W containing U as a Banach subspace there is a projection of norm one of W onto U .

Also the topological dual U^* of a Banach Space U has property E if and only if U is metrically isomorphic to a space $L^1(\mu)$ of all real integrable functions with respect to a suitable positive measure μ on a locally compact space [6, p. 345].

THEOREM 2. *If U and V are Banach Spaces and S and T are subspaces of U and V respectively such that T^* (the topological dual of T) and U^* have property E , then each bounded bilinear functional on $S \times T$ can be extended to $U \times V$.*

Proof. Let B_0 be a bounded bilinear functional on $S \times T$ with norm $\|B_0\|$. If we fix $s \in S$, then $B(s, t) = F_s(t)$ is a bounded linear functional on T , i.e. $F_s(t) \in T^*$. Let us denote the value of a linear functional F at x by $\langle x, F \rangle$. If G_0 is the map from S into T^* such that $G_0(s) = F_s$, then G_0 is a bounded linear operator from S into T^* and $\|G_0\| = \|B_0\|$. The linearity of G_0 follows from the relation $\langle t, G_0(s) \rangle = B_0(s, t)$ and the bilinearity of B , and

$$\|G_0\| = \sup_{\|s\| \leq 1} \|G_0(s)\| = \sup_{\substack{\|t\| \leq 1 \\ \|s\| \leq 1}} |\langle t, G_0(s) \rangle| = \sup_{\substack{\|t\| \leq 1 \\ \|s\| \leq 1}} |B_0(s, t)| = \|B_0\| .$$

Since T^* has property E we extend G_0 to a linear operator G on U into T^* with the same norm. Now let $B(u, t) = \langle t, G(u) \rangle$ for $(u, t) \in U \times T$ and it is easy to see that B is an extension of B_0 to $U \times T$. Using a similar procedure for the spaces U^* and T we obtain the desired extension to $U \times V$.

One natural way to try to extend bilinear functionals is to factor through the bilinear functional on $S \times T$ into a linear functional on the tensor product $S \otimes T$. Now the bounded bilinear functionals on $S \times T$ are isometric to the bounded linear functionals on $S \otimes T$ under the greatest cross norm topology on $S \otimes T$. (See Grothendieck, *Produits Tensoriels Topologiques Et Espaces Nucleaires*, Memoirs of A.M.S. No. 16, for related statements in topological vector spaces.) Hence if we extend the linear functional on $S \otimes T$ to $U \otimes V$ we have an extension of the bilinear functional on $S \times T$ to $U \times V$. However, Schatten [7, p. 57] has shown that the greatest cross norm topology on $U \otimes V$ is not an extension of the greatest cross norm topology on $S \otimes T$ in general. The next few theorems are closely related to those obtained by Schatten in proving this result.

THEOREM 3. *Suppose U and V are Banach Spaces with the bilinear extension property on the subspaces S and T , then every bounded linear operator from S into T^* (T into S^*) can be extended to a linear operator from U into T^* (V into T^*) with the same norm.*

Proof. Suppose F_0 is a bounded linear operator from S into T^* . Then there exists, as in the proof of Theorem 2, a bounded bilinear functional B_0 on $S \times T$ such that $B_0(s, t) = \langle t, F_0(s) \rangle$ and $\|B_0\| = \|F_0\|$. Since S and T have the extension property we may extend B_0 to a bounded bilinear functional B on $U \times V$, such that $\|B\| = \|B_0\|$. We may restrict B to $U \times T$ and find an F such that $B(u, t) = \langle t, F(u) \rangle$ on $U \times T$ with $\|F\| = \|F_0\|$ and F restricted to S is F_0 . Hence F is the desired extension of F_0 .

We also note that we could consider the set $U \times V$ and by a similar process we have an extension of F_0 to U but with final values in V^* .

THEOREM 4. *Suppose in addition to the hypothesis of Theorem 3 that $T^* = S$, then there is a projection of norm one of U onto S .*

Proof. Let F_0 be the identity operator from S into T^* . As in the proof of Theorem 3 associate B_0 with F_0 and $\|B_0\| = 1$. The extension then of F_0 to F gives us a linear operator of norm one of U

into T^* or S which is the identity on S , i.e., F is a projection of norm one.

THEOREM 5. *Let S be a subspace of the Banach Space U , then the existence of a projection of U on S of norm one implies that if V is a Banach Space then every bounded bilinear form on $S \times V$ can be extended to $U \times V$.*

Proof. Let B_0 be a bounded linear operator on $S \times V$, and let P be a projection of U onto S with norm one. It is clear that $B(u, v) = B_0(Pu, v)$ is the desired extension.

THEOREM 6. *Suppose S is a subspace of the Banach Space U , and that S is the topological dual of a Banach Space T , i.e. $S = T^*$, and that every bounded bilinear functional on $S \times T$ can be extended to $U \times T$, then if V is a Banach Space every bilinear functional on $S \times V$ can be extended to $U \times V$.*

Proof. By Theorem 4 there is a projection of norm one of U onto S and hence by Theorem 5 the result follows.

COROLLARY 1. *If S is a subspace of Hilbert Space U then for every Banach Space V and every bounded bilinear functional B on $S \times V$ there is an extension to $U \times V$.*

COROLLARY 2. *If U and V are Banach Spaces and S and T are subspaces of U and V respectively and there exists a projection of norm one of U on S and a projection of norm one of V on T then every bounded bilinear functional on $S \times T$ can be extended to $U \times V$.*

COROLLARY 3. *A Banach Space U is a Hilbert Space if and only if every bounded bilinear functional on $S \times S^*$ can be extended to $U \times S^*$ for any two dimensional subspace S of U .*

Proof. By Theorem 4 we have a projection of norm one on each two dimensional subspace, which implies that U is a Hilbert Space.

Although Corollary 2 shows that bilinear functionals on subspaces of Hilbert Spaces may always be extended, we repeat another proof given in [2] whose explicit construction yields other results immediately.

THEOREM 7. *Suppose S_1, S_2, \dots, S_n are subspaces respectively of the complex Hilbert Spaces X_1, X_2, \dots, X_n and $f(s_1, s_2, \dots, s_n)$ is a*

multilinear functional defined on $\times_{k=1}^n S_k$ with norm K . Then there exists a multilinear functional F on $\times_{k=1}^n X_k$ with norm K such that F is f on $\times_{k=1}^n S_k$.

Proof. First we show that f can be extended to a function defined on $\times_{k=1}^n (S_k \cup \{x_k\})$ for $x_k \in X_k, k = 1, 2, \dots, n$. In fact, let $x_k \perp S_k$, and define

$$h(a_1x_1 + s_1, \dots, a_nx_n + s_n) = K \prod_{k=1}^n a_k \|x_k\| + f(s_1, \dots, s_n)$$

where a_1, a_2, \dots, a_n are scalars and K is the norm of f . It is immediate that the extension is multilinear, and we now show that the norm is the same.

First we show by induction that

$$\left(\prod_{k=1}^n \|a_kx_k\| + \prod_{k=1}^n \|s_k\| \right) \leq \prod_{k=1}^n (\|a_kx_k\|^2 + \|s_k\|^2)^{1/2} \quad \text{for } n \geq 2.$$

For $n = 2$ this is Cauchy's inequality. Suppose it is true for $n = l - 1$, then

$$\begin{aligned} \left(\prod_{k=1}^l \|a_kx_k\| + \prod_{k=1}^l \|s_k\| \right) &\leq \left(\prod_{k=1}^{l-1} \|a_kx_k\| + \prod_{k=1}^{l-1} \|s_k\| \right) (\|a_lx_l\|^2 + \|s_l\|^2)^{1/2} \\ &\leq \prod_{k=1}^{l-1} (\|a_kx_k\|^2 + \|s_k\|^2)^{1/2} (\|a_lx_l\|^2 + \|s_l\|^2)^{1/2} \\ &\leq \prod_{k=1}^l (\|a_kx_k\|^2 + \|s_k\|^2)^{1/2}. \end{aligned}$$

Hence

$$\begin{aligned} |h(a_1x_1 + s_1, \dots, a_nx_n + s_n)| &\leq K \left(\prod_{k=1}^n \|a_kx_k\| + \prod_{k=1}^n \|s_k\| \right) \\ &\leq K \prod_{k=1}^n (\|a_kx_k\|^2 + \|s_k\|^2)^{1/2} = K \prod_{k=1}^n \|a_kx_k + s_k\|. \end{aligned}$$

The remainder of the proof follows from a straightforward application of Zorn's Lemma similar to that in the Hahn-Banach Theorem.

3. Corollaries. We will suppose in this section that U and V are Hilbert Spaces and S and T are subspaces of U and V respectively. From the proof of Theorem 7 we obtain the following

COROLLARY 4. *Suppose $S \subset U$ and B is a bilinear functional on $S \times S$ such that:*

- (i) $|B(s, t)| \leq K_1 \|s\| \|t\|$ for each s, t in S and
- (ii) $|B(s, s)| \geq K_2 \|s\|^2$ for each s in S ,

Then B may be extended to $U \times U$ such that (i) and (ii) hold on $U \times U$.

One also can easily see that a bounded symmetric bilinear functional on $S \times S$ ($B(u, v) = B(v, u)$) can be extended to a symmetric bilinear functional on $U \times U$. Furthermore a bounded sesquilinear functional on $S \times T$ can be extended to $U \times V$. One simply uses the fact that the sesquilinear form on $S \times T$ is a bilinear form on $S \times T'$ where T' is the so called complex conjugate of T .

The proofs of the following corollaries follow from the extension theorem and since they are similar to the proofs in the linear case, they will be omitted.

COROLLARY 5. *If $u \neq 0$ and $v \neq 0$ are in U and V respectively then there exists a bilinear functional B on $U \times V$ such that $B(u, v) = \|u\| \|v\|$ and $\|B\| = 1$.*

COROLLARY 6. *Suppose $x_0 \in U$ and distance from x_0 to S is $d_1 > 0$ and $y_0 \in V$ such that the distance from y_0 to T is $d_2 > 0$. Then there exists a bilinear functional B of norm one on $U \times V$ such that $B(s, t) = 0$ for $(s, t) \in S \times T$ and $B(x_0, y_0) = d_1 d_2$.*

COROLLARY 7. *Let $E \subset U, F \subset V$. A necessary and sufficient condition that $(x_0, y_0) \in U \times V$ to belong to $(\text{closure span } E) \times (\text{closure span } F)$ is that $B(x_0, y_0) = 0$ for every bounded linear functional which vanishes on $E \times F$.*

The next corollary follows from Theorem 1 and the extension theorem and indicates when one can solve infinite systems of equations in bilinear functionals.

COROLLARY 8. *Let M and N be indexing sets. Let $\{(u_\alpha, v_\beta); (\alpha, \beta) \in M \times N\} \subset U \times V$ and $\{C_{\alpha\beta}; (\alpha, \beta) \in M \times N\} \subset \text{complex numbers}$. A necessary and sufficient condition for the existence of a bounded bilinear functional B on $U \times V$ such that*

- (i) $B(u_\alpha, v_\beta) = C_{\alpha\beta}$ for each $(\alpha, \beta) \in M \times N$ and
- (ii) $\|B\| \leq K$

is that

$$|\sum d_\alpha e_\beta C_{\alpha\beta}| \leq K \|\sum d_\alpha u_\alpha\| \|\sum e_\beta v_\beta\|$$

hold for every finite subset of $M \times N$ and every choice of scalars d_α and e_β .

We note that if F and G are bounded linear functionals on S and T respectively then the bilinear functional defined by $B(s, t) = F(s)G(t)$ is bounded on $S \times T$ and can obviously be extended. On the other hand a nondegenerate bilinear functional on $U \times V$ can not be re-

presented in a form with the variables separated. In fact we have the following

THEOREM 8. *Suppose X and Y are normed linear spaces and B is a bounded bilinear functional on $X \times Y$ such that $B(x, y) = 0$ for all $y \in Y$ implies that x is zero. Then B can not be factored as $B(x, y) = \sum_{i=1}^n F_i(x)G_i(y)$ where $F_i \in X^*$, $G_i \in Y^*$ for $1 \leq i \leq n$, and $n < \dim X$.*

Proof. Suppose the conclusion is false. Let $K_i = \text{Kernal } F_i$, $1 \leq i \leq n$. Since K_i is a maximal closed subspace of X , hence $\dim \bigcap_{i=1}^n K_i \neq 0$. But for $x \in \bigcap_{i=1}^n K_i$, $B(x, y) = 0$ for all $y \in Y$.

It is known that if there exists a nondegenerate bilinear functional on $U \times V$ where U and V are finite dimensional then $\dim U = \dim V$.

THEOREM 9. *Suppose X and Y are normed linear spaces and B is a nondegenerate bounded bilinear functional on $X \times Y$. Then $\dim X \leq \dim Y^*$ and $\dim Y \leq \dim X^*$.*

Proof. Again as in the proof of Theorem 3 we associate a linear operator A from X into Y^* with the bilinear functional B . Since B is nondegenerate this implies that A is one to one and the result follows.

COROLLARY 9. *If U and V are Hilbert Spaces and there exists a nondegenerate bounded bilinear functional on $U \times V$ then $\dim U = \dim V$ where $\dim U$ and $\dim V$ may be either Hilbert Space dimension or Vector Space dimension.*

4. An application. A variant of the F. Riesz representation, the Lax-Milgram Theorem [4], is useful in proving the existence of solutions of partial differential equations. Let (\cdot, \cdot) denote the inner product in Hilbert Space.

THEOREM 10. (*Lax-Milgram*) *Let U be a Hilbert Space and B a bounded sesqui-linear functional on $U \times U$ such that there exists a $\delta > 0$ such that $B(u, u) \geq \delta \|u\|^2$. Then there exists a unique bounded linear operator T with a bounded inverse T^{-1} such that $(u, v) = B(u, Tv)$ when $u, v \in U$ and $\|T\| \leq 1/\delta$, $\|T^{-1}\| \leq \|B\|$.*

LEMMA. *If S is a subspace of a Hilbert Space U and B is a bounded sesquilinear functional on $S \times S$ such that $B(s, s) \geq \delta \|s\|^2$*

for some $\delta > 0$, then B can be extended to $U \times U$ with the same properties.

Proof. As in the proof of Theorem 5, let $x \perp S$ and define \bar{B} on $(x \cup S) \times (x \cup S)$ by

$$\bar{B}(\alpha x + s, \beta x + t) = \alpha\bar{\beta} \|B\| \|x\|^2 + B(s, t).$$

As before we now observe that \bar{B} is a sesqui-linear functional on $(x \cup S) \times (x \cup S)$ with norm $\|B\|$.

Now

$$\begin{aligned} \bar{B}(\alpha x + s, \alpha x + s) &= \|B\| |\alpha|^2 \|x\|^2 + B(s, s) \\ &\geq \delta(\|\alpha x\|^2 + \|s\|^2) \\ &= \delta\|\alpha x + s\|^2. \end{aligned}$$

An application of Zorn's Lemma completes the proof. We note that no claim to the uniqueness of the extension is made.

This Lemma immediately gives us the following extension of the Lax-Milgram Theorem.

THEOREM 11. *Let S be a subspace of the Hilbert Space U and B a sesquilinear functional satisfying the hypothesis of the Lax-Milgram Theorem. Then there is an extension of B to \bar{B} on $U \times U$ satisfying the hypothesis of Lax-Milgram Theorem and hence there is an extension of the linear operator T to \bar{T} from U onto U which satisfies the conclusion of the Lax-Milgram Theorem.*

We may also extend the Lax-Milgram Theorem in the following direction.

THEOREM 12. *Suppose U and V are Banach Spaces, V is reflexive, and that B is a bounded nondegenerate bilinear functional on $U \times V$. Then a necessary and sufficient condition that every bounded linear functional F on V have a unique representation of the form $F(v) = B(u, v)$ for some fixed $u \in U$ is that there exists an $m > 0$ such that for each $u \in U$, $\sup_{\|v\|=1} |B(u, v)| \geq m\|u\|$.*

Proof. Suppose the representation holds, then B induces a linear operator A from U into V^* in the usual manner. Since B is nondegenerate, A is one-to-one and $\|A\| = \|B\|$. Since every linear functional in V^* has this representation A must be onto. Hence A has a bounded inverse A^{-1} . However, a necessary and sufficient condition that A have a bounded inverse is that there exists an $m > 0$ such that $\|A(u)\| \geq m\|u\|$. But

$$\|A(u)\| = \sup_{\|v\|=1} |\langle v, A(u) \rangle| = \sup_{\|v\|=1} |B(u, v)|,$$

and it follows that the condition is necessary.

Now suppose that the condition is satisfied. Hence A is one-to-one and onto $R(A)$ which is closed in V^* . If $R(A) \neq V^*$ let $f^* \neq 0 \in R(A)$. By the Hahn Banach Theorem there exists $v^{**} \in V^{**}$ such that $v^{**}(f^*) = 1$ and $v^{**}(R(A)) = 0$. Let $v \in V$ correspond to v^{**} (V is reflexive). Hence $f^*(v) = 1$, and $B(u, v) = A(u)(v) = 0$ for all u which implies that $v = 0$ since B is nondegenerate. This contradiction shows $R(A) = V^*$.

REMARKS. Note that $\|A^{-1}\| \leq 1/m$. Also if there exists an $m_1 > 0$ such that for each $\|v\| = 1$, $\sup_{\|u\|=1} |B(u, v)| \geq m_1$, then this implies that the bilinear form is nondegenerate and hence the usual Lax-Milgram Lemma is a special case.

There are some corollaries to the Lax-Milgram Theorem due to Littman and Schechter [8, 568] which are useful in Elliptic Boundary Value Problems. The above theorem gives the following generalizations.

COROLLARY. *Suppose B is a bounded, nondegenerate, bilinear functional on the Banach Spaces U and V , and that V is reflexive. If there exists an $m > 0$ such that for each $u \in U$, $\sup_{\|v\|=1} |B(u, v)| > m\|u\|$, then for each proper closed subspace K of V there exists a $u \in U$ such that $B(u, v) = 0$ for each v in K .*

COROLLARY. *Suppose U and V are Banach Spaces, V is reflexive and that B is a bounded bilinear functional on $U \times V$. If there exist closed subspaces S and T of U and V respectively such that B satisfies the hypothesis of the preceding corollary on $S \times T$, then for each $u \in U$, there is a unique u^* in S such that $B(u - u^*, T) = 0$.*

Proof. For a fixed u , $B(u, t)$ is a bounded linear functional t^* on T . Therefore there is a unique u^* in S such that $B(u^*, t) = \langle t, t^* \rangle$ for all t in T . Hence $B(u, t) = B(u^*, t)$ for all t in T .

COROLLARY. *Suppose the hypothesis of the preceding corollary holds, then for each $u \in U$ there is a unique u^* in S and a u^{**} such that $B(u^{**}, t) = 0$ for all $t \in T$ and $u = u^* + u^{**}$.*

The author appreciates the helpful suggestions of the referee and in particular wishes to give credit to the referee for the brief proof of Theorem 12.

REFERENCES

1. R. C. Cooke, *Linear Operators*, MacMillan Co., London, 1953.
2. T. L. Hayden, *A conjecture on the extension of bilinear functionals* (to appear in the Amer. Math. Monthly)
3. L. V. Kantorovich and G. P. Akilov, *Functional Analysis in Normed Spaces*, MacMillan Co., New York 1964.
4. P. D. Lax and A. N. Milgram. *Parabolic equations*, Ann. of Math. **33** (1954), 167-190.
5. J. Lindenstrauss, *On the extension of operators*, Illinois J. Math. **8** (1964), 488-499.
6. L. Nachbin, *Some problems in extending and lifting linear transformations*, Proc. international symposium on linear spaces, Jerusalem, 1961, 340-352.
7. R. Schatten, *A Theory of Cross Spaces*, Annals of Math. Studies, No. 26, Princeton, 1950.
8. M. Schechter, *Remarks on elliptic boundary value problems*, Comm. Pure Appl. Math. **12** (1959), 561-578.

Received August 3, 1966.

UNIVERSITY OF KENTUCKY
LEXINGTON, KENTUCKY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California

J. P. JANS

University of Washington
Seattle, Washington 98105

J. DUGUNDJI

University of Southern California
Los Angeles, California 90007

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Charles A. Akemann, <i>Some mapping properties of the group algebras of a compact group</i>	1
A. V. Boyd, <i>Note on a paper by Uppuluri</i>	9
Thomas Craig Brown, <i>A semigroup union of disjoint locally finite subsemigroups which is not locally finite</i>	11
Richard Thomas Bumby and Everett C. Dade, <i>Remark on a problem of Niven and Zuckerman</i>	15
James Calvert, <i>An integral inequality with applications to the Dirichlet problem</i>	19
Jack Gary Ceder and Terrance Laverne Pearson, <i>On products of maximally resolvable spaces</i>	31
William Guignard Faris, <i>The product formula for semigroups defined by Friedrichs extensions</i>	47
Robert S. Freeman, <i>Closed operators and their adjoints associated with elliptic differential operators</i>	71
Thomas Lee Hayden, <i>The extension of bilinear functionals</i>	99
Gloria Conyers Hewitt, <i>Limits in certain classes of abstract algebras</i>	109
Tilla Weinstein, <i>The dilatation of some standard mappings</i>	117
Mitsuru Nakai, <i>On Evans' kernel</i>	125
Ernest Levane Roetman, <i>On the biharmonic wave equation</i>	139
Malcolm Jay Sherman, <i>Operators and inner functions</i>	159
Walter Laws Smith, <i>On the weak law of large numbers and the generalized elementary renewal theorem</i>	171
A. J. Ward, <i>On H-equivalence of uniformities: The Isbell-Smith problem</i>	189