

Pacific Journal of Mathematics

***AW**-ALGEBRAS ARE *QW**-ALGEBRAS**

BARRY E. JOHNSON

A W^* -ALGEBRAS ARE QW^* -ALGEBRAS

B. E. JOHNSON

G. A. Reid has introduced a class of B^* -algebras called QW^* -algebras which includes the W^* -algebras and which is included in the class of AW^* -algebras. In this paper it is shown that the QW^* -algebras are exactly the AW^* -algebras.

We shall use Reid's notation (see [2]) without further explanation.

THEOREM. *Let A be an AW^* -algebra. Then A is a QW^* -algebra.*

Proof. Let B be a norm-closed $*$ -subalgebra of A . Then, using [1; Theorem 2.3] we see that A contains a hermitian idempotent P such that PA is the right annihilator of B . Since B is a $*$ -subalgebra we see that the left annihilator of B is $(PA)^* = AP$. Thus $B_0 = AP \cap PA = PAP$ and $B_{00} = (I - P)A(I - P)$. It follows by [1; Theorem 2.4] that $B_{00} \supset B$ is an AW^* -algebra with identity $I - P$.

Using the Gelfand-Naimark Theorem [3; 244] we can consider B_{00} as an algebra of operators on a hilbert space H where the identity in B_{00} corresponds to the identity operator I_H in H . Let $(\mathcal{I}, \mathcal{S})$ be a double centraliser on B and let T be the element of $\mathcal{B}(H)$ corresponding to $(\mathcal{I}, \mathcal{S})$ under the isomorphism in [2; Proposition 3]. We have $TB \subset B, BT \subset B$ and wish to show that there is an element S of B_{00} with $SL = TL$ and $LS = LT$ for all $L \in B$.

We may clearly suppose T to be symmetric since the general case follows by considering separately the real and imaginary parts of T . Let K be the closed linear subspace of H generated by BH and P_K the orthogonal projection onto K . Let $\{F_\lambda\}$ be the spectral family of T [4; p. 275] and put $E_\lambda = P_K F_\lambda = F_\lambda P_K$. $\{E_\lambda\}$ is essentially the spectral family of T considered as an operator in K . Define

$$C_\lambda = \{P_K f(T); f \in C(\sigma(T)), f(\lambda') = 0 \text{ for } \lambda' \leq \lambda\}$$

$$D_\lambda = \{P_K f(T); f \in C(\sigma(T)), f(\lambda') = 0 \text{ for } \lambda' \geq \lambda\}$$

where $C(\sigma(T))$ is the set of continuous complex valued functions on $\sigma(T)$. The elements of C_λ, D_λ are essentially functions of T in $\mathcal{B}(K)$. Since the elements of C_λ and D_λ are limits in the uniform operator topology of sequences of polynomials in T we see that $C_\lambda B, D_\lambda B, BC_\lambda$ and BD_λ are subsets of B and hence of B_{00} . Using Kaplansky's result [1; Theorem 2.3] we can find an orthogonal projection $P_\lambda \in B_{00}$ such that $P_\lambda B_{00}$ is the right annihilator of BC_λ in B_{00} . Since B_{00} contains I_H we see $P_\lambda \in P_\lambda B_{00}$ and so $BC_\lambda P_\lambda = \{0\}$ and $P_\lambda C_\lambda B = \{0\}$. Thus for $\xi \in BH$, and hence for $\xi \in K, P_\lambda C_\lambda \xi = \{0\}$. However for $\xi \in H \ominus K, C_\lambda \xi = \{0\}$ and

so $P_\lambda C_\lambda \xi = \{0\}$ for all $\xi \in H$, that is $P_\lambda C_\lambda = \{0\}$. $P_K - E_{\lambda+0}$ is a strong operator limit of elements of C_λ [4; p. 263] so that $P_\lambda(P_K - E_{\lambda+0}) = 0$. Thus, with the usual ordering of projections, $I_H - P_\lambda \geq P_K - E_{\lambda+0}$.

We have $C_\lambda D_\lambda = \{0\}$ so that $D_\lambda B \subset P_\lambda B_{00}$ and $D_\lambda BH \subset P_\lambda B_{00}H = P_\lambda H$. Thus $D_\lambda K \subset P_\lambda H$ and, since $D_\lambda(H \ominus K) = \{0\}$ we have $D_\lambda H \subset P_\lambda H$. Again taking strong operator limits we obtain $E_{\lambda-0}H \subset P_\lambda H$ and so $P_\lambda \geq E_{\lambda-0}$.

If $\lambda \leq \mu$ we have $C_\lambda \supset C_\mu$ and so $P_\lambda B_{00} \subset P_\mu B_{00}$. This implies that $P_\lambda H \subset P_\mu H$ and so $P_\lambda \leq P_\mu$. If $\lambda < -\|T\|$ then $P_K \in C_\lambda$, $BC_\lambda = B$ and, since the annihilator of B in B_{00} is $\{0\}$, we have $P_\lambda = 0$. Similarly for $\lambda > \|T\|$, $P_\lambda = I_H$. Accordingly we can form the operator $S = \int_{-\infty}^{+\infty} \lambda dP_\lambda$, where the integral converges in operator norm, which lies in B_{00} . We also have $T = \int_{-\infty}^{+\infty} \lambda dE_\lambda$.

Suppose $\xi \in K$. The monotonically increasing function $\|E_\lambda \xi\|$ has only a countable number of discontinuities and so for each $\varepsilon > 0$ we can find $\lambda_0 \leq \lambda_1 \leq \dots \leq \lambda_n$, points of continuity of $\|E_\lambda \xi\|$ such that

$$\left\| S - \sum_{i=1}^n \lambda_i (P_{\lambda_i} - P_{\lambda_{i-1}}) \right\| < \varepsilon$$

$$\left\| T - \sum_{i=1}^n \lambda_i (E_{\lambda_i} - E_{\lambda_{i-1}}) \right\| < \varepsilon.$$

At a continuity point λ of $\|E_\lambda \xi\|$ we have

$$\begin{aligned} \xi &= E_\lambda \xi + (P_K - E_\lambda) \xi \\ &= E_{\lambda-0} \xi + (P_K - E_{\lambda+0}) \xi, \end{aligned}$$

thus, at continuity points of $\|E_\lambda \xi\|$, $P_\lambda \xi = E_{\lambda-0} \xi = E_\lambda \xi$ and so

$$\sum_{i=1}^n \lambda_i (P_{\lambda_i} - P_{\lambda_{i-1}}) \xi = \sum_{i=1}^n \lambda_i (E_{\lambda_i} - E_{\lambda_{i-1}}) \xi.$$

Hence $\|S\xi - T\xi\| < 2\varepsilon \|\xi\|$, and since this holds for all $\varepsilon > 0$, $S\xi = T\xi$ for all $\xi \in K$. Thus we have successively $SL\xi = TL\xi$ for all $L \in B$, $\xi \in H$; $SL = TL$ for all $L \in B$; and, using the fact that S, T, B are self-adjoint, $LS = LT$ for all $L \in B$. Thus $S \in B_{00}$ and determines the same double centraliser on B as T , that is $(\mathcal{S}, \mathcal{S})$ is determined by $S \in B_{00}$.

COROLLARY. *A is a QW*-algebra if and only if it is an AW*-algebra.*

Proof. Follows from the theorem and [2; Theorem 1].

REFERENCES

1. I. Kaplansky, *Projections in banach algebras*, Ann. of Math. **53** (1951), 235-249.
2. G. A. Reid, *A generalisation of W^* -algebras*, Pacific J. Math. **15** (1965), 1019-1026.
3. C. E. Rickart, *General Theory of Banach Algebras*, Van Nostrand, New York, 1960.
4. F. Riesz and B. Sz.-Nagy, *Functional Analysis*, Ungar, New York, 1955.

Received May 25, 1966.

THE UNIVERSITY
NEWCASTLE UPON TYNE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
Rice University
Houston, Texas 77001

J. P. JANS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Pacific Journal of Mathematics

Vol. 23, No. 1

March, 1967

M. J. C. Baker, <i>A spherical Helly-type theorem</i>	1
Robert Morgan Brooks, <i>On locally m-convex $*$-algebras</i>	5
Lindsay Nathan Childs and Frank Rimi DeMeyer, <i>On automorphisms of separable algebras</i>	25
Charles L. Fefferman, <i>A Radon-Nikodym theorem for finitely additive set functions</i>	35
Magnus Giertz, <i>On generalized elements with respect to linear operators</i>	47
Mary Gray, <i>Abelian objects</i>	69
Mary Gray, <i>Radical subcategories</i>	79
John A. Hildebrant, <i>On uniquely divisible semigroups on the two-cell</i>	91
Barry E. Johnson, <i>AW$*$-algebras are QW$*$-algebras</i>	97
Carl W. Kohls, <i>Decomposition spectra of rings of continuous functions</i>	101
Calvin T. Long, <i>Addition theorems for sets of integers</i>	107
Ralph David McWilliams, <i>On w^*-sequential convergence and quasi-reflexivity</i>	113
Alfred Richard Mitchell and Roger W. Mitchell, <i>Disjoint basic subgroups</i>	119
John Emanuel de Pillis, <i>Linear transformations which preserve hermitian and positive semidefinite operators</i>	129
Qazi Ibadur Rahman and Q. G. Mohammad, <i>Remarks on Schwarz's lemma</i>	139
Neal Jules Rothman, <i>An L^1 algebra for certain locally compact topological semigroups</i>	143
F. Dennis Sentilles, <i>Kernel representations of operators and their adjoints</i>	153
D. R. Smart, <i>Fixed points in a class of sets</i>	163
K. Srinivasacharyulu, <i>Topology of some Kähler manifolds</i>	167
Francis C.Y. Tang, <i>On uniqueness of generalized direct decompositions</i>	171
Albert Chapman Vosburg, <i>On the relationship between Hausdorff dimension and metric dimension</i>	183
James Victor Whittaker, <i>Multiply transitive groups of transformations</i>	189