

Pacific Journal of Mathematics

REMARKS ON SCHWARZ'S LEMMA

QAZI IBADUR RAHMAN AND Q. G. MOHAMMAD

REMARKS ON SCHWARZ'S LEMMA

Q. I. RAHMAN AND Q. G. MOHAMMAD

If $f(z)$ is an analytic function, regular in $|z| \leq 1$, $|f(z)| \leq 1$ for $|z| = 1$ and $f(0) = 0$, then by Schwarz's lemma

$$|f(re^{i\theta})| \leq r, \quad (0 \leq r \leq 1).$$

More generally, if $f(z)$ is regular inside and on the unit circle, $|f(z)| \leq 1$ on the circle and $f(a) = 0$, where $|a| < 1$, then

$$(1) \quad |f(z)| \leq |(z-a)/(1-\bar{a}z)|,$$

inside the circle. In other words,

$$(2) \quad |f(z)/(z-a)| < 1/|1-\bar{a}z|,$$

for $|z| \leq 1$. For a fixed a on the unit circle, let C_a denote the class of functions $f(z)$ which are regular in $|z| \leq 1$, vanish at the point $z = a$, and for which

$$\max_{|z|=1} |f(z)| = 1.$$

Any positive number A being given, it is clearly possible to construct a function $f(z)$ of the class C_a for which

$$\mathcal{M}_f(1) = \max_{|z|=1} |f(z)/(z-a)| > A,$$

i.e. $\mathcal{M}_f(1)$ is not uniformly bounded for $f \in C_a$. If $f(z)$ is restricted to a subclass of C_a there may exist a uniform bound for $\mathcal{M}_f(1)$ as $f(z)$ varies within the subclass. It is clear that such is the case for the important subclass consisting of all polynomials of degree at most n vanishing at $z = a$. The problem is to find the uniform bound.

We prove:

THEOREM. *If $p(z)$ is a polynomial of degree n such that $|p(z)| \leq 1$ on the unit circle, and $p(1) = 0$, then for $|z| \leq 1$.*

$$(3) \quad |p(z)/(z-1)| \leq n/2.$$

The example $(z^n - 1)/2$ shows that the bound in (3) is precise. The following corollary is immediate.

COROLLARY 1. *If $p(z)$ is a polynomial of degree n satisfying the conditions of the theorem, then $|p'(1)| \leq n/2$.*

This is interesting in view of the fact that if $p(1) \neq 0$ all we can say [2, p. 357] is that $|p'(1)| \leq n$.

If $p(z)$ is a polynomial of degree n and $|p(z)| \leq 1$ on the unit circle then the polynomial

$$\frac{p(z) - p(1)}{1 + |p(1)|}$$

satisfies the hypotheses of our theorem. Consequently

$$\left| \frac{p(z) - p(1)}{z - 1} \right| \leq \frac{n}{2}(1 + |p(1)|),$$

or

$$|p(z) - p(1)| \leq \frac{n}{2}(1 + |p(1)|)|z - 1|.$$

Thus we have:

COROLLARY 2. *If $p(z)$ is a polynomial of degree n and $|p(z)| < 1$ on the unit circle, then for $|z| \leq 1$,*

$$|p(z)| \leq |p(1)| + \frac{n}{2}(1 + |p(1)|)|z - 1|.$$

Proof of the theorem. Set $p(z)/(z - 1) = \tilde{p}(z)$ and let the maximum of $|\tilde{p}(e^{i\theta})|$ for $-\pi \leq \theta \leq \pi$ occur when $\theta = 2\theta_0$. We may suppose that $|e^{2i\theta_0} - 1| < 2/n$, otherwise there is nothing to prove. Let

$$t(\theta) = e^{-i(n-1)\theta} \tilde{p}(e^{2i\theta}),$$

and choose γ such that $e^{i\gamma}t(\theta_0)$ is real. Consider the real trigonometric polynomial

$$T(\theta) \equiv \operatorname{Re} \{e^{i\gamma}t(\theta)\}.$$

Since $|e^{i\gamma}t(\theta)|$ has its maximum at θ_0 , the real trigonometric polynomial $T(\theta)$ has its maximum modulus at θ_0 where it is actually a local maximum, i.e. $T'(\theta_0) = 0$. The function $2 \sin \theta T(\theta)$ is a real trigonometric polynomial of degree n such that

$$|2 \sin \theta T(\theta)| = |e^{-i\theta}(e^{2i\theta} - 1)T(\theta)| \leq |p(e^{2i\theta})| \leq 1$$

for $-\pi \leq \theta \leq \pi$.

A result of van der Corput and Schaake [1] states that if $F(\theta)$ is a real trigonometric polynomial of degree n and $|F(\theta)| \leq 1$ for real θ , then

$$(4) \quad n^2 F(\theta)^2 + (F'(\theta))^2 \leq n^2.$$

Applying this result to the trigonometric polynomial $2 \sin \theta T(\theta)$ we

get

$$n^2 \sin^2 \theta T^2(\theta) + \{\cos \theta T(\theta) + \sin \theta T'(\theta)\}^2 \leq \frac{n^2}{4}$$

for $-\pi \leq \theta \leq \pi$. Setting $\theta = \theta_0$ we get

$$\{1 + (n^2 - 1) \sin^2 \theta_0\} T^2(\theta_0) \leq \frac{n^2}{4},$$

since $T'(\theta_0) = 0$. Hence

$$|T(\theta_0)| \leq (n/2)(1 + (n^2 - 1) \sin^2 \theta_0)^{-1/2},$$

and the result follows.

REMARK 1. From the method of proof it is clear that if $p(z)$ is a polynomial of degree n , such that $|p(z)| \leq 1$ on the unit circle and $p(\pm 1) = 0$, then for $|z| \leq 1$,

$$(5) \quad |p(z)/(z^2 - 1)| \leq n/4.$$

Or, more generally, if $p(z) = 0$ whenever z is a root of $z^k - 1 = 0$, and $|p(z)| \leq 1$ for $|z| \leq 1$, then

$$(6) \quad \max_{|z| \leq 1} |p(z)/(z^k - 1)| \leq n/(2k).$$

REMARK 2. For every $\varepsilon > 0$ and a lying inside the unit circle we can construct a polynomial $p(z)$ which vanishes at $z = a$, $|p(z)| \leq 1$ for $|z| \leq 1$ but

$$\max_{|z| \leq 1} |p(z)/(z - a)| > \frac{1}{1 - |a|} - \varepsilon.$$

In fact, let

$$\frac{1}{1 - \bar{a}z} = \sum_{\nu=0}^{\infty} a_{\nu} z^{\nu}$$

for $|z| < 1/|a|$, and choose N so large that

$$\left| \frac{1}{1 - \bar{a}z} - \sum_{\nu=0}^N a_{\nu} z^{\nu} \right| < \varepsilon' = \frac{(1 - |a|)\varepsilon}{2 - \varepsilon(1 - |a|^2)}$$

for $|z| \leq 1$. With this choice of N ,

$$\left| (z - a) \sum_{\nu=0}^N a_{\nu} z^{\nu} \right| < 1 + (1 + |a|)\varepsilon'$$

for $|z| \leq 1$ and

$$\frac{1}{1 + (1 + |a|)\varepsilon'} (z - a) \sum_{\nu=0}^N a_{\nu} z^{\nu}$$

is therefore a polynomial of the type we wanted to construct.

However, if we restrict ourselves to polynomials of degree at most $n(>1)$ then we can easily prove that for $|z| \leq 1$

$$(7) \quad |p(z)/(z - a)| < n.$$

For $a = 0$ the result is included in Schwarz's lemma. If $a \neq 0$ we observe that

$$p(z) = \int_a^z p'(w) dw$$

and therefore

$$|p(z)| \leq |z - a| \max_{|w| \leq 1} |p'(w)| < n |z - a|$$

by an inequality due to M. Riesz [2, p. 357] wherein equality holds only when $p(z)$ is a constant multiple of z^n . This is the same as (7).

REFERENCES

1. J. G. van der Corput and G. Schaake, *Ungleichungen für Polynome und trigonometrische Polynome*, *Compositio Mathematica* **2** (1935), 321-361. *Berichtigung zu: Ungleichungen für Polynome und trigonometrische Polynome*, *Compositio Mathematica* **3** (1936), 128.
2. M. Riesz, *Eine trigonometrische Interpolationformel und einige Ungleichungen für Polynome*, *Jahresbericht der Deutschen Mathematiker-Vereinigung* **23** (1914), 354-368.

Received April 15, 1966.

UNIVERSITÉ DE MONTRÉAL

AND

UNIVERSITY OF JAMMU AND KASHMIR

SRINAGAR (KASHMIR), INDIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
Rice University
Houston, Texas 77001

J. P. JANS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Pacific Journal of Mathematics

Vol. 23, No. 1

March, 1967

M. J. C. Baker, <i>A spherical Helly-type theorem</i>	1
Robert Morgan Brooks, <i>On locally m-convex $*$-algebras</i>	5
Lindsay Nathan Childs and Frank Rimi DeMeyer, <i>On automorphisms of separable algebras</i>	25
Charles L. Fefferman, <i>A Radon-Nikodym theorem for finitely additive set functions</i>	35
Magnus Giertz, <i>On generalized elements with respect to linear operators</i>	47
Mary Gray, <i>Abelian objects</i>	69
Mary Gray, <i>Radical subcategories</i>	79
John A. Hildebrant, <i>On uniquely divisible semigroups on the two-cell</i>	91
Barry E. Johnson, <i>AW$*$-algebras are QW$*$-algebras</i>	97
Carl W. Kohls, <i>Decomposition spectra of rings of continuous functions</i>	101
Calvin T. Long, <i>Addition theorems for sets of integers</i>	107
Ralph David McWilliams, <i>On w^*-sequential convergence and quasi-reflexivity</i>	113
Alfred Richard Mitchell and Roger W. Mitchell, <i>Disjoint basic subgroups</i>	119
John Emanuel de Pillis, <i>Linear transformations which preserve hermitian and positive semidefinite operators</i>	129
Qazi Ibadur Rahman and Q. G. Mohammad, <i>Remarks on Schwarz's lemma</i>	139
Neal Jules Rothman, <i>An L^1 algebra for certain locally compact topological semigroups</i>	143
F. Dennis Sentilles, <i>Kernel representations of operators and their adjoints</i>	153
D. R. Smart, <i>Fixed points in a class of sets</i>	163
K. Srinivasacharyulu, <i>Topology of some Kähler manifolds</i>	167
Francis C.Y. Tang, <i>On uniqueness of generalized direct decompositions</i>	171
Albert Chapman Vosburg, <i>On the relationship between Hausdorff dimension and metric dimension</i>	183
James Victor Whittaker, <i>Multiply transitive groups of transformations</i>	189