

Pacific Journal of Mathematics

A NOTE ON PARTIALLY ORDERED COMPACTA

JAMES HARVEY CARRUTH

A NOTE ON PARTIALLY ORDERED COMPACTA

J. H. CARRUTH

In this note it is shown that each compact metric space endowed with a closed partial order admits an equivalent radially convex metric. It is also shown that certain functions from subsets of a partially ordered compactum into the Hilbert Cube are extendable to order preserving homeomorphisms on the whole space.

The notion of a radially convex metric on a partially ordered space was introduced by Kock and McAuley in 1964 [1]. A metric d is *radially convex* with respect to a partial order Γ on X if $(x, y) \in \Gamma$, $(y, z) \in \Gamma$, and $y \neq z$ imply $d(x, y) < d(x, z)$. The natural question "Does each partially ordered metric space admit a radially convex metric?" arises from [1] and [2]. This question is answered in the affirmative for compact spaces. The topological closure of a set $A \subset X$ will be denoted by A^* . The symbol \square will denote the empty set. If Γ is a partial order on X and $A \subset X$ then

$$L(A) = \{x \mid (x, a) \in \Gamma \text{ for some } a \in A\}$$

and

$$M(A) = \{x \mid (a, x) \in \Gamma \text{ for some } a \in A\}.$$

THEOREM 1. *If Γ is a closed partial order on the compact metric space X , then there exists an equivalent metric on X which is radially convex with respect to Γ .*

Proof. Let \mathcal{V} be a countable base for the topology of X and let \mathcal{U} be the collection of finite unions of members of \mathcal{V} . Then \mathcal{U} is countable as is

$$\mathcal{B} = \{(U, V) \mid U, V \in \mathcal{U} \text{ and } M(U^*) \cap L(V^*) = \square\}.$$

Let $\{(U_i, V_i) \mid i = 1, 2, \dots\}$ be an enumeration of \mathcal{B} . For each i , the function f_i defined by

$$f_i(z) = \begin{cases} 1 & \text{if } z \in U_i^* \\ 0 & \text{if } z \in V_i^* \end{cases}$$

is continuous and order preserving. Hence, by a result of Nachbin [3], f_i is extendable to a continuous order preserving function g_i from X into $[0, 1]$. Let Φ be the product function Pg_i which takes X into the Hilbert Cube H . Again using results of [3], it is easy to verify

that $(x, y) \in (X \times X) \setminus \Gamma$ if and only if there exists an integer i such that $f_i(x) > f_i(y)$. It follows that Φ is an order preserving homeomorphism from X into H whose inverse is also order preserving. Hence, we may assume that X is a subspace of H and that Γ is the natural partial order on H restricted to X . It is obvious that the usual metric on H is radially convex with respect to the natural partial order. It is also obvious that radial convexity is hereditary and the proof is complete.

We note that included in the proof of Theorem 1 is the fact that the Hilbert Cube with the natural partial order is a universal partially ordered compactum in the imbedding sense.

If X is partially ordered by Γ , the element x of X is *maximal* (*minimal*) if $M(x) = \{x\}$ ($L(x) = \{x\}$). The element $\{r_i\}$ of the Hilbert Cube H will be called an *interior point* of H if $0 < r_i < 1$ for each i .

THEOREM 2. *Let Γ be a closed partial order on the compact metric space X . Suppose x is maximal, z is minimal, $x \neq z$, and $Y = \{y_i: i = 1, 2, \dots, n\}$ is a finite chain (with respect to Γ) of points of X which are neither maximal nor minimal. Let h be a one-to-one order preserving function from Y into H such that $h(y_i)$ is an interior point of H for each i . Then, there exists an order preserving homeomorphism Φ taking X into H whose restriction to Y is h and with the property that $\Phi(x)_i = 1$ and $\Phi(z)_i = 0$ for each integer i .*

Proof. Let \mathcal{V} be a countable base for the topology of X and let \mathcal{U} be the collection of finite union of members of \mathcal{V} . Assume that $y_i < y_{i+1}$ for each $i = 1, 2, \dots, n - 1$. Let \mathcal{B} be the collection of $n + 2$ tuples (U, V_1, \dots, V_n, W) satisfying the following properties:

- (i) $U, V_1, \dots, V_n, W \in \mathcal{U}$
- (ii) The sets $M(U^*), L(V_1^*) \cap M(V_1^*), \dots, L(V_n^*) \cap M(V_n^*)$, and $L(W^*)$ are pairwise disjoint.
- (iii) $x \in U, y_i \in V_i$ for each i , and $z \in W$.

For each $(U, V_1, \dots, V_n, W)_i \in \mathcal{B}$, define f_i on

$$U^* \cup (\cup \{V_j^*: j = 1, 2, \dots, n\}) \cup W^*$$

by

$$f_i(a) = \begin{cases} 1 & \text{if } a \in U^* \\ h(y_j)_i & \text{if } a \in V_j^* \\ 0 & \text{if } a \in W^* . \end{cases}$$

As in the proof of Theorem 1, we extend f_i to g_i taking X into $[0,1]$ and let Φ be the product of the g_i 's.

It is not difficult to show that the functions $\{g_i\}$ separate points.

This completes the proof.

Note. The inverse of the homeomorphism Φ in Theorem 2 need not be order preserving. In fact, if $L(x) \neq X$, Φ^{-1} is not order preserving.

The importance of the interiority of the points $h(y_i)$ should not be overlooked. Indeed, if $h(y_i)$ is not an interior point of H for some i , the conclusion of Theorem 2 may fail to hold. The following example exhibits this fact.

EXAMPLE. Let X be the space consisting of the unit circle in the plane along with the arc from $(1, 0)$ to $(2, 0)$. Define Γ by $((a, b), (c, d)) \in \Gamma$ if and only if $a \leq c$ and $0 \leq b \cdot d$. Then, Γ is a closed partial order on the compact metric space X . Let $z = (-1, 0)$, $y = (1, 0)$, and $x = (2, 0)$. Now let

$$r_i = \begin{cases} 1 & \text{if } i = 1 \\ 0 & \text{if } i \neq 1. \end{cases}$$

No order preserving function Φ satisfying $\Phi(z)_i = 0$, $\Phi(y)_i = r_i$, and $\Phi(x)_i = 1$ for each i can be a homeomorphism.

REFERENCES

1. R. J. Koch and L. F. McAuley, *Semigroups on trees*, Fund. Math. **50** (1962), 341-346.
2. ———, *Semigroups on continua ruled by arcs*, Fund. Math. **56** (1964), 1-8.
3. L. Nachbin, *Topology and Order*, Van Nostrand Math. Studies, 1965.
4. L. E. Ward, Jr., *Binary relations in topological spaces*, Acad. Brasileria de Ciencias, **26** (1954), 357-373.

Received February 6, 1967, and in revised form May 10, 1967. Presented to the American Mathematical Society on November 12, 1961. This note forms part of the author's doctoral dissertation prepared under the direction of Professor R. J. Koch at Louisiana State University.

UNIVERSITY OF TENNESSEE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
Rice University
Houston, Texas 77001

J. P. JANS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California 90024.

Each author of each article receives 50 reprints free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners of publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 24, No. 2

June, 1968

John Suemper Alin and Spencer Ernest Dickson, <i>Goldie's torsion theory and its derived functor</i>	195
Steve Armentrout, Lloyd Lesley Lininger and Donald Vern Meyer, <i>Equivalent decomposition of R^3</i>	205
James Harvey Carruth, <i>A note on partially ordered compacta</i>	229
Charles E. Clark and Carl Eberhart, <i>A characterization of compact connected planar lattices</i>	233
Allan Clark and Larry Smith, <i>The rational homotopy of a wedge</i>	241
Donald Brooks Coleman, <i>Semigroup algebras that are group algebras</i>	247
John Eric Gilbert, <i>Convolution operators on $L^p(G)$ and properties of locally compact groups</i>	257
Fletcher Gross, <i>Groups admitting a fixed-point-free automorphism of order 2^n</i>	269
Jack Hardy and Howard E. Lacey, <i>Extensions of regular Borel measures</i> ...	277
R. G. Huffstutler and Frederick Max Stein, <i>The approximation solution of $y' = F(x, y)$</i>	283
Michael Joseph Kascic, Jr., <i>Polynomials in linear relations</i>	291
Alan G. Konheim and Benjamin Weiss, <i>A note on functions which operate</i>	297
Warren Simms Loud, <i>Self-adjoint multi-point boundary value problems</i>	303
Kenneth Derwood Magill, Jr., <i>Topological spaces determined by left ideals of semigroups</i>	319
Morris Marden, <i>On the derivative of canonical products</i>	331
J. L. Nelson, <i>A stability theorem for a third order nonlinear differential equation</i>	341
Raymond Moos Redheffer, <i>Functions with real poles and zeros</i>	345
Donald Zane Spicer, <i>Group algebras of vector-valued functions</i>	379
Myles Tierney, <i>Some applications of a property of the functor Ef</i>	401