

# Pacific Journal of Mathematics

**ON THE TETRAHEDRAL GRAPH**

MARTIN AIGNER

## ON THE TETRAHEDRAL GRAPH

MARTIN AIGNER

Generalizing the concept of the triangular association scheme, Bose and Laskar introduced the tetrahedral graph the vertices of which are the  $\binom{n}{3}$  unordered triplets selected from  $n$  symbols with two points adjacent if and only if their corresponding triplets have two symbols in common. If we let  $d(x, y)$  denote the distance between two vertices  $x, y$  and  $\Delta(x, y)$  the number of vertices adjacent to both  $x$  and  $y$ , then the tetrahedral graph possesses the following 4 properties:

- (B0) the number of vertices is  $\binom{n}{3}$
- (B1) it is connected and regular of degree  $3(n - 3)$
- (B2) if  $d(x, y) = 1$  then  $\Delta(x, y) = n - 2$
- (B3) if  $d(x, y) = 2$  then  $\Delta(x, y) = 4$ .

The question whether these conditions characterize tetrahedral graphs (no loops or parallel edges permitted) was answered in the affirmative by Bose and Laskar for  $n > 16$ . In the present paper characterizations of tetrahedral graphs are derived by strengthening each one of (B1), (B2), (B3) and these results are utilized to prove the sufficiency of (B0)–(B3) for  $n=6$ . (For  $n < 4$  the problem is void,  $n = 4, 5$  are trivial cases.)

All graphs considered in this paper are finite undirected without loops or parallel edges. As is readily seen the line-graph  $G$  of the complete graph with  $n$  vertices may be defined as a graph whose vertices are the  $\binom{n}{2}$  unordered pairs taken from  $n$  symbols so that two pairs are adjacent if and only if they have a symbol in common. Letting  $d(x, y)$  denote the distance between  $x$  and  $y$  and  $\Delta(x, y)$  the number of vertices that are adjacent to both  $x$  and  $y$ , then  $G$  has the following properties:

- (A0) the number of vertices is  $\binom{n}{2}$
- (A1)  $G$  is connected and regular of degree  $2(n - 2)$ .
- (A2)  $d(x, y) = 1$  implies  $\Delta(x, y) = n - 2$
- (A3)  $d(x, y) = 2$  implies  $\Delta(x, y) = 4$ .

Conner [2], Shrikhande [7], Hoffman [3, 4] and Li-chien [5, 6] showed that (A0)–(A3) completely characterize linegraphs of complete graphs except for  $n = 8$  where 3 nonisomorphic graphs satisfying (A0)–(A3) exist. Bose and Laskar [1] took up the similar problem concerning unordered triplets chosen from  $n$  symbols we mentioned above.

For  $n > 16$  (B0)–(B3) characterize tetrahedral graphs as was shown by Bose and Laskar in [1].

For  $n < 4$  the characterization problem is meaningless.

For  $n = 4$  a graph  $G$  satisfying (B0)–(B3) is necessarily the complete graph with 4 vertices and the vertices may be identified with the 4 unordered triplets chosen from 4 symbols.

For  $n = 5$  the conditions (B0)–(B3) are identical with (A0)–(A3) hence a graph  $G$  satisfying (B0)–(B3) may be assigned as vertices the 10 unordered pairs of symbols taken from a set of 5 symbols with two vertices adjacent if and only if their corresponding pairs share a symbol. Replacing each pair by its complement in the set of the 5 symbols we obtain a graph  $G$  with triplets assigned to its vertices and  $G$  is readily seen to be tetrahedral.

In the following we assume  $n \geq 6$ .  $K_i$  will denote the complete graph with  $i$  vertices,  $S(x)$  the set of the vertices adjacent to  $x$ ,  $T(x, y)$  the set of the vertices adjacent to both  $x$  and  $y$ .

## II. Characterizations of tetrahedral graphs.

LEMMA 1. *For a graph  $G$  satisfying (B0)–(B3) the following properties are equivalent:*

(C1) *For all  $x \in G$  the subgraph induced by  $S(x)$ <sup>1</sup> can be partitioned into  $3K_{n-3}$ 's:*

$$X = \{x_1, x_2, \dots, x_{n-3}\}, Y = \{y_1, y_2, \dots, y_{n-3}\}, Z = \{z_1, z_2, \dots, z_{n-3}\}$$

*such that  $\{x_i, y_i, z_i\}$  induces a  $K_3$  for  $i = 1, \dots, n - 3$ .*

(C2) *For all  $x, y \in G$  with  $d(x, y) = 1$  the subgraph induced by  $T(x, y)$  consists of a  $K_{n-4}$  and a  $K_2$  such that no vertex in  $K_{n-4}$  is adjacent to either vertex in  $K_2$ .*

(C3) *For all  $x \in G$  the subgraph induced by  $S(x)$  can be partitioned into  $3K_{n-3}$ 's such that for any pair  $y, z \in S(x)$  with  $d(y, z) = 2$  there are exactly 2 other vertices  $v, w \in S(x)$  which are adjacent to both  $y, z$ .*

*Proof.* It is evident that property (C1) implies both (C2) and (C3). On the other hand assume (C2) and let  $x_1 \in S(x)$ . In  $T(x, x_1)$  let  $x_2, \dots, x_{n-3}$  be the vertices in  $K_{n-4}$  and let  $y_1, z_1$  be those in  $K_2$ . It follows from (C2) that in  $T(x, y_1)$  the  $K_2$ -part is constituted by  $x_1, z_1$  and further that the  $n - 4$  remaining vertices  $y_2, \dots, y_{n-3}$  form the  $K_{n-4}$ -part and are distinct from  $x_2, \dots, x_{n-3}$ . Similarly for the pair  $x, z_1$  the set  $T(x, z_1)$  is made up by  $x_1, y_1$  as  $K_2$ -part and by  $n - 4$  vertices  $z_2, \dots, z_{n-3}$  different from  $x_1, y_i (i = 1, \dots, n - 3)$ . Hence  $S(x)$  has the form displayed in Fig. 1. (B2) implies that each of  $x_i, y_i, z_i (i = 2, \dots, n - 3)$  is adjacent to exactly 2 vertices in  $S(x)$  outside its own  $K_{n-3}$ .

<sup>1</sup> By the subgraph induced by a set  $S$  of vertices in  $G$  we mean the subgraph which has  $S$  as vertex-set and includes all edges between any two points in  $S$ .

<sup>2</sup> By (B2) it is clear that there exist no edges joining vertices of one  $K_{n-3}$  to another other than those of the specified  $K_3$ 's.




FIG. 1

If e.g.  $x_i$  were adjacent to  $y_j, y_k (2 \leq j, k)$  then the subgraph induced by  $T(x, y_j)$  would consist of a  $K_{n-4}$  and a  $K_2$  with  $y_k \in K_{n-4}, x_i \in K_2$  and  $d(x_i, y_k) = 1$ , thus violating (C2). Hence each of  $x_i, y_i, z_i$  is adjacent to exactly one vertex of the two  $K_{n-3}$ 's not containing it—hence (C1) holds.

Next let us assume (C3). Let  $X, Y, Z$  be the 3  $K_{n-3}$ 's of  $S(x)$  as in (C1). Given  $x_1 \in X$ : In order to prove (C1) we have to exclude the following two possibilities:

(A)  $x_1$  is adjacent to two vertices, say  $y_1, y_2 \in Y$ , lying in the same  $K_{n-3}$ .

(B)  $x_1$  is adjacent to say  $y_1 \in Y, z_1 \in Z$  in different  $K_{n-3}$ 's but  $d(y_1, z_1) = 2$ .

Suppose (A): The set  $Y - \{y_1, y_2\}$  is nonempty (since  $n \geq 6$ ) and by (C3) no  $y \in Y - \{y_1, y_2\}$  can be adjacent to any  $x \in X$ . Hence let us assume  $y_3 \in Y - \{y_1, y_2\}$  is adjacent to  $z_1, z_2 \in Z$ . Since  $z_1$  is adjacent to exactly two points in  $S(x)$  outside  $Z$  (one of them being  $y_3$ ) we conclude there is at most one vertex in  $S(x)$  adjacent to both  $x_1$  and  $z_1$ , thus contradicting (C3).

Suppose (B): By (C3) either  $y_1$  is adjacent to some  $z \in Z$  in which case  $z_1$  must be adjacent to some  $x_2 \in X$ , or  $y_1$  is adjacent to some  $x_2 \in X$  with  $z_1$  adjacent to some  $y_2 \in Y$  or to  $x_3$  also. Either possibility brings us back to case (A) with  $z_1$  respectively  $y_1$  playing the role of  $x_1$  in (A).

REMARK 1. In a graph  $G$  satisfying (B0)–(B3) condition (C1) implies (C3'): For any pair of vertices  $x_1, y_1$  with  $d(x_1, y_1) = 2$  the subgraph induced by the 4 vertices  $x_2, x_3, y_2, y_3$  adjacent to both  $x_1, y_1$  is a cycle.

*Proof.* In the subgraph induced by  $S(x_3), x_1$  and  $y_1$  are in different  $K_{n-3}$ 's with  $x_1$  adjacent to a vertex, say  $x_2$ , in the  $K_{n-3}$  containing  $y_1$ , and  $y_1$  in turn adjacent to  $y_2$  in the  $K_{n-3}$  containing  $x_1$ . We have  $d(x_2, y_2) = 2$  and no other vertices in  $S(x_3)$  are adjacent to both  $x_1, y_1$ . Now let us consider  $S(x_2)$ . There  $x_3, y_1$  are in the same  $K_{n-3}, x_1$  is in another and  $y_1$  is adjacent to a vertex  $y_3$  in the  $K_{n-3}$  which contains  $x_1$ . Since  $y_3$  evidently is different from  $y_2$ , it must be the fourth point adjacent to  $x_1, y_1$ ; furthermore  $d(x_3, y_3) = 2, d(x_2, y_3) = 1$ . Similarly one

gets  $d(y_2, y_3) = 1$ ; hence  $x_2, x_3, y_2, y_3$  induce a cycle.

REMARK 2. It can be shown with only a little difficulty that for  $n = 6$  the converse of Remark 1 holds i.e. (C3') implies (C1) in a graph satisfying (B0)–(B3). As we will not make use of this fact subsequently the proof is omitted.

LEMMA 2. Let a graph  $G$  satisfy (B0)–(B3) and (C1). Let  $X, Y, Z$  be as in Lemma 1, i.e.  $S(x) = X \cup Y \cup Z$ . Then

$$S(z_i) = (T(x, z_i) - \{x_i, y_i\}) \cup \{x\} \cup (T(x_i, z_i) - \{x, y_i\}) \cup \{x_i\} \\ \cup (T(y_i, z_i) - \{x, x_i\}) \cup \{y_i\} \quad \text{for all } i$$

with each one of the sets on the right hand side inducing a  $K_{n-3}$ . For  $S(x_i), S(y_i)$  the analogous statements hold.

*Proof.* Follows instantly from (C1).

Note that Lemma 2 implies that  $S(z_i)$  is completely determined by  $S(x), S(x_i), S(y_i)$ .

THEOREM 1. A graph  $G$  is tetrahedral if and only if  $G$  satisfies (B0)–(B3) and any (and hence all) of the conditions (C1)–(C3).

*Proof.* Necessity follows readily from the definition. To prove the sufficiency let us first interpret tetrahedral graphs geometrically.

For  $n \geq 6$  let  $C_n = \{(i, j, k) \mid 1 \leq i, j, k \leq n; i, j, k \text{ integral}\}$  i.e., the set of all integral lattice points of the 3-cube with sides extending from 1 to  $n$ . Let  $C'_n = \{(i, j, k) \mid 1 \leq i, j, k \leq n; i \neq j \neq k \neq i; i, j, k \text{ integral}\}$  then  $|C'_n| = n(n-1)(n-2)$ . Now it is evident that  $G$  is tetrahedral if and only if its vertices can be identified with the lattice points in  $C'_n$  (each vertex appears exactly 6 times in  $C'_n$ ) such that two vertices are adjacent if and only if they lie on a straight line parallel to a coordinate-axis.

Thus in order to prove the theorem it suffices to show that the vertices of a graph  $G$  satisfying (B0)–(B3) and say (C1) can be arranged in  $C'_n$  in the above fashion.

For simplicity let us denote the vertices of  $G$  by the natural numbers from 1 to  $\binom{n}{3}$ .

Let

$$\left\{ \begin{array}{l} 2, \dots, n-2 \\ n-1, \dots, 2n-5 \\ 2n-4, \dots, 3n-8 \end{array} \right\}$$

be the 3  $K_{n-3}$ 's constituting  $S(1)$  with numbers in the same column induc-

ing  $K_3$ 's. Place 1 at the spots  $(3, 2, 1), (2, 3, 1), (1, 3, 2)$  in  $C'_n$  and  $\{2, \dots, n - 2\}$  on the lattice points  $\{(i, 2, 1) \mid 4 \leq i \leq n\}$  in this order,  $\{n - 1, \dots, 2n - 5\}$  on  $\{(i, 3, 1) \mid 4 \leq i \leq n\}$ , and  $\{2n - 4, \dots, 3n - 8\}$  on  $\{(i, 3, 2) \mid 4 \leq i \leq n\}$ .

Next we look at the set  $S(2)$ . Besides  $\{1, 3, \dots, n - 2\}$  inducing a  $K_{n-3}$ , there are two more  $K_{n-3}$ 's one headed by  $n - 1$ , the other by  $2n - 4$ . ( $n - 1, 2n - 4$  cannot be in the same  $K_{n-3}$  since both are adjacent to 1.)

Thus

$$S(2) = \begin{pmatrix} 1, & 3, \dots, n - 2 \\ n - 1, 3n - 7, \dots, 4n - 12 \\ 2n - 4, 4n - 11, \dots, 5n - 16 \end{pmatrix}$$

with numbers in the same column inducing  $K_3$ 's.

Now place 2 at the spots  $\{2, 4, 1\}$  and  $(1, 4, 2)$  in  $C'_n$ ,  $\{n - 1, 3n - 7, \dots, 4n - 12\}$  on the line  $\{(i, 4, 1) \mid 3 \leq i \leq n\}$ ,  $\{2n - 4, 4n - 11, \dots, 5n - 16\}$  on the line  $\{(i, 4, 2) \mid 3 \leq i \leq n\}$ . Now the situation is as follows (Fig. 2): We claim:

$$\begin{aligned} d(n, 3n - 7) &= d(2n - 3, 4n - 11) = 1 \\ d(n + 1, 3n - 6) &= d(2n - 2, 4n - 10) = 1 \\ &\vdots \\ d(n, 4n - 11) &= d(2n - 3, 3n - 7) = 2 \\ d(n + 1, 4n - 10) &= d(2n - 2, 3n - 6) = 2 \\ &\vdots \end{aligned}$$

It now follows from  $d(2, n) = 2$  and Remark 1 that the fourth point  $i$  beside  $1, 3, n - 1$  adjacent to both  $2, n$  has to satisfy  $d(i, 1) = 2$ ,  $d(i, 3) = d(i, n - 1) = 1$ . Hence  $i$  must be  $3n - 7$  and furthermore  $d(n, 4n - 11) = 2, d(2n - 3, 4n - 11) = 1, d(2n - 3, 3n - 7) = 2$ . A similar argument proves the other assertions. That no other points among the ones introduced thus far are adjacent beside those already mentioned also follows easily with the help of Remark 1.

Next we consider  $S(3)$ :  $\{1, 2, 4, \dots, n - 2\}$  induce one  $K_{n-3}$ ; we have already found that  $n, 2n - 3, 3n - 7, 4n - 11$  are also in  $S(3)$ . Since  $n, 2n - 3$  are both adjacent to 1 they must be in different  $K_{n-3}$ 's, and so must  $3n - 7, 4n - 11$  be in different  $K_{n-3}$ 's. Hence

$$S(3) = \begin{pmatrix} 1, & 2, & 4, \dots, n - 2 \\ n, 3n - 7, 5n - 15, \dots, 6n - 21 \\ 2n - 3, 4n - 11, 6n - 20, \dots, 7n - 26 \end{pmatrix}$$

with elements in the same column inducing  $K_3$ 's.




FIG. 2

Place 3 at  $(2, 5, 1)$  and  $(1, 5, 2)$  in  $C'_n, \{n, 3n - 7, \dots, 6n - 21\}$  on the line  $\{(i, 5, 1) \mid 3 \leq i \leq n\}$ ,  $\{2n - 3, \dots, 7n - 26\}$  on the line  $\{(i, 5, 2) \mid 3 \leq i \leq n\}$ .

We claim:

- (1)  $d(n + 1, 5n - 15) = d(3n - 6, 5n - 15) = 1$   
 $d(2n - 2, 6n - 20) = d(4n - 10, 6n - 20) = 1$
- (2)  $d(n + 1, 6n - 20) = d(3n - 6, 6n - 20) = 1$   
 $d(2n - 2, 5n - 15) = d(4n - 10, 5n - 15) = 1$

and similarly for the lines parallel to the  $y$ -axis starting at  $n + 2, n + 3, \dots$  resp.  $2n - 1, 2n, \dots$ . (1) follows as before by considering  $3, n + 1; 3, 3n - 6; 3, 2n - 2, 3, 4n - 10$ , and (2) is a consequence of (1). Again we note that no other edges beside those already mentioned exist between vertices 1 to  $7n - 26$ .

In this way one considers all sets  $S(a)$  for  $1 \leq a \leq n - 2$  and fills the lattice points  $\{(i, j, k) \mid 1 \leq i \leq n, 1 \leq j \leq n, k = 1, 2\}$  in  $C'_n$  in the same fashion, thus obtaining  $\binom{n-1}{2} + \binom{n-2}{2} = (n-2)^2$  vertices.

Now we turn to vertices corresponding to points  $\{(i, j, k) \mid j = 3, k = 1\}$  in  $C'_n$ . In  $S(n - 1)$  two  $K_{n-3}$ 's are already known:  $\{1, n, n + 1, \dots, 2n - 5\}$ ,  $\{2, 3n - 7, \dots, 4n - 12\}$  with  $1 = d(1, 2) = d(n, 3n - 7) = \dots$  and further  $2n - 4 \in S(n - 1)$  Hence

$$S(n-1) \left\{ \begin{array}{l} 1, \quad n, \quad n+1, \dots, 2n-5 \\ 2, \quad 3n-7, \quad 3n-6, \dots, 4n-12 \\ 2n-4, (n-2)^2+1, (n-2)^2+2, \dots, (n-2)^2+(n-4) \end{array} \right\}$$

with numbers in the same column inducing  $K_3$ 's.

Place  $n-1$  at  $(1, 4, 3)$  and  $\{2n-4, (n-2)^2+1, \dots, (n-2)^2+(n-4)\}$  on the line  $\{(i, 4, 3) \mid 2 \leq i \leq n\}$  in  $C'_n$ .

We claim:

$$\begin{aligned} d(2n-3, (n-2)^2+1) &= d(4n-11, (n-2)^2+1) = 1 \\ d(2n-2, (n-2)^2+2) &= d(4n-10, (n-2)^2+2) = 1 \\ &\vdots \end{aligned}$$

This assertion is verified by considering the pairs  $2n-3, n-1; 4n-11, n-1; 2n-2, n-1; 4n-10, n-1; \dots$  and applying Remark 1.

In this manner we fill up all the lattice points  $\{(i, j, k) \mid 1 \leq i \leq n, 4 \leq j \leq n, k=3\}$  thus obtaining  $\binom{n-3}{2}$  new vertices. By Lemma 2 the vertices adjacent to  $2n-4, 2n-3, \dots$  i.e. to points on the line  $\{(i, j, k) \mid 4 \leq i, 3=j, 2=k\}$  have already been taken care of, so we may turn to points on the line  $\{(i, j, k) \mid j=4, k=1\}$ . We place  $3n-7$  at  $(1, 5, 4)$  and proceed in the usual manner.

Proceeding in the same fashion we gradually fill up all the lattice points  $\{(i, j, k) \mid j > k\}$  in  $C'_n$  obtaining finally

$$\binom{n-1}{2} + \binom{n-2}{2} + \dots + \binom{3}{2} + \binom{2}{2} = \binom{n}{3}$$

i.e., all vertices of  $G$ .

It is easily seen that reflection about the plane  $y = z$  fills the other half of  $C'_n$  and that by means of this construction one actually arrives at a tetrahedral graph.

### III. The case $n = 6$ .

LEMMA 3. *Given a graph  $G$  which satisfies (B0)–(B3). Let  $D_i(x) = \{y \in G \mid d(x, y) = i\}$   $i \geq 1$ . Then*

$$(3) \quad |D_1(x)| = 3(n-3)$$

$$(4) \quad |D_2(x)| = 3 \binom{n-3}{2}$$

$$(5) \quad |\bigcup_{i>2} D_i(x)| = \binom{n-3}{3}$$

for all vertice  $x \in G$ .

*Proof.* (3) is condition (B1) of the hypothesis. (B2) implies that



there are exactly  $3(n-3) - (n-1) = 2(n-4)$  edges joining an arbitrary vertex in  $D_1(x)$  with  $D_2(x)$ . Now since by (B3) any vertex in  $D_2(x)$  is adjacent to exactly 4 vertices in  $D_1(x)$  we have the equality  $2(n-4) \cdot 3(n-3) = 4 \cdot |D_2(x)|$  and hence (4). (5) is now a consequence of (3) and (4).

COROLLARY. For  $n \leq 8$  the diameter of  $G \leq 3$ .

(5) and the fact that  $\binom{n-3}{3} < 3(n-3)$  for  $n \leq 8$  immediately prove this assertion.

THEOREM 2. For  $n = 6$  a graph  $G$  satisfying (B0)–(B3) is tetrahedral.

*Proof.* In the light of Theorem 1 we have to show that in this case (B0)–(B3) imply any (and hence all) of the conditions (C1)–(C3'). Let us prove that (C2) follows from (B0)–(B3).

Let  $x$  be any vertex, then by Lemma 3

$$|D_1(x)| = 9, |D_2(x)| = 9, |D_3(x)| = 1.$$

Furthermore let  $z$  be such that  $d(x, z) = 3$  then for any

$$y \in D_1(x) \cup D_2(x) (= D_1(z) \cup D_2(z))$$

$$(6) \quad d(x, y) = 1 \iff d(z, y) = 2.$$

For simplicity let us denote the vertices of  $G$  by  $1, 2, \dots, 20$  and suppose that

$$\begin{aligned} D_1(1) &= \{2, 3, \dots, 10\} \\ D_2(1) &= \{11, 12, \dots, 19\} \\ D_3(1) &= \{20\} \end{aligned}$$

with  $d(2, 19) = d(3, 18) = \dots = d(10, 11) = 3$ .

By (B2) 2 is adjacent to 4 vertices in  $D_1(1)$ —say 3, 4, 5, 6. To prove (C2) we have to show each one of 3, 4, 5, 6 is joined to exactly one other vertex of this set.

(6) now implies

$$d(2, 11) = d(2, 12) = d(2, 13) = d(2, 14) = 1.$$

*Case (A).* 3 is adjacent to each one of  $\{4, 5, 6\}$ . Then by (B2)  $d(3, 7) = d(3, 8) = d(3, 9) = d(3, 10) = 2$  and hence 1, 4, 5, 6, 11, 12, 13, 14 would all be adjacent to both 2 and 3, contradicting (B2).

*Case (B).* 3 is adjacent to two of  $\{4, 5, 6\}$  say 4, 5. Then by a similar argument 1, 4, 5 and three vertices among  $\{11, 12, 13, 14\}$  would be adjacent to 2 and 3, a contradiction.

*Case (C).* 3 is adjacent to none of  $\{4, 5, 6\}$ . Then say  $d(3, 7) =$

$d(3, 8) = d(3, 9) = 1$  and hence by (6)  $d(3, 14) = d(3, 13) = d(3, 12) = 2$  leaving only 1 and 11 as vertices adjacent to both 2 and 3—thus again contradicting (B2).

Hence the only possible case: 3 (and similarly 4, 5, 6) is adjacent to exactly one among  $\{3, 4, 5, 6\}$ , thus proving the theorem.

#### REFERENCES

1. R. C. Bose and R. Laskar, *A characterization of tetrahedral graphs* (to appear).
2. W. S. Conner, *The uniqueness of the triangular association scheme*, Ann. Math. Stat. **29** (1958), 262-266.
3. A. J. Hoffman, *On the uniqueness of the triangular association scheme*, Ann. Math. Stat. **31** (1960), 492-497.
4. ———, *On the exceptional case in a characterization of the arcs of a complete graph*, I.B.M.J. **4** (1960), 487-496.
5. Chang Li-chien, *The uniqueness and non-uniqueness of the triangular association schemes*, Science Record, Math. New Ser. **3** (1959), 604-613.
6. ———, *Association schemes of partially balanced designs with parameters  $v = 28$ ,  $n_1 = 12$ ,  $n_2 = 15$  and  $p_{11}^2 = 4$* , Science Record, Math. New Ser. **4** (1960), 12-18.
7. S. S. Shrikhande, *In a characterization of the triangular association scheme*, Ann. Math. Stat. **30** (1959), 39-47.

Received May 19, 1967.

WAYNE STATE UNIVERSITY



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. ROYDEN

Stanford University  
Stanford, California

J. DUGUNDJI

Department of Mathematics  
Rice University  
Houston, Texas 77001

J. P. JANS

University of Washington  
Seattle, Washington 98105

RICHARD ARENS

University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL WEAPONS CENTER

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California 90024.

Each author of each article receives 50 reprints free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners of publishers and have no responsibility for its content or policies.

# Pacific Journal of Mathematics

Vol. 25, No. 2

October, 1968

|   |     |
|---|-----|
| Martin Aigner, <i>On the tetrahedral graph</i> . . . . .  | 219 |
| Gregory Frank Bachelis, <i>Homomorphisms of annihilator Banach algebras</i> . . . . .                               | 229 |
| Phillip Alan Griffith, <i>Transitive and fully transitive primary abelian groups</i> . . . . .                      | 249 |
| Benjamin Rigler Halpern, <i>Fixed points for iterates</i> . . . . .   | 255 |
| James Edgar Keesling, <i>Mappings and dimension in general metric spaces</i> . . . . .                              | 277 |
| Al (Allen Frederick) Kelley, Jr., <i>Invariance for linear systems of ordinary differential equations</i> . . . . . | 289 |
| Hayri Korezlioglu, <i>Reproducing kernels in separable Hilbert spaces</i> . . . . .                                 | 305 |
| Gerson Louis Levin and Wolmer Vasconcelos, <i>Homological dimensions and Macaulay rings</i> . . . . .               | 315 |
| Leo Sario and Mitsuru Nakai, <i>Point norms in the construction of harmonic forms</i> . . . . .                     | 325 |
| Barbara Osofsky, <i>Noncommutative rings whose cyclic modules have cyclic injective hulls</i> . . . . .             | 331 |
| Newton Tenney Peck, <i>Extreme points and dimension theory</i> . . . . .  | 341 |
| Jack Segal, <i>Quasi dimension type. II. Types in 1-dimensional spaces</i> . . . . .                                | 353 |
| Michael Schilder, <i>Expected values of functionals with respect to the Ito distribution</i> . . . . .              | 371 |
| Grigorios Tsagas, <i>A Riemannian space with strictly positive sectional curvature</i> . . . . .                    | 381 |
| John Alexander Williamson, <i>Random walks and Riesz kernels</i> . . . . .  | 393 |