

Pacific Journal of Mathematics

**EXPECTED VALUES OF FUNCTIONALS WITH RESPECT TO
THE ITO DISTRIBUTION**

MICHAEL SCHILDER

EXPECTED VALUES OF FUNCTIONALS WITH RESPECT TO THE ITO DISTRIBUTION

MICHAEL SCHILDER

Stochastic differential equations of the type (written symbolically)

$$(1.1) \quad x^{(n)}(t) + m(t, x^{(n-1)}(t), x^{(n-2)}(t), \dots, x'(t), x(t), t) = z'(t) \\
 x(S) = a_0, x'(S) = a_1, \dots, x^{(n-1)}(S) = a_{n-1} \quad S \leq t \leq T$$

where $z(t)$ is Brownian motion, arise in physics and engineering and are also the object of study of pure mathematicians. In this paper it will be shown that the integral associated with the distribution of the function $x(\cdot)$ may be expressed in terms of a Wiener integral with a weighting functional (the Radon-Nikodym derivative). Thus the expected values of functionals with respect to the distribution of $x(\cdot)$ can be easily and concisely expressed. Also it will be shown that certain partial differential equations of physics naturally have their solutions associated with this integral.

To proceed more precisely, following Ito [7], we rewrite the first order case of (1.1) as

$$(1.1)' \quad x(t) + \int_S^t m(u, x(u))du = z(t)$$

and ask about solutions of (1.1)'. With certain conditions on $m(t, x)$ it is possible to conclude that for almost every $z(\cdot)$ (Wiener or Brownian motion measure) in $C[S, T]$, where $C[S, T]$ is the class of continuous functions on the interval $[S, T]$ whose value at S is a , that (1.1)' has a unique solution $x(\cdot)$. See [2] for example. We express this by writing

$$(1.2) \quad x = Gz .$$

Clearly now, the most natural definition for the distribution of $x(\cdot)$ is as follows:

$$P(x(\cdot) \in A) = P_w(z \in G^{-1}A) .$$

P_w is Wiener measure, and A is a set of functions contained in $C[S, T]$ which is such that $G^{-1}A$ is Wiener measurable. The distribution $P(\cdot)$ so defined will be called the Ito distribution and will be denoted from now on by $P_I(\cdot)$; the associated integral will be denoted by $E_I(\cdot)$.

This approach has been taken before. (See [4] and [5].) However in this paper it will be shown that explicit expressions for the expected values of various functionals arising in physics and engi-

neering can be given in terms of Wiener integrals. In particular a number of the thermodynamic functions can be so expressed. Further, the results of this paper shed some light on the Feynman-Kac [8] formula for Wiener integrals.

2. The integral. In this section, the definition of the Ito distribution which was given in the introduction, will be given rigorously and the integral with respect to it will be defined.

Let T_k be the integral operator

$$T_k(x)(t) = \int_S^t x(u) du + a_{n-k}$$

and let $y(t) = x^{(n-1)}(t)$.

We note that (1.1) may be written

$$(2.1) \quad y(t) + \int_S^t m(u, y(u), T_2(y)(u), \dots, T_n(T_{n-1}, \dots, (T_2 y))(u)) du \\ = z(t).$$

As in the introduction, if certain conditions are put on $m(t, x_{n-1}, \dots, x_0)$, then it follows that for almost every $z \in C[S, T]$ ($z(S) = a_{n-1}$) Wiener measure there is a unique solution to (2.1). We express this again by writing $y = Gz$.

As in the introduction, this transformation induces a σ -field and a measure on the y space, which we again call the Ito measure.

From now on the notation for the sets over which integration is performed will be left out if the meaning is clear.

We now have the following theorem:

THEOREM 1. *If $F(x^{(n-1)}, \dots, x)$ is a functional which is measurable and integrable with respect to the Ito distribution just defined and if for almost every $z \in C[S, T]$ (2.1) has a well defined solution, then*

$$\int F(x^{(n-1)}, \dots, x) dP_I(x^{(n-1)}, \dots, x) \\ = \int F(Gz, T_2 Gz, \dots, T_n(T_{n-1}, \dots, T_2(Gz))) dP_W(z).$$

Proof. The proof follows immediately from the definitions.

There is the rather interesting and useful corollary.

COROLLARY. *Suppose $a_{n-j}(t)$ are continuous real valued functions for $S \leq t \leq T, j = 1, \dots, n$, and that A is the matrix*

$$A(t) = \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ -a_0(t) & -a_1(t) & -a_2(t) & \cdots & a_{n-1}(t) \end{pmatrix}.$$

Let $\Phi(t)$ be the (matrix) fundamental solution to the equation

$$\dot{\Phi}_{ik}(t) = \sum_j^n A_{ij}(t)\Phi_{jk}(t).$$

Then

$$(i) \quad x(t) = \sum_{kl} \Phi_{1k}(t) \int_S \Phi_{kl}^{-1}(u) A_{l_n}(u) z(u) du$$

is a solution to the integro-differential equation

$$x^{(n-1)}(t) + \int_S \sum a_{n-j}(u) x^j(u) du = z(t)$$

$$(ii) \quad x^{(j)}(t) = \sum_{kl} \Phi_{jk}(t) \int_S \Phi_{kl}^{-1}(u) A_{l_n}(u) z(u) du + \delta_{jn-1} z(t)$$

(iii) The Ito distribution defined by

$$(2.2) \quad \tilde{x}^{(n)}(t) + \sum_j a_{n-j}(t) \tilde{x}^j(t) = \dot{z}(t)$$

and

$$\tilde{x}(s) = \tilde{x}^{(1)}(s) = \tilde{x}^{(2)}(s) = \cdots = \tilde{x}^{(n-1)}(s) = z(s) = 0$$

is Gaussian with

$$E_I\{\tilde{x}^{(j)}(t)\} = 0 \quad j = 0, 1, \dots, n-1 \quad \text{and} \quad S \leq t \leq T$$

and

$$\begin{aligned} E_I\{\tilde{x}^{(j_1)}(t_1)\tilde{x}^{(j_2)}(t_2)\} &= \sum_{k_1 l_1 k_2 l_2} \int_S \int_S \Phi_{j_1 k_1}(t_1) \Phi_{k_1 l_1}^{-1}(u_1) A_{l_1 n}(u_1) \Phi_{j_2 k_2}(t_2) \Phi_{k_2 l_2}^{-1}(u_2) \\ &A_{l_2 n}(u_2) \min(u_1 - S, u_2 - S) du_1 du_2 \\ &+ \sum_{k_1 l_1} \Phi_{j_1 k_1}(t_1) \int_S \Phi_{k_1 l_1}^{-1}(u_1) A_{l_1 n}(u_1) \min(u_1 - S, t_2 - S) du_1 \delta_{j_1 n-1} \\ &+ \sum_{k_2 l_2} \Phi_{j_2 k_2}(t_2) \int_S \Phi_{k_2 l_2}^{-1}(u_2) A_{l_2 n}(u_2) A_{l_2 n}(u_2) \min(u_2 - S, t_1 - S) du_2 \delta_{j_2 n-1} \\ &+ \delta_{j_1 n-1} \delta_{j_2 n-1} \min(t_1 - S, t_2 - S). \end{aligned}$$

Proof. (i) and (ii) follow by direct verification, which can be accomplished without differentiating z by using integration by parts. To find the distribution $\tilde{x}^{(j)}(t)$, the method of characteristic functions is used. By definition, the distribution of $\tilde{x}^{(j)}(t)$ is that of $x^j(t)$. Thus from theorem 1, this definition, and (i) and (ii),

$$\begin{aligned} \int \exp [iv\tilde{x}^{(j)}(t)]P_I(\tilde{x}) &= \int \exp [ivx^{(j)}(t)]P_I(x) \\ &= E_w \left(\exp \left[iv \left(\sum_{kl} \Phi_{jk}(t) \int_S^t \Phi_{kl}^{-1}(u) A_{l_n}(u) z(u) du + \delta_{j_{n-1}} z(t) \right) \right] \right). \end{aligned}$$

The last Wiener integral is known to be

$$\begin{aligned} \exp \left[-v^2/2 \left(\sum_{k_1 l_1 k_2 l_2} \int_S^t \int_S^t \Phi_{jk_1}(t) \Phi_{k_1 l_1}^{-1}(u_1) A_{l_1 n}(u_1) \right. \right. \\ \times \Phi_{jk_2}(t) \Phi_{k_2 l_2}^{-1}(u_2) A_{l_2 n}(u_2) \min(u_1 - S, U_2 - S) du_1 du_2 \\ \left. \left. + 2 \sum_{k_1 l_1} \Phi_{jk_1}(t) \int_S^t \Phi_{k_1 l_1}^{-1}(u) A_{l_1 n}(u) \min(u - S, t - S) du \delta_{j_{n-1}} + \delta_{j_{n-1}} t \right) \right]. \end{aligned}$$

Thus $\tilde{x}^{(j)}(t)$ $j = 0, \dots, n-1$ is Gaussian with mean zero and variance

$$\begin{aligned} \sum_{k_1 l_1 k_2 l_2} \int_S^t \int_S^t \Phi_{jk_1}(t) \Phi_{k_1 l_1}^{-1}(u_1) A_{l_1 n}(u_1) \Phi_{jk_2}(t) \\ \times \Phi_{k_2 l_2}^{-1}(u_2) A_{l_2 n}(u_2) \min(u_1 - S, u_2 - S) du_1 du_2 \\ + 2 \sum_{k_1 l_1} \Phi_{jk_1}(t) \int_S^t \Phi_{k_1 l_1}^{-1}(u) A_{l_1 n}(u) \min(u - S, t - S) du \delta_{j_{n-1}} + \delta_{j_{n-1}} t. \end{aligned}$$

To calculate $E_I(\tilde{x}^{j_1}(t_1)\tilde{x}^{j_2}(t_2))$, we note that

$$\begin{aligned} E_I(\tilde{x}^{j_1}(t_1)\tilde{x}^{j_2}(t_2)) &= E_I(x^{j_1}(t_1)x^{j_2}(t_2)) \\ &= E_w \left(\left(\sum_{k_1 l_1} \Phi_{j_1 k_1}(t_1) \int_S^{t_1} \Phi_{k_1 l_1}^{-1}(u_1) A_{l_1 n}(u_1) z(u_1) du_1 + \delta_{j_1 n-1} z(t_1) \right) \right. \\ &\quad \left. \left(\sum_{k_2 l_2} \Phi_{j_2 k_2}(t_2) \int_S^{t_2} \Phi_{k_2 l_2}^{-1}(u_2) A_{l_2 n}(u_2) z(u_2) du_2 + \delta_{j_2 n-1} z(t_2) \right) \right) \\ &= \sum_{k_1 l_1 k_2 l_2} \int_S^{t_1} \int_S^{t_2} \Phi_{j_1 k_1}(t_1) \Phi_{k_1 l_1}^{-1}(u_1) A_{l_1 n}(u_1) \Phi_{j_2 k_2}(t_2) \\ &\quad \times \Phi_{k_2 l_2}^{-1}(u_2) A_{l_2 n}(u_2) \min(u_1 - S, u_2 - S) du_1 du_2 \\ &\quad + \sum_{k_1 l_1} \Phi_{j_1 k_1}(t_1) \int_S^{t_1} \Phi_{k_1 l_1}^{-1}(u_1) A_{l_1 n}(u_1) \min(u_1 - S, t_2 - S) du, \delta_{j_1 n-1} \\ &\quad + \sum_{k_2 l_2} \Phi_{j_2 k_2}(t_2) \int_S^{t_2} \Phi_{k_2 l_2}^{-1}(u_2) A_{l_2 n}(u_2) \min(u_2 - S, t_1 - S) du_1 \delta_{j_2 n-1} \\ &\quad + \delta_{j_1 n-1} \delta_{j_2 n-1} \min(t_1 - S, t_2 - S), \end{aligned}$$

which completes the proof of the corollary. For another from of this corollary, see [10].

3. The transformation theorem. It, unfortunately, is not nearly as easy to invert (in the sense of (1.2)) the general nonlinear n^{th} order ordinary differential equation as it is the linear one. It, therefore, is necessary to use more general methods to study the Ito distribution. It is an interesting fact that inverting (2.1) is in some

way equivalent to changing variables in the Wiener integral. In this section, this idea will be made more precise. The analysis leans heavily on the work of Cameron and Martin who developed the transformation theory of the Wiener integral with their students.

THEOREM 2. *Suppose $m(t, x_{n-1}, \dots, x_1, x_0)$ is a real valued, continuous, and continuously differentiable function on R^{n+1} . Suppose also that for almost every $z \in C[S, T]$ (2.1) has a unique solution. Then, if $F(x^{(n-1)}, x^{(n-2)}, \dots, x', x)$ is a measurable and integrable functional on the Ito space defined by (1.1) and the conventions agreed upon at the beginning of § 2,*

$$\begin{aligned}
 (3.1) \quad & \int F(x^{(n-1)}, x^{(n-2)}, \dots, x', x) dP_I(x^{(n-1)}, x^{(n-2)}, \dots, x', x) \\
 &= \int F(y, T_2 y, \dots, T_n(T_{n-1}, \dots, (T_2 y))) \\
 & \quad \times \exp \left[\int_S^T p(t, T_2 y(t), \dots, T_n(T_{n-1}, \dots, (T_2 y))(t)) dt \right] \\
 & \quad \times \exp [-M(T, T_2 y(T), \dots, T_n(T_{n-1}, \dots, (T_2 y))(T))] dP_w(y) \\
 & \quad \times \exp [M(S, a_{n-1}, \dots, a_0)] .
 \end{aligned}$$

where

$$M(t, x_{n-1}, \dots, x_0) = \int_0^{x_{n-1}} m(t, u, x_{n-2}, \dots, x_0) du$$

and

$$p(t, x_{n-1}, \dots, x_0) = \frac{1}{2} m_{x_{n-1}} + M_t + \sum_{j=2}^n M_{x_{n-j}} x_{n-j} - \frac{1}{2} m^2 .$$

It will be noted that Theorem 2 allows one to express the Ito integral without solving (2.1) for almost every $z \in C[S, T]$.

Proof. From Theorem (I) it follows that

$$\begin{aligned}
 & \int F(x^{(n-1)}, x^{(n-2)}, \dots, x) dP(x^{(n-1)}, \dots, x_0) \\
 &= \int F(y, T_2 y, \dots, T_n(T_{n-1}, \dots, (T_2 y))) dP_w(z) .
 \end{aligned}$$

The result then follows immediately from [1] equation (3.6).

Note that Cameron uses a slightly different definition of the Wiener integral that usual. The result here is also generalized to include the case of arbitrary final time T and arbitrary initial values a_{n-1}, \dots, a_0 , and time S .

4. Partial differential equations. In this section, we will show that the solutions of certain partial differential equations of physics are naturally given by integrals of the form (3.1). Let

$$H(x_{n-1}, \dots, x_0; b_{n-1}, \dots, b_0) = \begin{cases} 1 & \text{if } x_{n-1} \leq b_{n-1}, \dots, x_0 \leq b_0 \\ 0 & \text{otherwise.} \end{cases} \text{ And let}$$

$$p(t, x_{n-1}, \dots, x_0) = \frac{1}{2}m_{x_{n-1}}(t, x_{n-1}, \dots, x_0) + M_t(t, x_{n-1}, \dots, x_0) + \sum M_{x_{n-j}}(t, x_{n-1}, \dots, x_0)x_{n-j} - \frac{1}{2}m^2(t, x_{n-1}, \dots, x_0).$$

We note that with the conventions agreed upon at the beginning of § 2,

$$P(b_{n-1}, \dots, b_0; a_{n-1}, \dots, a_0; t) = \text{probability } (x^{(n-1)}(t) \leq b_{n-1}, \dots, x(t) \leq b_0 \mid x^{(n-1)}(S) = a_{n-1}, \dots, x(S) = a_0)$$

is a well defined function of $2n + 2$ variables. The following theorem follows immediately from Theorem 2.

THEOREM 3. *If $m(t, x_{n-1}, \dots, x_0)$ satisfies the conditions of Theorem 2, then,*

$$\begin{aligned} &P(b_{n-1}, \dots, b_0; a_{n-1}, \dots, a_0; t) \\ &= \int H(y(t), \dots, T_n(T_{n-1}, \dots, (T_2 y))(t); b_{n-1}, \dots, b_0) \\ &\quad \times \exp \left[\int_S^T p(u, y(u), \dots, T_n(T_{n-1}, \dots, (T_2 y))(u)) du \right] \\ &\quad \times \exp [-M(T, y(T), \dots, T_n(T_{n-1}, \dots, (T_2 y)(T)))] dP_w(y) \\ &\quad \times \exp [M(S, a_{n-1}, \dots, a_0)]. \end{aligned}$$

Theorem 3 shows that the function $P(b_{n-1}, \dots, b_0; a_{n-1}, \dots, a_0; t)$ may be expressed in terms of a Wiener integral. It will presently be shown for the first order equation, anyway, that the Ito distribution, defined in § 2 is a Markov process, and thus it follows that $P(b_{n-1}, \dots, b_0; a_{n-1}, \dots, a_0; t)$ is its distribution function. It also follows that $P(b_{n-1}, \dots, b_0; a_{n-1}, \dots, a_0; t)$ solves the backward and forward diffusion equations (the Fokker-Planck equation). We state and prove this rigorously in Theorem 4, for first order equations.

THEOREM 4. *Let $\delta(a - b)$ be the delta function and let $P(b; a; S+; S)$ mean the limit as T approaches S from above. Suppose $m(T, x)$ satisfies the conditions of Theorem 2, and $m(T, x) \leq K(1 + x^2)^{1/2}$ where K is a positive constant. Suppose also that*

$$p(t, x) = \frac{1}{2}m_x + M_t - \frac{1}{2}m^2$$

is uniformly Holder continuous and bounded from below. Then it follows that the differential equations

$$(4.1) \quad \frac{\partial}{\partial S}[P(b; a; T; S)] + M(S, a)\frac{\partial}{\partial a}[P] + \frac{1}{2}\frac{\partial^2}{\partial a^2}[P] = 0$$

$$P(b; a; S^+; S) = 1 \text{ if } a \leq b \\ = 0 \text{ if } a > b$$

$$(4.2) \quad [P' \frac{\partial}{\partial T}(b; a; T; S)] + \frac{\partial}{\partial b}[M(T, b)P'] - \frac{1}{2}\frac{\partial^2}{\partial b^2}[P'] = 0$$

$$P(b; a; S^+; S) = \delta(a - b)$$

have well defined solutions. If

$$(4.3) \quad P(a; b; T; S)$$

$$= \int H(y(T); b) \exp\left[\int_S^T p(t, y(t))dt - M(T; y(T))\right] dP_w(y) \exp[M(S, a)],$$

it follows that P solves (4.1), and if

$$(4.4) \quad P'(b; a; T; S) = \int \exp\left[\int_S^T p(t, y(t))dt\right] dP_w(y | y(T) = b) \\ \times \exp[M(T, b) - M(S, a)] \\ \times (2\pi(T - S))^{-1/2} \exp[-1/2(T - S)(b - a)^2]$$

then it follows that P' solves (4.2).

$P_w(y | y(t) = b)$ is Wiener measure (probability) conditioned on $y(S) = a, y(T) = b$.

Proof. It will first be shown that

$$\frac{\partial}{\partial b}[P(b; a; T; S)] = P'(b; a; T; S) \text{ where } P' \text{ is defined by (4.4).}$$

$$P'(b; a; T; S) = \lim_{h \rightarrow 0} [P(b + h, a; T; S) - P(b; a; T; S)] \frac{1}{h} \\ = \lim_{h \rightarrow 0} \frac{1}{h} \int \exp\left[\int_S^T p(t, y(t))dt - M(T, y(T))\right] \\ \times [H(y(T); b + h) - H(y(T); b)] dP_w(y) \exp[M(S, a)].$$

$$\text{Since } \lim_{h \rightarrow 0} \frac{1}{h} \int [H(y(T); b + h) - H(y(T); b)] dP_w(y)$$

$$= \lim_{h \rightarrow 0} (2\pi(T - S))^{-1/2} \int_b^{b+h} \exp[-(1/2(T - S))(u - a)^2] du \\ = (2\pi(T - S))^{-1/2} \exp[-(1/2(T - S))(b - a)^2],$$

it follows that $\partial/\partial b[P] = P'$ as was to be shown.

From Doob [3] Chap. VI, § 3 and the assumption of this theorem, it follows that process $x(t)$, defined by (2.1) and our conventions, is a Markov process and that the following limits exist and have the indicated values.

$$(4.5) \quad \lim_{h \rightarrow 0} \frac{1}{h} \int [x(S+h) - x(S)] dP_t(x) = m(S, a)$$

$$(4.6) \quad \lim_{h \rightarrow 0} \frac{1}{h} \int [x(S+h) - x(S)]^2 dP_t(x) = 1,$$

and that

$$(4.7) \quad P_a(|x(T) - x(S)| > \varepsilon) = o(T - S). \quad (\text{any } \varepsilon > 0)$$

From [9] Theorem 1, it follows that P' has sufficient derivatives for (4.2) to make sense; from [6] and the above, it then follows that P' solves (4.2) except for (possibly) the initial conditions. Since P' has sufficient derivatives for (4.2) to make sense, it can be seen that P has sufficient derivatives for (4.1) to make sense and thus from [6] and the above, again it follows that P solves (4.1) except for (possibly) the initial conditions. The initial conditions for both (4.1) and (4.2) follow easily from (4.7). This shows Theorem 4.

The integral in (4.4) is, of course, very close to the Feynman integral of quantum mechanics; and, by way of Theorem 4, one is led to a rather interesting physical interpretation of the Schrodinger equation.

We now take a slightly different direction and note that $U(t, S, a) = E_t(y(t)) = \int y(t) dP_t(y)$ has a physical interpretation as the expected position of a particle moving in one dimension at time t which, at time S , was at a . $\partial/\partial t[U(t, S, a)]|_{t=S}$ then has a physical interpretation as the expected velocity of the particle at time S given the position of time S is a . It is interesting that this expression solves an equation similar to the one-dimensional Navier-Stokes equation. This fact will be shown in Theorem 5.

THEOREM 5¹. *Suppose $m(T, x)$ satisfies the conditions of Theorem 4 and $m_{xx}(t, x)$ exists and is continuous and let*

$$(4.9) \quad U(T, t, S, a) = E_t(y(t)) \\ = \int y(t) \exp \left[\int_S^T p(u, y(u)) du - M(T, y(T)) \right] dP_w(y) \exp [M(S, a)]$$

¹ Professor M. Donsker has shown that a ratio of Wiener integrals similar to (4.9) satisfies (4.10). Professor Donsker, however, used a different method of proof.

then

$$\lim_{t \downarrow S} 1/(t - S)[U(T, t, S, a) - U(T, S, S, a)]$$

exists, and if

$$V(T, S, a) = \lim_{t \downarrow S} 1/(t - S)[U(T, t, S, a) - U(T, S, S, a)]$$

V satisfies the equation,

$$(4.10) \quad -V_S + VV_a = -p_x(a, S) + \frac{1}{2}V_{aa}$$

with terminal condition $V(T, T, a) = m(T, a)$. The term $\exp[M(S, a)]$ may be expressed as

$$(4.11) \quad \exp[M(S, a)] = 1/\left[\exp\left[\int_S^T p(t, y(t))dt - M(T, y(T)) \right] dP_w(y) \right].$$

Proof. From (4.5), it follows immediately that the indicated limit exists and that $V(T, S, a) = m(S, a)$, and therefore $V(T, T, a) = m(T, a)$. But $M(T, a)$ is defined as

$$M(T, a) = \int_0^a m(T, u)du$$

and $p(S, a)$ is defined as

$$p(S, a) = \frac{1}{2}m_a(S, a) + M_s(S, a) - \frac{1}{2}m^2(S, a)$$

and thus

$$p_a(S, a) = \frac{1}{2}m_{aa}(S, a) + m_s(S, a) - m(S, a)m_a(S, a).$$

(4.10) thus follows immediately. (4.11) follows from the relationship

$$1 = P(\infty, a, T, S) = \exp\left[\int_S^T p(t, y(t))dt - M(T, y(T)) \right] dP_w(y) \exp[M(S, a)].$$

Theorem 5 shows that the solution to (4.10) may be expressed in terms of Wiener integrals, if it is known that the solution to (4.10) exists and has certain properties. The fact that the solution for earlier times is given in terms of later times and that the sign of V_s is minus may be fixed by reversing the direction of the Wiener paths.

More details and applications of the theory presented here will be given in a later work.

REFERENCES

1. R. H. Cameron, *Nonlinear volterra functional equations*, Journal d'Analuse Mathématique **5** (1957), 136-182.
2. ———, *Differential equations involving a parametric function*, Proc. Amer. Math. Soc. **8** (1957), 834-840.
3. J. L. Doob, *Stochastic processes*, New York, 1953.
4. E. B. Dynkin, *Markov Processes*, New York, 1965.
5. I. V. Girsanov, *On the transformation of a class of stochastic processes by means of an absolutely continuous change of measure*, Teor. Veroyatnost. i Primen **5** (1960), 314-330.
6. B. V. Gnedenko, *The Theory of Probability*, New York, 1962.
7. K. Ito, *On stochastic differential equations*, Mem. Am. Math. Soc. **4** (1951).
8. M. Kac, *On some connections between probability theory and differential and integral equations*, Proc. Second Berkeley Symposium on Math. Statistics and Probability, 189-215, University of California Press, Berkeley and Los Angeles, 1951.
9. M. Rosenblatt, *On a class of Markov processes*, Trans. Amer. Math., Soc. **71** (1951).
10. M. C. Wang and G. E. Uhlenbeck, *On the Theory of the Brownian Motion II*, Rev. of Mod. Phy. Vol. 17, Numbers 2 and 3 (1945).

Received February, 24, 1967.

AEROSPACE RESEARCH CENTER
GENERAL PRECISION SYSTEMS INC.
LITTLE FALLS, N. J.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN

Stanford University
Stanford, California

J. DUGUNDJI

Department of Mathematics
Rice University
Houston, Texas 77001

J. P. JANS

University of Washington
Seattle, Washington 98105

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California 90024.

Each author of each article receives 50 reprints free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners of publishers and have no responsibility for its content or policies.

Martin Aigner, <i>On the tetrahedral graph</i>	219
Gregory Frank Bachelis, <i>Homomorphisms of annihilator Banach algebras</i>	229
Phillip Alan Griffith, <i>Transitive and fully transitive primary abelian groups</i>	249
Benjamin Rigler Halpern, <i>Fixed points for iterates</i>	255
James Edgar Keesling, <i>Mappings and dimension in general metric spaces</i>	277
Al (Allen Frederick) Kelley, Jr., <i>Invariance for linear systems of ordinary differential equations</i>	289
Hayri Korezlioglu, <i>Reproducing kernels in separable Hilbert spaces</i>	305
Gerson Louis Levin and Wolmer Vasconcelos, <i>Homological dimensions and Macaulay rings</i>	315
Leo Sario and Mitsuru Nakai, <i>Point norms in the construction of harmonic forms</i>	325
Barbara Osofsky, <i>Noncommutative rings whose cyclic modules have cyclic injective hulls</i>	331
Newton Tenney Peck, <i>Extreme points and dimension theory</i>	341
Jack Segal, <i>Quasi dimension type. II. Types in 1-dimensional spaces</i>	353
Michael Schilder, <i>Expected values of functionals with respect to the Ito distribution</i>	371
Grigorios Tsagas, <i>A Riemannian space with strictly positive sectional curvature</i>	381
John Alexander Williamson, <i>Random walks and Riesz kernels</i>	393