

Pacific Journal of Mathematics

CURVATURE IN HILBERT GEOMETRIES. II

PAUL JOSEPH KELLY AND ERNST GABOR STRAUS

CURVATURE IN HILBERT GEOMETRIES II

PAUL KELLY AND E. G. STRAUS

The interior of a closed convex curve C in the Euclidean plane can be given a Hilbert metric, which is preserved by projective mappings. Let p, q be points interior to C and let u, v be the points of intersection of the line pq with C . The Hilbert distance $h(p, q)$ is defined by

$$h(p, q) = \left| \log \frac{d(u, p)d(v, q)}{d(v, p)d(u, q)} \right|,$$

where $d(x, y)$ denotes Euclidean distance. If C contains at most one line segment then $h(p, q)$ is a proper metric and the metric lines are the open chords of C carried by the Euclidean lines. Following Busemann [1, p. 237], we define the (qualitative) curvature at a point p as positive or negative if there exists a neighborhood U of p such that for every $x, y \in U$ we have

$$2h(\bar{x}, \bar{y}) \geq h(x, x) \quad \text{respectively} \quad 2h(\bar{x}, \bar{y}) \leq h(x, y),$$

where \bar{x}, \bar{y} are the Hilbert midpoints of p and x and of p and y respectively.

In an earlier paper [2] we proved that any point p at which the sign of the curvature is determined is a projective center of C ; that is, there exists a projective transformation which maps p into an affine center of the image of C . We also stated the conjecture that a Hilbert geometry has no point of positive curvature. It is the purpose of this paper to prove that conjecture.

Let C be centrally symmetric about the origin O . We may further assume that C has vertical lines of support at its points of intersection with the x -axis. Thus we may describe the upper arc of C by $y = y(x)$ and the lower arc by $y = -y(-x)$. Consider the points

$$a = (\varepsilon x, \varepsilon y(x)), \quad b = (\varepsilon x, -\varepsilon y(-x)), \quad 0 < \varepsilon < 1.$$

Then

$$\begin{aligned} h(O, a) &= \log \frac{\sqrt{x^2 + y(x)^2} \cdot (1 + \varepsilon) \sqrt{x^2 + y(x)^2}}{(1 - \varepsilon) \sqrt{x^2 + y(x)^2} \cdot \sqrt{x^2 + y(x)^2}} \\ (1) \quad &= \log \frac{1 + \varepsilon}{1 - \varepsilon} = 2\varepsilon + \frac{2}{3} \varepsilon^3 + O(\varepsilon^5). \end{aligned}$$

If $\bar{a} = (\lambda x, \lambda y(x))$ is the Hilbert midpoint of O and a , then according to (1) we have

$$2 \log \frac{1 + \lambda}{1 - \lambda} = \log \frac{1 + \varepsilon}{1 - \varepsilon}$$

so that

$$(2) \quad \lambda = \frac{1 - \sqrt{1 - \varepsilon^2}}{\varepsilon} = \frac{1}{2} \varepsilon + \frac{1}{8} \varepsilon^3 + O(\varepsilon^5)$$

Now let $h_1 = h(a, b)$ so that

$$\begin{aligned} h_1 &= \log \left[\frac{y(\varepsilon x) + \varepsilon y(-x)}{y(\varepsilon x) - \varepsilon y(x)} \cdot \frac{y(-\varepsilon x) + \varepsilon y(x)}{y(-\varepsilon x) - \varepsilon y(-x)} \right] \\ &= \log \frac{1 + \varepsilon \frac{y(-x)}{y(\varepsilon x)}}{1 - \varepsilon \frac{y(x)}{y(\varepsilon x)}} \cdot \frac{1 + \varepsilon \frac{y(x)}{y(-\varepsilon x)}}{1 - \varepsilon \frac{y(1-x)}{y(-\varepsilon x)}} \\ (3) \quad &= \varepsilon(y(x) + y(-x)) \left(\frac{1}{y(\varepsilon x)} + \frac{1}{y(-\varepsilon x)} \right) \\ &\quad + \frac{\varepsilon^3}{3} (y^3(x) + y^3(-x)) \left(\frac{1}{y^3(\varepsilon x)} + \frac{1}{y^3(-\varepsilon x)} \right) + O(\varepsilon^5). \end{aligned}$$

We now suppose that the curve C is twice differentiable at its points of intersection with the y -axis. We then have

$$y(\varepsilon x) = y(0) + y'(0)\varepsilon x + \frac{1}{2} y''(0)\varepsilon^2 x^2 + o(\varepsilon^2),$$

so that

$$(4) \quad \frac{1}{y(\varepsilon x)} = \frac{1}{y(0)} \left[1 - \frac{y'(0)}{y(0)} \varepsilon x + \left(\left(\frac{y'(0)}{y(0)} \right)^2 - \frac{1}{2} \frac{y''(0)}{y(0)} \right) \varepsilon^2 x^2 \right] + o(\varepsilon^2),$$

$$\frac{1}{y^3(\varepsilon x)} = \frac{1}{y^3(0)} + O(\varepsilon).$$

Substituting (4) in (3) we have

$$\begin{aligned} h_1 &= \varepsilon(y(x) + y(-x)) \frac{2}{y(0)} \left[1 + \left(\left(\frac{y'(0)}{y(0)} \right) - \frac{1}{2} \frac{y''(0)}{y(0)} \right) \varepsilon^2 x^2 \right] \\ &\quad + o(\varepsilon^3) + \frac{\varepsilon^3}{3} (y^3(x) + y^3(-x)) \frac{2}{y^3(0)} + O(\varepsilon^4) \\ (5) \quad &= \frac{2\varepsilon}{y(0)} (y(x) + y(-x)) + \frac{\varepsilon^3}{y(0)} \left[(y(x) + y(-x)) \right. \\ &\quad \left. \left(2 \left(\frac{y'(0)}{y(0)} \right)^2 - \frac{y''(0)}{y(0)} \right) + \frac{2}{3y(0)^2} (y^3(x) + y^3(-x)) \right] + o(\varepsilon^3). \end{aligned}$$

If $h_2 = h(\bar{a}, \bar{b})$, where \bar{b} is the Hilbert midpoint of O and b , then (5) holds for h_2 if we replace ε by λ . Using (2) we get

$$(6) \quad h_2 = \frac{\varepsilon}{y(0)}(y(x) + y(-x)) + \frac{\varepsilon^3}{8y(0)}(y(x) + y(-x)) \left[2 + 2x^2 \left(\frac{y'(0)}{y(0)} \right)^2 - x^2 \frac{y''(0)}{y(0)} + \frac{2}{3y^2(0)} \frac{y^3(x) + y^3(-x)}{y(x) + y(-x)} \right] + o(\varepsilon^3).$$

From (5) and (6) we have

$$(7) \quad (2h_2 - h_1) \frac{y(0)}{\varepsilon^3(y(x) + y(-x))} = \frac{1}{2} - \frac{3}{2} x^2 \left(\frac{y'(0)}{y(0)} \right)^2 + \frac{3}{4} x^2 \frac{y''(0)}{y(0)} - \frac{1}{2y^2(0)} \frac{y^3(x) + y^3(-x)}{y(x) + y(-x)} + o(1).$$

Thus the origin could be a point of positive curvature only if the right side of (7) is nonnegative for all sufficiently small values of ε , which means that the leading term is nonnegative for all x . For small x we have

$$\frac{y^3(x) + y^3(-x)}{y(x) + y(-x)} = y^2(0) \left[1 + x^2 \left(3 \left(\frac{y'(0)}{y(0)} \right)^2 + \frac{y''(0)}{y(0)} \right) \right] + o(x^2),$$

so that the leading term on the right side of (7) is

$$(8) \quad \frac{1}{4} x^2 \frac{y''(0)}{y(0)} - 3x^2 \left(\frac{y'(0)}{y(0)} \right)^2 + o(x^2).$$

For small x this term can be nonnegative only if $y'(0) = y''(0) = 0$. Thus if O is a point of positive curvature and C is twice differentiable at $(0, y(0))$ then C has zero curvature at $(0, y(0))$. But C is twice differentiable almost everywhere and any point of such differentiability can be taken as a point of intersection with the y -axis after a suitable affine transformation. Hence C must have zero curvature almost everywhere. This implies that there must be points of infinite curvature (if corner points are included in this category).

Let us start again, as before, but with $(0, y(0))$ taken as a point of infinite curvature. This means that, if $m(x)$ is the slope of a line of support of C at $(x, y(x))$, then

$$(9) \quad \lim_{\varepsilon_1, \varepsilon_2 \rightarrow 0^+} \frac{m(-\varepsilon_1) - m(\varepsilon_2)}{\min \{\varepsilon_1, \varepsilon_2\}} = \infty.$$

We can rewrite (3) as

$$(10) \quad h_1 = \varepsilon(y(x) + y(-x)) \left(\frac{1}{y(\varepsilon x)} + \frac{1}{y(-\varepsilon x)} \right) + O(\varepsilon^3)$$

and analogously

$$(11) \quad h_2 = \frac{\varepsilon}{2} (y(x) + y(-x)) \left(\frac{1}{y\left(\frac{\varepsilon}{2}x\right)} + \frac{1}{y\left(-\frac{\varepsilon}{2}x\right)} \right) + O(\varepsilon^3).$$

Thus

$$(12) \quad \begin{aligned} 2h_2 - h_1 &= \varepsilon(y(x) + y(-x)) \left[\left(\frac{1}{y\left(\frac{\varepsilon}{2}x\right)} - \frac{1}{y(\varepsilon x)} \right) \right. \\ &\quad \left. - \left(\frac{1}{y(-\varepsilon x)} - \frac{1}{y\left(-\frac{\varepsilon}{2}x\right)} \right) \right] + O(\varepsilon^3) \\ &= -2\varepsilon^2 x (y(x) + y(-x)) (M(\varepsilon_1 x) - M(-\varepsilon_2 x)) + O(\varepsilon^3), \end{aligned}$$

where $M(x)$ is the slope of a line of support to the curve $y = 1/y(x)$ at the point $(x, 1/y(x))$ and $(\varepsilon/2) < \varepsilon_1, \varepsilon_2 < \varepsilon$. Since the curve $C': y = 1/y(x)$ has infinite positive curvature at $(0, 1/(y(0)))$ our comment (9) applied to C' yields

$$+\infty = \lim_{\varepsilon_1, \varepsilon_2 \rightarrow 0} \frac{M(\varepsilon_1 x) - M(-\varepsilon_2 x)}{\min\{\varepsilon_1 x, \varepsilon_2 x\}} \leq 2 \lim_{x \rightarrow 0} \frac{M(\varepsilon_1 x) - M(-\varepsilon_2 x)}{x\varepsilon}.$$

Thus, from (12) we get

$$(13) \quad \lim_{\varepsilon \rightarrow 0} \frac{2h_2 - h_1}{\varepsilon^3} = -\infty.$$

Hence for sufficiently small ε , we have $2h_2 - h_1 < 0$ and O cannot be a point of positive curvature.

From (7) it can be seen that not every projective center is a point at which the curvature has a sign. For example, if $y'(0) = y''(0) = 0$, then the leading term on the right of (7) is positive for all x . In other words the center O is not a point of curvature if there exist a circumscribed parallelogram that touches C at the midpoints of its sides and C has zero curvature at one of the points of tangency. Even if C is twice differentiable everywhere, it does not seem easy to determine whether the complicated inequality expressed by the right side of (7) and linking three points of C is always satisfied.

REFERENCES

1. H. Busemann, *The Geometry of Geodesics*, Academic Press, New York, 1957.
2. Paul Kelly and E. G. Straus, *Curvature in Hilbert geometries*, Pacific J. Math. (1958), 119-125.

Received September 17, 1967. The second author was sponsored in part by NSF Grant GP-5497.

UNIVERSITY OF CALIFORNIA, SANTA BARBARA AND LOS ANGELES

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN

Stanford University
Stanford, California

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

J. P. JANS

University of Washington
Seattle, Washington 98105

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California 90024.

Each author of each article receives 50 reprints free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners of publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 25, No. 3

November, 1968

Philip Marshall Anselone and Theodore Windle Palmer, <i>Collectively compact sets of linear operators</i>	417
Philip Marshall Anselone and Theodore Windle Palmer, <i>Spectral analysis of collectively compact, strongly convergent operator sequences</i>	423
Edward A. Bender, <i>Characteristic polynomials of symmetric matrices</i>	433
Robert Morgan Brooks, <i>The structure space of a commutative locally convex algebra</i>	443
Jacob Feldman and Frederick Paul Greenleaf, <i>Existence of Borel transversals in groups</i>	455
Thomas Muirhead Flett, <i>Mean values of power series</i>	463
Richard Vernon Fuller, <i>Relations among continuous and various non-continuous functions</i>	495
Philip Hartman, <i>Convex sets and the bounded slope condition</i>	511
Marcel Herzog, <i>On finite groups containing a CCT-subgroup with a cyclic Sylow subgroup</i>	523
James Secord Howland, <i>On the essential spectrum of Schroedinger operators with singular potentials</i>	533
Thomas William Hungerford, <i>On the structure of principal ideal rings</i>	543
Paul Joseph Kelly and Ernst Gabor Straus, <i>Curvature in Hilbert geometries. II</i>	549
Malempati Madhusudana Rao, <i>Linear functionals on Orlicz spaces: General theory</i>	553
Stanley F. Robinson, <i>Theorems on Brewer sums</i>	587
Ralph Tyrrell Rockafellar, <i>A general correspondence between dual minimax problems and convex programs</i>	597
Richard Benjamin Sher, <i>Defining subsets of E^3 by cubes</i>	613
Howard Jacob Weiner, <i>Invariant measures and Cesàro summability</i>	621