

Pacific Journal of Mathematics

**SOME CONTINUITY PROPERTIES OF THE SCHNIRELMANN
DENSITY**

R. L. DUNCAN

SOME CONTINUITY PROPERTIES OF THE SCHNIRELMANN DENSITY

R. L. DUNCAN

Let S denote the set of all infinite increasing sequences of positive integers. For all $A = \{a_n\}$ and $B = \{b_n\}$ in S , define the metric $\rho(A, B) = 0$ if $A = B$, i.e., if $a_n = b_n$ for all n and $\rho(A, B) = 1/k$ otherwise, where k is the smallest value of n for which $a_n \neq b_n$. Similar metrics have been considered previously [1, 2].

Our purpose here is to discuss several continuity properties of the Schnirelmann density $d(A) = \inf A(n)/n$, where $A(n)$ is the number of elements of A not exceeding n . In particular, we obtain a characterization of the set of all sequences having density zero as the set of points of continuity of $d(A)$.

THEOREM 1. *$d(A)$ is upper semicontinuous on S .*

Proof. For each $\varepsilon > 0$ there is a k such that $A(k)/k < d(A) + \varepsilon$. If $\rho(A_n, A) = 1/k_n \rightarrow 0$, then $k_n \rightarrow \infty$ and there is an N such that $k_n > k$ for all $n > N$. Hence $d(A_n) \leq A_n(k)/k = A(k)/k < d(A) + \varepsilon$ for all $n > N$ and the desired result follows.

Let $L_a = \{A \in S \mid d(A) = a\}$ ($0 \leq a \leq 1$) denote the level sets of $d(A)$ and define $M_a = \{A \in S \mid d(A) \geq a\}$. Also, let \bar{L}_a denote the closure of L_a .

THEOREM 2. $\bar{L}_a = M_a$.

Proof. If $\lim \rho(A_n, A) = 0$ and $A_n \in L_a$ for all n , then $d(A) \geq \limsup d(A_n) = a$ by Theorem 1 and $\bar{L}_a \subset M_a$.

Now let $B = \{b_k\} \in M_a$ and $d(B) = b \geq a > 0$. Also, let $A = \{a_k\}$, where $a_k = 1 + [k - 1/a]$, and define $B_k = \{b_1, \dots, b_k, a_{k+1}, a_{k+2}, \dots\}$. Then $an \leq A(n) < an + 1$ and it follows that $d(A) = a$. Also, $k = B(b_k) \geq bb_k$ and $b_k \leq k/b < [k/b] + 1 \leq [k/a] + 1 = a_{k+1}$. Hence, B_k is an increasing sequence, and $B_k(n)/n \geq a$ for all n and k . Thus $d(B_k) = \lim B_k(n)/n = a$ and $B_k \in L_a$ for all k . Since $\lim \rho(B_k, B) = 0$, we have $M_a \subset \bar{L}_a$ for $a > 0$.

Finally, if $a = 0$ we define $B_k = \{b_1, \dots, b_k, b_{k+1}^2, b_{k+2}^2, \dots\}$. Then it is obvious that $B_k \in L_0$ and $\lim \rho(B_k, B) = 0$. Hence $M_0 \subset \bar{L}_0$ and $\bar{L}_a = M_a$ for $0 \leq a \leq 1$.

COROLLARY. $\bar{L}_0 = S$.

It follows from the above corollary and Theorem 1 that $d(A)$ is continuous at A if and only if $d(A) = 0$. It also follows from Theorem

2 that L_a is closed if and only if $a = 1$ and it is easily shown that L_a is never open. However, it is a consequence of the following theorem that L_a is a G_δ set and that a description of the graph of $d(A)$ can be given [3].

THEOREM 3. $d(A)$ is a function of Baire class one.

Proof. Let

$$d_n(A) = \inf_{1 \leq k \leq n} \frac{A(k)}{k}.$$

Then $d_n(A) = d_n(B)$ if $\rho(A, B) < 1/n$ and it follows that $d_n(A)$ is continuous (uniformly) on S . It remains to be shown that $\lim d_n(A) = d(A)$ for all $A \in S$.

Now let k_n be the smallest value of k for which $d_n(A) = A(k)/k$. Then $\lim d_n(A)$ exists since $d_n(A) \geq d_{n+1}(A) \geq 0$ for all n . Also, $d_n(A) = A(k_n)/k_n \geq d(A)$ for all n . Hence $\lim d_n(A) \geq d(A)$. Since the sequence $\{k_n\}$ is nondecreasing and these numbers are integers, we have either $k_n = k$ for all $n \geq N$ or $k_n \rightarrow \infty$. In the first case it is clear that $d_n(A) = d(A)$ for all $n \geq N$ and $\lim d_n(A) = d(A)$. Suppose that $k_n \rightarrow \infty$. Then $A(k)/k \geq A(k_n)/k_n = d_n(A)$ for all $k \leq k_n$. Hence $A(k)/k \geq \lim d_n(A)$ for all k and $\lim d_n(A) \leq d(A)$ since the sequence $\{d_n(A)\}$ is nonincreasing. Thus $\lim d_n(A) = d(A)$ in this case also.

REFERENCES

1. R. L. Duncan, *A topology for sequences of integers I*, Amer. Math. Monthly **66** (1959), 34-39.
2. ———, *A topology for sequences of integers II*, Amer. Math. Monthly **67** (1960), 537-539.
3. E. S. Thomas, Jr., *Some characterizations of functions of Baire class 1*, Proc. Amer. Math. Soc. **17** (1966), 456-461.

Received July 18, 1967.

LOCK HAVEN STATE COLLEGE AND
PENNSYLVANIA STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN

Stanford University
Stanford, California

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS

University of Washington
Seattle, Washington 98105

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Pacific Journal of Mathematics

Vol. 26, No. 1

November, 1968

Efraim Pacillas Armendariz, <i>Closure properties in radical theory</i>	1
Friedrich-Wilhelm Bauer, <i>Postnikov-decompositions of functors</i>	9
Thomas Ru-Wen Chow, <i>The equivalence of group invariant positive definite functions</i>	25
Thomas Allan Cootz, <i>A maximum principle and geometric properties of level sets</i>	39
Rodolfo DeSapio, <i>Almost diffeomorphisms of manifolds</i>	47
R. L. Duncan, <i>Some continuity properties of the Schnirelmann density</i>	57
Ralph Jasper Faudree, Jr., <i>Automorphism groups of finite subgroups of division rings</i>	59
Thomas Alastair Gillespie, <i>An invariant subspace theorem of J. Feldman</i>	67
George Isaac Glauberman and John Griggs Thompson, <i>Weakly closed direct factors of Sylow subgroups</i>	73
Hiroshi Haruki, <i>On inequalities generalizing a Pythagorean functional equation and Jensen's functional equation</i>	85
David Wilson Henderson, <i>D-dimension. I. A new transfinite dimension</i>	91
David Wilson Henderson, <i>D-dimension. II. Separable spaces and compactifications</i>	109
Julien O. Hennefeld, <i>A note on the Arens products</i>	115
Richard Vincent Kadison, <i>Strong continuity of operator functions</i>	121
J. G. Kalbfleisch and Ralph Gordon Stanton, <i>Maximal and minimal coverings of $(k - 1)$-tuples by k-tuples</i>	131
Franklin Lowenthal, <i>On generating subgroups of the Moebius group by pairs of infinitesimal transformations</i>	141
Michael Barry Marcus, <i>Gaussian processes with stationary increments possessing discontinuous sample paths</i>	149
Zalman Rubinstein, <i>On a problem of Ilyeff</i>	159
Bernard Russo, <i>Unimodular contractions in Hilbert space</i>	163
David Lee Skoug, <i>Generalized Ilstow and Feynman integrals</i>	171
William Charles Waterhouse, <i>Dual groups of vector spaces</i>	193