

Pacific Journal of Mathematics

GENERALIZED ILSTOW AND FEYNMAN INTEGRALS

DAVID LEE SKOUG

GENERALIZED ILSTOW AND FEYNMAN INTEGRALS

D. L. SKOUG

Let $C[a, b]$ denote the space of continuous functions $x(t)$ defined on $[a, b] \ni x(a) = 0$. This space is called Wiener space. Using the Wiener integral we define, for each nonnegative integer M , what we call the M Ilstow, M complex Wiener, M Feynman, limiting M complex Wiener, and limiting M Feynman integrals of a functional $F(x)$ on $C[a, b]$ and show various relationships which exist between these integrals. In particular we give necessary and sufficient conditions for a finite dimensional functional $F(x)$ to be M Ilstow integrable on $C[a, b]$.

We consider the set of linear functionals $x(t_1), \dots, x(t_n)$ where $a = t_0 < t_1 < \dots < t_n = b$ and obtain conditions on $g_j(u) \ni$ the functional

$$(1.1) \quad F(x) = g_1[x(t_1)] \cdots g_n[x(t_n)]$$

is M Ilstow and limiting M Feynman integrable on $C[a, b]$. We then apply these results to the functional

$$F(t, \xi, x) = \exp\left(\int_a^b \theta[t - s, x(s) + \xi] ds\right) \sigma[x(t) + \xi]$$

where $0 \leq t \leq t_0$, $-\infty < \xi < \infty$ and $x \in C[0, t_0]$ and show that for appropriate functions $\theta(t, \xi)$ and $\sigma(\xi)$, the limiting M Feynman integral $\hat{G}(t, \xi, q)$ of $F(t, \xi, x)$ exists for

$$(t, \xi, q) \in (0, t_0) \otimes R_1 \otimes \{R_1 - \{0\}\}$$

and satisfies there the integral equation

$$(1.2) \quad \begin{aligned} \hat{G}(t, \xi, q) &= \left(\frac{-iq}{2\pi t}\right)^{1/2} \int_{-\infty}^{\infty} \sigma(u) \exp\left(\frac{qi(\xi - u)^2}{2t}\right) du \\ &+ \left(\frac{-iq}{2\pi}\right)^{1/2} \int_0^t (t-s)^{-1/2} ds \\ &\times \int_{-\infty}^{\infty} \theta(s, u) \hat{G}(s, u, q) \exp\left(\frac{qi(\xi - u)^2}{2(t-s)}\right) du. \end{aligned}$$

For $M = 0$ the definitions of the above mentioned integrals reduce to the definitions of the Ilstow, complex Wiener, Feynman, limiting complex Wiener, and limiting Feynman integrals as defined by R. H. Cameron in [2]. He used the Ilstow integral as an intermediate integral in his definition of the Feynman integral. The word "Ilstow" is a contraction of "inverse Laplace Stieltjes transform of Wiener's".

Many of the theorems in this paper are generalizations of theorems in [2]. However, the techniques developed in § 4, applied when $M = 0$, allow us to reduce the hypothesis of several theorems of [2]. In particular, we obtain a condition for the Ilstow and limiting Feynman

integrability of $F(x)$ defined by (1.1), that only requires each g'_i to be essentially of bounded variation on any bounded interval, instead of each g''_i being continuous as required by Theorem 7.2 of [2].

These more general integrals allow us to weaken the conditions required on σ and still obtain the existence of $\hat{G}(t, \xi, q)$.

2. Generalized Ilstow and complex Wiener integrals and the banach space $B(M, \lambda)$.

DEFINITION 2.1. Let M be a nonnegative integer. Let $F(\rho x)$ be Lebesgue \otimes Wiener integrable on $[0, \rho_0] \otimes C[a, b]$ and assume $\exists f_M(s)$ on $[0, \infty) \ni f_M(0) = 0$, f_M is left continuous and of bounded variation on every bounded interval $[0, s_0]$, $\int_0^\infty e^{-\lambda s} |df_M(s)| < \infty$ for some $\lambda > 0$, and for almost all sufficiently large λ ,

$$\int_{C[a, b]} F(\lambda^{-\frac{1}{2}}x) dx = \lambda^{M/2} \int_0^\infty e^{-\lambda s} df_M(s) .$$

Then $F(x)$ will be said to be M Ilstow integrable on $C[a, b]$ and $f_M(s)$ is called the M Ilstow integral of $F(x)$ with parameter s and we write $f_M(s) = \int_{C[a, b]}^{ILM} F(x) dx$.

Notation. Let $I_M = \{F(x) \mid F(x) \text{ is } M \text{ Ilstow integrable on } C[a, b]\}$.

We use the next lemma in the proof of several theorems.

LEMMA 2.1. Suppose f is of bounded variation on every bounded interval of $[0, \infty)$ such that for some $\lambda_0 \geq 0$, $\int_0^\infty e^{-\lambda_0 s} |df(s)| < \infty$. Then for $\lambda > \lambda_0$

$$(2.1) \quad \int_0^\infty e^{-\lambda s} df(s) = \lambda^{\frac{1}{2}} \int_0^\infty e^{-\lambda s} dg(s)$$

where

$$(2.2) \quad g(s) = 2\pi^{-\frac{1}{2}} \int_0^s (s - \rho)^{1/2} df(\rho) .$$

In addition, g is absolutely continuous on every bounded interval $[0, s_0]$ and $g(0) = 0$. Finally for $\lambda > \lambda_0$

$$(2.3) \quad \int_0^\infty e^{-\lambda s} |dg(s)| \leq \lambda^{-\frac{1}{2}} \int_0^\infty e^{-\lambda s} |df(s)| .$$

Proof. Same as proof of Lemma 5 on pp. 346–347 of [2].

THEOREM 2.1. $I_M \subseteq I_{M+1}$ for $M = 0, 1, 2, \dots$. Furthermore if $F(x) \in I_M$ then

$$(2.4) \quad f_{M+1}(s) = 2\pi^{-\frac{1}{2}} \int_0^s (s - \rho)^{\frac{1}{2}} df_M(\rho)$$

and

$$(2.5) \quad f_{M+2}(s) = 2\pi^{-\frac{1}{2}} \int_0^s (s - \rho)^{\frac{1}{2}} df_{M+1}(\rho) = \int_0^s f_M(\rho) d\rho$$

so that the $(M + 2)$ Ilstow integral of $F(x)$ is differentiable a.e. on $[0, \infty)$ and its derivative is $f_M(s)$.

Proof. Assume $F(x) \in I_M$ with M Ilstow integral $f_M(s)$. Define $f_{M+1}(s)$ by equation (2.4) and note that (2.1)–(2.3) imply $F(x) \in I_{M+1}$. To obtain the second equality of (2.5), integrate the middle expression by parts, substitute for f_{M+1} from (2.4) and interchange order of integration.

DEFINITION 2.2. Let $\lambda \neq 0$ be $\ni \operatorname{Re} \lambda \geq 0$. Let $F(x) \in I_M$ with M Ilstow integral $f_M(s)$. Then if

$$(2.6) \quad \int_0^\infty |e^{-\lambda s}| |df_M(s)| < \infty$$

we say that $F(x)$ is M complex Wiener integrable on $C[a, b]$ with revar parameter λ (λ is called the reciprocal variance or revar parameter since in [1], λ^{-1} denoted the variance parameter) and we define the value of the M complex Wiener integral to be

$$\int_{C[a, b]}^{W_\lambda^M} F(x) dx = \lambda^{M/2} \int_0^\infty e^{-\lambda s} df_M(s)$$

where if M is odd the square roots have values in the right half plane. If $\operatorname{Re} \lambda = 0$ the M complex Wiener integral will be called an M Feynman integral. For $\lambda = -iq$, q real, we denote the M Feynman integral of $F(x)$ by $\int_{C[a, b]}^{F_q^M} F(x) dx$.

Note that a functional $F(x)$ is M Feynman integrable if and only if its M Ilstow integral is of bounded variation on $[0, \infty)$. In order to remove this somewhat restrictive condition we shall generalize the above definition slightly.

DEFINITION 2.3. If in Definition 2.2 we replace (2.6) by

$$(2.7) \quad \int_0^\infty |e^{-s(\lambda + \gamma)}| |df_M(s)| < \infty$$

for all $\eta \ni \operatorname{Re} \eta > 0$ and if the right member of

$$\int_{C[a,b]}^{\rightarrow W_\lambda^M} F(x) dx = \lambda^{M/2} \lim_{\substack{\eta \rightarrow 0 \\ (\operatorname{Re} \eta > 0)}} \int_0^\infty e^{-s(\lambda+\eta)} df_M(s)$$

exists, we say that $F(x)$ has a limiting M complex Wiener integral with revar parameter λ and we give it the value and notation specified above.

Notation. The symbol $A \rightleftharpoons B$ shall mean the existence of A implies the existence of B and the equality $A = B$.

In terms of the analytic Wiener integral, defined by Cameron on p. 289 of [2], and denoted by $\int_{C[a,b]}^{anW_\lambda} F(x) dx$, we have the following theorem.

THEOREM 2.2. For $\operatorname{Re} \lambda \geq 0$

$$\int_{C[a,b]}^{W_\lambda^M} F(x) dx \rightleftharpoons \int_{C[a,b]}^{\rightarrow W_\lambda^M} F(x) dx \rightleftharpoons \int_{C[a,b]}^{anW_\lambda} F(x) dx$$

and if $F \in I_M$ then for almost all sufficiently large positive λ we have

$$\int_{C[a,b]} F(\lambda^{-\frac{1}{2}}x) dx = \int_{C[a,b]}^{W_\lambda^M} F(x) dx.$$

Proof. Same as proof of Theorem 1 of [2] as the additional factor $\lambda^{M/2}$ doesn't affect analyticity.

Notation. Let $W(M, \lambda) = \{F(x) \mid F(x) \text{ is } M \text{ complex Wiener integrable on } C[a, b] \text{ with revar parameter } \lambda\}$.

$\overrightarrow{W}(M, \lambda) = \{F(x) \mid F(x) \text{ is limiting } M \text{ complex Wiener integrable on } C[a, b] \text{ with revar parameter } \lambda\}$.

The following theorem follows directly from Lemma 2.1 and Theorems 2.1 and 2.2.

THEOREM 2.3. $W(M, \lambda) \subseteq W(M+1, \lambda)$ for $\operatorname{Re} \lambda > 0$ and $M = 0, 1, 2, \dots$. $\overrightarrow{W}(M, \lambda) \subseteq \overrightarrow{W}(M+1, \lambda)$ for $\operatorname{Re} \lambda \geq 0$ and $M = 0, 1, 2, \dots$. Furthermore if $F(x) \in W(M, \lambda)$ then the M and $(M+1)$ complex Wiener integrals of $F(x)$ are equal (both being equal to the analytic Wiener integral of $F(x)$ by Theorem 2.2). A similar statement holds in case $F(x) \in \overrightarrow{W}(M, \lambda)$.

We now define for $M = 0, 1, 2, \dots$ and $\lambda \geq 0$, a space $B(M, \lambda)$ of functionals of $C[a, b]$ and discuss some of its properties. In particular $B(0, \lambda) = B_\lambda$ where B_λ was defined on p. 297 of [2].

DEFINITION 2.4. Let $\lambda \geq 0$ and $M \in \{0, 1, 2, \dots\}$ be given. Let $F(x)$ be a functional defined on $C[a, b] \ni F(\rho x)$ is Lebesgue \otimes Wiener measurable in (ρ, x) on $[0, \lambda^{-\frac{1}{2}}] \otimes C[a, b]$ and \ni

$$(2.8) \quad N_\lambda(F) = \int_\lambda^\infty e^{\lambda-\rho} d\rho \int_{C[a, b]} |F(\rho^{-\frac{1}{2}}x)| dx < \infty$$

where we interpret $\lambda^{-\frac{1}{2}}$ to be $+\infty$ if $\lambda = 0$. Assume further that there exists a left continuous function $f(s)$ of bounded variation on every bounded subinterval of $[0, \infty) \ni$

$$(2.9) \quad N'_\lambda(M, F) = \int_0^\infty e^{-\lambda s} |df(s)| < \infty$$

and such that for almost all $\rho > \lambda$

$$(2.10) \quad \rho^{M/2} \int_0^\infty e^{-\rho s} df(s) = \int_{C[a, b]} F(\rho^{-\frac{1}{2}}x) dx.$$

Then we say $F \in \tilde{B}(M, \lambda)$ and define the norm of F to be $N_\lambda(M, F) = N_\lambda(F) + N'_\lambda(M, F)$. Note that $N_\lambda = N'_\lambda$ where N'_λ is defined by (2.1) of [2] while $N'_\lambda(M, \cdot) = N''_\lambda(\cdot)$ if and only if $M = 0$ where N''_λ is defined by (2.2) of [2].

DEFINITION 2.5. We say that two elements F_1 and F_2 of $\tilde{B}(M, \lambda)$ are equivalent if for almost all (ρ, x) on $[0, \lambda^{-\frac{1}{2}}] \otimes C[a, b]$ we have $F_1(\rho x) = F_2(\rho x)$. We define $B(M, \lambda)$ as the space of equivalence classes of elements of $\tilde{B}(M, \lambda)$.

THEOREM 2.4. The space $B(M, \lambda)$ has the following properties:

- (a) $B(M, \lambda) \subseteq I_M$ for $\lambda \geq 0$ and $M = 0, 1, 2, \dots$.
- (b) $B(M, \lambda)$ is a Banach space with norm $N_\lambda(M, \cdot)$.
- (c) $B(M, \lambda) \subseteq B(M+1, \lambda)$ for $\lambda > 0$, and $M = 0, 1, 2, \dots$.
- (d) $0 \leq \lambda_1 \leq \lambda_2 \Rightarrow B(M, \lambda_1) \subseteq B(M, \lambda_2)$.

Proof. Property (a) follows from the definitions of $B(M, \lambda)$ and I_M . Property (b) follows from Theorem 2 of [2]. Property (c) follows from equations (2.3), (2.4), (2.9) and (2.10). Property (d) follows as $0 \leq \lambda_1 \leq \lambda_2$ implies that $N_{\lambda_2}(M, F) \leq N_{\lambda_1}(M, F)$.

3. Finite dimensional functionals. Let $\alpha_1(t), \dots, \alpha_n(t)$ be real functions of bounded variation on $[a, b]$ and let $g(u_1, \dots, u_n)$ be a real or complex function defined almost everywhere on R_n . Let

$$(3.1) \quad F(x) = g \left[\int_a^b \alpha_1(t) dx(t), \dots, \int_a^b \alpha_n(t) dx(t) \right].$$

Then $F(x)$ will be called an " n -dimensional functional" on $C[a, b]$. If $g(u_1, \dots, u_n)$ is Lebesgue measurable on R_n we call F a measurable n -dimensional functional. If $\alpha_1(t), \dots, \alpha_n(t)$ are an orthonormal set on $[a, b]$, (3.1) will be said to express $F(x)$ in "normalized form".

Our next theorem is a generalization of Theorem 3 of [2] and gives necessary and sufficient conditions that a finite dimensional functional be M Ilstow integrable and a formula for the M Ilstow integral.

THEOREM 3. *Let $F(x)$ be a measurable n -dimensional functional on $C[a, b]$ expressed in normalized form by (3.1). Let $M \in \{0, 1, 2, \dots\}$. Let $K = 0$ if $n \leq M$ while if $n > M$ let K equal $(n - M)/2$ or $(n + 1 - M)/2$ whichever is an integer. Then $F(x)$ is M Ilstow integrable on $C[a, b]$ if and only if the following four conditions are satisfied:*

(1) *For sufficiently large λ , the quantity $N_\lambda(F)$ is finite, where*

$$N_\lambda(F) = (2\pi)^{-n/2} \int_\lambda^\infty e^{\lambda - \rho} d\rho \int_{R_n} |g(\rho^{-1/2}u_1, \dots, \rho^{-1/2}u_n)| \exp\left(-\frac{(u_1^2 + \dots + u_n^2)}{2}\right) du.$$

(2) *The function $I_n(M, s)$ defined by*

$$(3.2) \quad I_n(M, s) = \int_{u_1^2 + \dots + u_{M+2k}^2 < 2s}^{(M+2k)} g(u_1, \dots, u_n) du_1 \dots du_{M+2k}$$

has $(K - 1)$ continuous derivatives all of which of course vanish for $s \leq 0$.

(3) *The $(K - 1)$ th derivative $I_n^{(K-1)}(M, s)$ has a left hand derivative, which we denote by $I_n^{(K)}(M, s)$, which is of bounded variation on every bounded interval and which is a true K th derivative of $I_n(M, s)$ except on the countable set where it has jumps.*

(4) *For sufficiently large λ , the quantity $N'_\lambda(F, M)$ is finite, where*

$$N'_\lambda(F, M) = \left(\frac{1}{2\pi}\right)^{(2k+M)/2} \int_0^\infty e^{-\lambda s} |d_s I_n^{(K)}(M, s)|.$$

Moreover when conditions (1) through (4) are satisfied we have for $s \geq 0$

$$\int_{C[a, b]}^{IL_{W^M_s}} F(x) dx = \left(\frac{1}{2\pi}\right)^{(2k+M)/2} I_n^{(K)}(M, s)$$

Note that when $K = 0$, conditions (2) and (3) reduce to $I_n(M, s)$ being of bounded variation on every bounded interval.

COROLLARY TO THEOREM 3. *For*

$$\begin{aligned} M = 1, 2, \dots, I_{M-1} \subsetneq I_M, B(M-1, 0) \\ \not\subseteq B(M, 0), B(M, 0) \not\subseteq B(M-1, 0), W(M-1, 0) \\ \not\subseteq W(M, 0) \quad \text{and} \quad W(M, 0) \not\subseteq W(M-1, 0). \end{aligned}$$

Proof. Let $g(u_1, \dots, u_j)$ be the characteristic function of a j -dimensional sphere of radius $\sqrt{2}$ about the origin. Let $\alpha_1(t), \dots, \alpha_j(t)$ be an orthonormal set on $[a, b]$. Let

$$F_j(x) = g\left[\int_a^b \alpha_1(t) dx(t), \dots, \int_a^b \alpha_j(t) dx(t)\right].$$

Then for $M = 1, 2, \dots$, $F_{M+2}(x) \in I_M$ and $F_{M+2}(x) \notin I_{M-1}$. This follows by noting that by use of (2.4) and (3.2) we obtain for $s \geq 0$

$$I_{M+2}(M, s) = \frac{(2\pi)^{(M+2)/2}}{\Gamma\left(\frac{M+4}{2}\right)} \min\{1, s^{(M+2)/2}\}$$

so that by Theorem 3, $F_{M+2}(x) \in I_M$ while

$$\begin{aligned} I_{M+2}(M-1, s) &= I_{M+2}(M+1, s) \\ &= \begin{cases} \frac{2(2\pi s)^{(M+3)/2}}{\Gamma\left(\frac{M+5}{2}\right)}, & 0 \leq s \leq 1 \\ \frac{2\sqrt{2}(2\pi)^{(M+2)/2}}{\Gamma\left(\frac{M+2}{2}\right)} \int_0^1 (s-\rho)^{1/2} \rho^{M/2} d\rho, & s > 1 \end{cases} \end{aligned}$$

so that $\lim_{s \rightarrow 1+} I_{M+2}^{(2)}(M-1, s) = -\infty$ and hence $F(x) \notin I_{M-1}$.

Similarly for $M = 1, 2, \dots$, one can show that

$$F_M(x) \in B(M, 0) \cap W(M, 0) \quad \text{and} \quad F_M(x) \notin B(M+1, 0) \cup W(M+1, 0).$$

For $M = 3, 4, \dots$ it follows that $F_M(x) \in B(M-2, 0) \cap W(M-2, 0)$ while $F_M(x) \notin B(M-3, 0) \cup W(M-3, 0)$. Finally letting $g(u_1, u_2) \equiv 1$ and writing $F(x)$ in form (3.1) one obtains $F(x) \in B(0, 0) \cap W(0, 0)$ and $F(x) \notin B(1, 0) \cup W(1, 0)$.

4. **Products of functions of (xt_j) .** In this section we consider the set of linear functionals, $x(t_1), \dots, x(t_n)$, and obtain conditions under which products of the form

$$(4.1) \quad F(x) = g_1[x(t_1)] \cdots g_n[x(t_n)]$$

are M Ilstow and limiting M Feynman integrable on $C[a, b]$. $F(x)$ as given by (4.1) is an n -dimensional functional but here we wish to obtain simpler conditions for M Ilstow integrability than those given in Theorem 3. In Theorem 7.1 of [2], R. H. Cameron showed that each g'_i being continuous and g_i and g'_i both being absolutely integrable on R was sufficient to insure Ilstow integrability. Here we will only require each of the g_i 's to have a derivative essentially of bounded variation on any bounded interval, and satisfy certain growth conditions.

In Theorem 4 we will find it necessary to express certain Lebesgue-Lebesgue Stieltjes iterated integrals as Laplace Stieltjes transforms. We shall now develop the required notation and lemmas to achieve this objective.

LEMMA 4.1. *Assume $f(u)$ is absolutely continuous on $[c, d]$ and has a derivative $f'(u)$ which is essentially of bounded variation on $[c, d]$. Let $\phi(u) \equiv \text{ess}_{h \rightarrow 0+} \lim f'(u + h)$ and assume $\text{Var}(\phi, [c, d]) < \infty$. Let $a \in [c, d]$ and let $g(u) = \{f(u) - f(a)\}/(u - a)$ for $u \in [c, d]$, $u \neq a$ and let $g(a) = \phi(a)$. Then $g(u)$ is of bounded variation on $[c, d]$ and $\text{Var}(g, [c, d]) \leq \text{Var}(\phi, [c, d])$.*

Proof. This lemma follows from the observation that for all $u \in [c, d]$ we have $g(u) = \int_0^1 \phi[(u - a)t + a] dt$.

LEMMA 4.2. *Assume $f_1(u), \dots, f_n(u)$ are of bounded variation on $[c, d]$. Let $1 \leq k \leq n$ and assume that $g_1(u)$ and $g_2(u)$ are such that for all $u \in [c, d]$ we have $|f_i(u)| \leq g_1(u)$ for $i = 1, 2, \dots, k$, while $|f_i(u)| \leq g_2(u)$ for $i = k + 1, \dots, n$. Let $K_1 = \max_{u \in [c, d]} g_1(u)$ and $K_2 = \max_{u \in [c, d]} g_2(u)$. Then*

$$\text{Var}\left(\prod_{i=1}^n f_i, [c, d]\right) \leq K_1^{k-1} K_2^{n-k-1} \left[K_2 \sum_1^k \text{Var}(f_i, [c, d]) + K_2 \sum_{k+1}^n \text{Var}(f_i, [c, d]) \right].$$

Notation. Let m be a positive odd integer. For $i = 1, 2, \dots, m$, let k_i be a positive constant and let $\phi_i(s) = (s^2)/(k_i)$. Let $u_0 \equiv 0$, $\psi_1(u) = \sum_{i=1}^m \phi_i(u_i - u_{i-1})$, and $\psi_2(u) = u_1^2 + \phi_2(u_2) + \sum_{i=3}^m \phi_i(u_i - u_{i-1})$. For $\sigma \geq 0$ and $j = 1, 2$, let $S_j(\sigma) = \{u \mid \psi_j(u) < \sigma\}$. Note that $S_j(\sigma)$ is an open m -dimensional ellipsoid with its center at the origin.

LEMMA 4.3. *For $i = 1, 3, \dots, m$, assume $g_i(u_i)$ is bounded and Lebesgue integrable on every bounded interval of R_1 . For $i = 2, 4, \dots, m - 1$, assume $h_i(u_{i-1}, u_i, u_{i+1})$ is Borel measurable in its three vari-*

ables and is right continuous and of bounded variation with respect to u_i on every bounded interval $[-2d, 2d]$ for any $u_{i-1}, u_{i+1} \in [-d, d]$. For notational purposes we will denote the function $h_i(u_{i+1}, u_i, u_{i-1})$ by $h_i(u_i)$. Also assume that $\text{Var}(h_i(s), -2d \leq s \leq 2d)$ is bounded on $u_{i-1}, u_{i+1} \in [-d, d]$. Then for $j = 1, 2$, $A_j(\sigma)$ exists and is of bounded variation on $0 \leq \sigma \leq \nu$ for any $\nu > 0$ where $A_j(\sigma)$ is the iterated Lebesgue-Lebesgue Stieltjes integral

$$A_j(\sigma) = \int_{-\infty}^{\infty} g_1(u_1) du_1 \cdots \int_{-\infty}^{\infty} g_m(u_m) du_m \int_{-\infty}^{\infty} d_{u_2} h_2(u_2) \int_{-\infty}^{\infty} d_{u_4} h_4(u_4) \\ \cdots \int_{-\infty}^{\infty} \chi_{s_j(\sigma)}(u_1, u_2, \cdots, u_m) d_{u_{m-1}} h_{m-1}(u_{m-1})$$

which for notational purposes we shall write as

$$A_j(\sigma) = \int_{0 \leq \psi_j(u) < \sigma}^{(m)} \cdots \int \prod_{k=1}^{(m-1)/2} g_{2k-1}(u_{2k-1}) \prod_{k=1}^{(m+1)/2} d_{u_{2k}} h_{2k}(u_{2k}) du_m \cdots du_3 du_1.$$

Further if $f(\sigma)$ is continuous on $[0, \nu]$ then

$$(4.2) \quad \int_0^\nu f(\sigma) dA_j(\sigma) = \int_{0 \leq \psi_j(u) < \sigma}^{(m)} \cdots \int f(\psi_j(u)) \left(\prod_{k=1}^{(m+1)/2} g_{2k-1}(u_{2k-1}) \right) \\ \times \left(\prod_{k=1}^{(m-1)/2} d_{u_{2k}} h_{2k}(u_{2k}) \right) du_m du_{m-2} \cdots du_3 du_1.$$

In addition if $f(\sigma)$ is continuous on $[0, \infty)$ then the existence (as a finite number) and absolute convergence of the right side of the following equation implies the existence of the left and equality.

$$(4.3) \quad \lim_{C \rightarrow \infty} \int_0^C f(\sigma) dA_j(\sigma) = \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} f(\psi_j(u)) \left(\prod_{k=1}^{(m+1)/2} g_{2k-1}(u_{2k-1}) \right) \\ \times \left(\prod_{k=1}^{(m-1)/2} d_{u_{2k}} h_{2k}(u_{2k}) \right) du_m \cdots du_3 du_1.$$

Proof. As $S_j(\sigma)$ is an open set in R_m one can show that

$$\int_{-\infty}^{\infty} d_{u_2} h_2(u_2) \int_{-\infty}^{\infty} d_{u_4} h_4(u_4) \cdots \int_{-\infty}^{\infty} \chi_{s_j(\sigma)}(u_1, u_2, \cdots, u_m) d_{u_{m-1}} h_{m-1}(u_{m-1})$$

is bounded and Borel measurable in its $(m+1)/2$ variables u_1, u_3, \cdots, u_m . Hence $A_j(\sigma)$ exists for $0 \leq \sigma \leq \nu$. By breaking up the g_i 's and h_i 's appropriately one can write $A_j(\sigma)$ as the difference of two monotonic increasing functions and so $A_j(\sigma)$ is of bounded variation on every bounded interval. Equation (4.2) follows from the general limiting sum type of argument. As $\nu \rightarrow \infty$, $S_j(\sigma) \rightarrow R_m$ and so by dominated convergence in each variable we obtain (4.3).

In the following lemma and theorem we will use the notation

developed by Cameron in Theorem 7.1 of [2].

LEMMA 4.4. Assume $h(u)$ is absolutely continuous on every bounded interval of R_1 and $\phi(u) \equiv \text{ess}_{h \rightarrow 0+} \lim h'(u + h)$ is of bounded variation on every bounded interval of R_1 . Also assume that for some $A > 0$, $B \geq 0$ and $0 \leq \gamma < 2$ we have $|h(u)| \leq Ae^{B|u|^\gamma}$ and $\int_{-V}^V |d\phi(s)| \leq Ae^{BV^\gamma}$ for all $u \in R_1$ and $V > 0$. Then for $0 \leq i < j < k$, $\lambda > 0$ and $u_{i,k}^j \in R_1$ we have

$$\begin{aligned}
 \int_{-\infty}^{\infty} h(u_j) E_j^i E_k^j du_j &= \lambda^{-1/2} \Gamma_{i,k}^j [h(u_j)] + \int_{-\infty}^{\infty} E_j^i E_k^j \hat{\Gamma}_{i,k}^j h(u_j) du_j \\
 (4.4) \qquad \qquad \qquad &= \lambda^{-1/2} \Gamma_{i,k}^j [h(u_j)] + \lambda^{-1} \alpha_{i,k}^j (t_j - t_i) \\
 &\quad \times \int_{-\infty}^{\infty} E_j^i E_k^j d_{u_j} \left[\frac{h(u_j) - h(u_{i,k}^j)}{u_j - u_{i,k}^j} \right]
 \end{aligned}$$

where the integral in the last term of (4.4) exists both as a Riemann Stieltjes integral and as a Lebesgue Stieltjes integral and where $\hat{\Gamma}_{i,k}^j h(u_j) = h(u_j) - h(u_{i,k}^j)$.

Proof. The first equality follows directly by noting that

$$\lambda^{-\frac{1}{2}} E_k^i = \int_{-\infty}^{\infty} E_j^i E_k^j du_j$$

while the second equality follows from integration by parts which is justified by the above growth conditions.

THEOREM 4. Let $a = t_0 < t_1 < \dots < t_n = b$. For $i = 1, 2, \dots, n-1$ assume $g_i(u)$ is absolutely continuous on every bounded subinterval of R_1 and $\phi_i(u) \equiv \text{ess}_{h \rightarrow 0+} \lim g_i'(u + h)$ is of bounded variation on every bounded subinterval of R_1 . Furthermore assume for some $0 \leq \gamma < 2$ and $B > 0$

$$(4.5) \qquad |g_i(u)| \leq B$$

$$(4.6) \qquad \int_{-\infty}^{\infty} |g_i(s)| ds \leq B$$

$$(4.7) \qquad \int_{-V}^V |d\phi_i(s)| \leq Be^{BV^\gamma}$$

$$(4.8) \qquad \int_{-V}^V |dg_i(s)| \leq Be^{BV^\gamma}$$

$$(4.9) \qquad |\phi_i(u)| \leq Be^{B|u|^\gamma}$$

for $i = 1, 2, \dots, n-1$, $u \in R_1$ and $V > 0$. Furthermore assume

$$\int_{-\infty}^{\infty} |g_n(s)| ds \leq B.$$

Let $F(x)$ be given as in (4.1). Then there exists a function $\psi(\lambda)$, analytic for $\operatorname{Re} \lambda > 0$, continuous for $\operatorname{Re} \lambda \geq 0$, and which for real $\lambda > 0$ satisfies

$$\psi(\lambda) = \int_{C[a,b]} F(\lambda^{-\frac{1}{2}}x) dx.$$

Furthermore $F(x)$ is M Ilstow and limiting M Feynman integrable for all values of the parameter for $M \geq 1$ and $F(x) \in B(1, \lambda)$ for all $\lambda > 0$.

Proof. By using techniques similar to those used by Cameron in proving Theorem 7.1 of [2] (in particular using (4.4) in place of (7.8) of [2]) we obtain for $\lambda > 0$

$$\begin{aligned}
 \lambda^{-1/2} \int_{C[a,b]} F(\lambda^{-1/2}x) dx &= \int_{-\infty}^{\infty} \frac{g_n(u_n) g_{0,n}^{\#}(0, u_n)}{\sqrt{2\pi(b-a)}} \exp\left(-\frac{\lambda u_n^2}{2(b-a)}\right) du_n \\
 &+ \sum_{\substack{2 \\ \cup \\ j=1}}^{Q_j(n-1)} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \frac{h^{\#}(\rho_1) g_n(\rho_{q+1})}{\sqrt{2\pi(T'_{q+1} - T_q)}} \\
 &\times \prod_{i=1}^q \left(\frac{g_{v_i}(\rho_i) \alpha_{v_i, v_{i+1}}^{v_i+1} (T_i - T'_i)^{1/2}}{2\pi(T'_i - T_{i-1})^{1/2}} \right) \\
 &\times \exp\left\{ -\frac{\lambda}{2} \left(\sum_{i=1}^{q+1} \frac{(\rho_i - \sigma_{i-1})^2}{T'_i - T_{i-1}} + \sum_{i=1}^q \frac{(\sigma_i - \rho_i)^2}{T_i - T'_i} \right) \right\} \\
 (4.10) \quad &\times \left(\prod_{i=1}^q d_{\sigma_i} H_i^{\#}(\rho_i, \sigma_i, \rho_{i+1}) \right) d\rho_{q+1} d\rho_q \cdots d\rho_1 \\
 &+ \sum_{Q_3(n-1)} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} \frac{g_n(\rho_{q+1}) \alpha_{0, v_2}^1 (T_i - T'_i)}{2\pi[2\pi(T'_{q+1} - T_q)]^{1/2}} \\
 &\times \left(\prod_{i=2}^q \frac{g_{v_i}(\rho_i) (T_i - T'_i)^{1/2} \alpha_{v_i, v_{i+1}}^{v_i+1}}{2\pi(T'_i - T_{i-1})^{1/2}} \right) \\
 &\times \exp\left\{ -\frac{\lambda}{2} \left(\sum_{i=2}^{q+1} \frac{(\rho_i - \sigma_{i-1})^2}{T'_i - T_{i-1}} + \sum_{i=1}^q \frac{(\sigma_i - \rho_i)^2}{T_i - T'_i} + w^2 \right) \right\} \\
 &\times \left(\prod_{i=1}^q d_{\sigma_i} H_i^{\#}(\rho_i, \sigma_i, \rho_{i+1}) \right) d\rho_{q+1} d\rho_q \cdots d\rho_3 d\rho_2 dw
 \end{aligned}$$

where $Q_j(n-1)$ is the class of all finite sets of integers $\nu_1, \nu_2, \dots, \nu_{q+1}$ satisfying

$$Q_1(n-1) = \{\nu_1, \dots, \nu_{q+1} \mid \nu_1 = 1, \nu_{i+1} \geq \nu_i + 2, i = 1, 2, \dots, q, \nu_{q+1} = n\}$$

$$Q_2(n-1) = \{\nu_1, \dots, \nu_{q+1} \mid \nu_1 > 1, \nu_{i+1} \geq \nu_i + 2, i = 1, 2, \dots, q, \nu_{q+1} = n\}$$

$$Q_3(n-1) = \{\nu_1, \dots, \nu_{q+1} \mid \nu_1 = 0, \nu_{i+1} \geq \nu_i + 2, i = 1, 2, \dots, q, \nu_{q+1} = n\}$$

$$(4.11) \quad H_i^{\#}(\rho_i, \sigma_i, \rho_{i+1})$$

$$\begin{aligned}
 &= \frac{\prod_{k=\nu_i+1}^{\nu_{i+1}-1} g_k(\alpha_{\nu_i+1, \nu_{i+1}}^k \sigma_i + \beta_{\nu_i+1, \nu_{i+1}}^k \rho_{i+1}) - \prod_{k=\nu_i+1}^{\nu_{i+1}-1} g_k(\alpha_{\nu_i, \nu_{i+1}}^k \rho_i + \beta_{\nu_i, \nu_{i+1}}^k \rho_{i+1})}{\sigma_i - \alpha_{\nu_i, \nu_{i+1}}^{\nu_i+1} \rho_i - \beta_{\nu_i, \nu_{i+1}}^{\nu_i+1}}
 \end{aligned}$$

and where in the sum over $Q_3(n-1)$, $\rho_1 \equiv 0$, and the integrals in the two sums exist both as iterated Riemann-Riemann Stieltjes integrals and as iterated Lebesgue-Lebesgue Stieltjes integrals.

Now define $\phi_i^*(\sigma_i, \rho_{i+1})$ by the equation

$$(4.12) \quad \begin{aligned} \phi_i^*(\sigma_i, \rho_{i+1}) = & \sum_{k=\nu_i+1}^{\nu_{i+1}-1} \left\{ \alpha_{\nu_i+1, \nu_i+1}^k \phi_k(\alpha_{\nu_i+1, \nu_i+1}^k \sigma_i + \beta_{\nu_i+1, \nu_i+1}^k \rho_{i+1}) \right. \\ & \times \left. \prod_{\substack{j=\nu_i+1 \\ j \neq k}}^{\nu_{i+1}-1} g_j(\alpha_{\nu_i+1, \nu_i+1}^j \sigma_i + \beta_{\nu_i+1, \nu_i+1}^j \rho_{i+1}) \right\}. \end{aligned}$$

Then as $g'_k = \phi_k$ a.e. we see that

$$\frac{d}{d\sigma_i} \left[\prod_{k=\nu_i+1}^{\nu_{i+1}-1} g_k(\alpha_{\nu_i+1, \nu_i+1}^k \sigma_i + \beta_{\nu_i+1, \nu_i+1}^k \rho_{i+1}) \right] = \phi_i^*(\sigma_i, \rho_{i+1}) \text{ a.e.}$$

Thus for any $d > 0$ and $\rho_i, \rho_{i+1} \in [-d, d]$ we apply Lemma 4.1 and 4.2 and obtain

$$(4.13) \quad \begin{aligned} & \text{Var}(H_i^*(\rho_i, s, \rho_{i+1}), -2d \leq s \leq 2d) \\ & \leq \text{Var}(\phi_i^*(s, \rho_{i+1}), -2d \leq s \leq 2d) \\ & \leq (\nu_{i+1} - \nu_i - 1)^2 B^{\nu_{i+1}-\nu_i-1} e^{2B} (3d)^r. \end{aligned}$$

Hence we see that H_i^* is of bounded variation as a function of σ_i on every bounded interval and as $0 \leq \gamma < 2$ the integrals on the right side of (4.10) converge absolutely for $\lambda > 0$.

Next we apply Lemma 4.3 to the right side of (4.10) and obtain by use of (4.3)

$$(4.14) \quad \int_{C[a,b]} F(\lambda^{-\frac{1}{2}} x) dx = \lambda^{\frac{1}{2}} \int_0^\infty e^{-\lambda u} df(u)$$

where $f(u) = f_1(u) + f_2(u) + f_3(u)$ and

$$\begin{aligned} f_1(u) &= \int_{V_1(u)} \frac{g_n(s) g_{0,n}^{\#}(0, s) ds}{\sqrt{2\pi(b-a)}} \\ f_2(u) &= \sum_{\substack{2 \\ j=1}} Q_j(n-1) \int_{V_2(u)} \cdots \int_{V_2(u)}^{(2q+1)} \frac{g_n(\rho_{q+1}) h^{\#}(\rho_1)}{[2\pi(T'_{q+1} - T_q)]^{1/2}} \left(\prod_{i=1}^q \frac{g_{\nu_i}(\rho_i) \alpha_{\nu_i, \nu_{i+1}}^{\nu_i+1} (T_i - T'_i)^{1/2}}{2\pi(T'_i - T_{i-1})^{1/2}} \right) \\ & \quad \times \left(\prod_{i=1}^q d_{\sigma_i} H_i^*(\rho_i, \sigma_i, \rho_{i+1}) \right) d\rho_{q+1} \cdots d\rho_2 d\rho_1 \\ f_3(u) &= \sum_{Q_3(n-1)} \int_{V_3(u)} \cdots \int_{V_3(u)}^{(2q+1)} \frac{g_n(\rho_{q+1}) \alpha_{0, \nu_2}^1 (T_1 - T'_1)^{1/2}}{2\pi[2\pi(T'_{q+1} - T_q)]^{1/2}} \\ & \quad \times \left(\prod_{i=2}^q \frac{g_{\nu_i}(\rho_i) \alpha_{\nu_i, \nu_{i+1}}^{\nu_i+1} (T_i - T'_i)^{1/2}}{2\pi(T'_i - T_{i-1})^{1/2}} \right) \\ & \quad \times \left(\prod_{i=1}^q d_{\sigma_i} H_i^*(\rho_i, \sigma_i, \rho_{i+1}) \right) d\rho_{q+1} \cdots d\rho_2 dw \end{aligned}$$

$$\begin{aligned}
V_1(u) &= \{s \mid s^2 < 2u(b-a)\} \\
V_2(u) &= \left\{ [\rho_1, \dots, \rho_{q+1}, \sigma_1, \dots, \sigma_q] \mid \right. \\
&\quad \times \sum_{i=1}^{q+1} \frac{(\rho_i - \sigma_{i-1})^2}{T'_i - T_{i-1}} + \sum_{i=1}^q \frac{(\sigma_i - \rho_i)^2}{T_i - T'_i} < 2u \left. \right\} \\
V_3(u) &= \left\{ [w, \rho_2, \dots, \rho_{q+1}, \sigma_1, \dots, \sigma_q] \mid w^2 \right. \\
&\quad \left. + \sum_{i=2}^{q+1} \frac{(\rho_i - \sigma_{i-1})^2}{T'_i - T_{i-1}} + \sum_{i=1}^q \frac{(\sigma_i - \rho_i)^2}{T_i - T'_i} < 2u \right\}.
\end{aligned}$$

In order to show that $F(x)$ is M Ilstow integrable for $M=1$ we need to obtain bounds on the variation of $f(u)$ for $u \geq 0$. First we note that for $j=1, 2$, and 3 , $f_j(u)$ is left continuous (by definition of $V_j(u)$) and of bounded variation on every bounded subinterval of $[0, \infty)$. Now for $j=1, 2$ and 3 , let $\hat{f}_j(u)$ denote what we obtain by replacing $g_{0,n}^\#, h^\#, g_{\nu_i}$, and $d_{\sigma_i} H_i^\#(\rho_i, \sigma_i, \rho_{i+1})$ by $|g_{0,n}^\#|, |h^\#|, |g_{\nu_i}|$ and $|d_{\sigma_i} H_i^\#(\rho_i, \sigma_i, \rho_{i+1})|$ respectively in $f_j(u)$. Then clearly $\hat{f}_j(u) \uparrow$ and for $0 \leq u_1 < u_2 < \infty$ we have

$$\text{Var}(f_j, [u_1, u_2]) \leq \text{Var}(\hat{f}_j, [u_1, u_2]) = \hat{f}_j(u_2) - \hat{f}_j(u_1).$$

Thus we see that it will suffice to obtain bounds on $\hat{f}_j(u)$. First note that for all $u \geq 0$

$$\hat{f}_1(u) \leq \int_{-\infty}^{\infty} \frac{|g_n(s)g_{0,n}^\#(0, s)|}{\sqrt{2\pi(b-a)}} ds \leq B^n[2\pi(b-a)]^{-1/2}.$$

Next we see that $V_2(u) \subseteq \hat{V}_2(u)$ and $V_3(u) \subseteq \hat{V}_3(u)$ where

$$\begin{aligned}
\hat{V}_2(u) &= \{[\rho_1, \dots, \rho_{q+1}, \sigma_1, \dots, \sigma_q] \mid |\rho_i| \leq \hat{u}, |\sigma_i| \leq 2\hat{u}\} \\
\hat{V}_3(u) &= \{[w, \rho_2, \dots, \rho_{q+1}, \sigma_1, \dots, \sigma_q] \mid |\rho_i| \leq \hat{u}, |\sigma_i| \leq 2\hat{u}, w^2 \leq 2u\}
\end{aligned}$$

and $\hat{u} = [2u(b-a)]^{\frac{1}{2}}$. Thus for $[\rho_1, \dots, \rho_{q+1}, \sigma_1, \dots, \sigma_q] \in \hat{V}_2(u)$ we have $\alpha_{\nu_i, \nu_{i+1}}^{\nu_i+1} \rho_i + \beta_{\nu_i, \nu_{i+1}}^{\nu_i+1} \rho_{i+1} \in [-2\hat{u}, 2\hat{u}]$ for $i=1, 2, \dots, q$, so that by Lemmas 4.1 and 4.2 and equation (4.13) we see that

$$\begin{aligned}
\text{Var}(H_i^\#(\rho_i, s, \rho_{i+1}), -2\hat{u} \leq s \leq 2\hat{u}) \\
\leq (\nu_{i+1} - \nu_i - 1)^2 B^{\nu_{i+1} - \nu_i - 1} \exp 2B(3\hat{u})^r.
\end{aligned}$$

Hence by replacing $V_2(u)$ by $\hat{V}_2(u)$ in $\hat{f}_2(u)$ we obtain

$$\begin{aligned}
\hat{f}_2(u) &\leq \sum_{j=1}^2 \sum_{Q_j^{(n-1)}} \left(\prod_{i=1}^q (\nu_{i+1} - \nu_i - 1)^2 B^{\nu_{i+1} - \nu_i - 1} e^{2B(3\hat{u})^\gamma} \right) \left(\prod_{i=1}^q \frac{\alpha_{\nu_i, \nu_{i+1}}^{\nu_i+1} (T_i - T'_i)^{1/2}}{2\pi(T'_i - T_{i-1})^{1/2}} \right) \\
&\quad \times \int_{-\hat{u}}^{\hat{u}} \cdots \int_{-\hat{u}}^{\hat{u}} \frac{|h^*(\rho_1)g_n(\rho_{q+1})|}{[2\pi(T'_{q+1} - T_q)]^{1/2}} \left(\prod_{i=1}^q |g_{\nu_i}(\rho_i)| \right) d\rho_{q+1} \cdots d\rho_1 \\
&\leq \sum_{j=1}^2 \sum_{Q_j^{(n-1)}} \frac{B^n e^{2Bq(3\hat{u})^\gamma}}{[2\pi(T'_{q+1} - T_q)]^{1/2}} \prod_{i=1}^q \frac{(T_i - T'_i)^{1/2} (\nu_{i+1} - \nu_i - 1)^2 \alpha_{\nu_i, \nu_{i+1}}^{\nu_i+1}}{2\pi(T'_i - T_{i-1})^{1/2}} \\
&< \frac{2^{n-1} B^n e^{Bn[3(2u(b-a))^{1/2}]^\gamma}}{3[2\pi(t_1 - a)]^{1/2}}
\end{aligned}$$

since $q \leq (n/2)$, while by (2.6) of [4] and (7.38) of [2],

$$\begin{aligned}
&\sum_{j=1}^2 \sum_{Q_j^{(n-1)}} [2\pi(T'_{q+1} - T_q)]^{-1/2} \prod_{i=1}^q \frac{(T_i - T'_i)^{1/2} (\nu_{i+1} - \nu_i - 1)^2 \alpha_{\nu_i, \nu_{i+1}}^{\nu_i+1}}{2\pi(T'_i - T_{i-1})^{1/2}} \\
&\leq [2\pi(t_1 - a)]^{-1/2} \sum_{j=1}^2 \left(\frac{1}{2\pi} \right)^q \prod_{i=1}^q (\nu_{i+1} - \nu_i - 1)^2 \\
&\leq [2\pi(t_1 - a)]^{-1/2} 3^{-1} \exp \{ (n-1)\pi^{-1/3} \} \\
&< 2^{n-1} 3^{-1} [2\pi(t_1 - a)]^{-1/2}.
\end{aligned}$$

Thus $0 \leq \gamma < 2$ implies $\int_0^\infty e^{-\lambda u} d\hat{f}_2(u) < \infty$ for $\lambda > 0$.

Similarly we can show that for $u > 0$

$$\hat{f}_3(u) \leq 2^{(2n+1)/2} B^n \exp(nB(3[2u(b-a)]^{\frac{1}{2}})^\gamma) 3^{-1} \pi^{-\frac{1}{2}} [2u(b-a)]^{\frac{1}{2}}$$

and hence for $\lambda > 0$, $\int_0^\infty e^{-\lambda u} d\hat{f}_3(u) < \infty$. Thus for $\lambda > 0$

$$N_\lambda(F) = \int_\lambda^\infty e^{-\lambda \rho} d\rho \int_{C[a,b]} |F(\rho^{-\frac{1}{2}}x)| dx \leq B^n$$

and

$$N'_\lambda(1, F) = \int_0^\infty e^{-\lambda u} |df(u)| \leq \sum_{j=1}^3 \int_0^\infty e^{-\lambda u} d\hat{f}_j(u) < \infty.$$

so that $F(x) \in B(1, \lambda)$ for $\lambda > 0$ and in particular, $F(x)$ is M Ilstow integrable for $M \geq 1$.

Next for $\lambda \ni \operatorname{Re} \lambda \geq 0$ let

$$(4.15) \quad \psi(\lambda) = \lambda^{n/2} \int_{-\infty}^\infty \cdots \int_{-\infty}^\infty g_1(u_1) \cdots g_n(u_n) E_1^0 \cdots E_n^{n-1} du_n \cdots du_1.$$

Then by (7.5) of [2] we have $\psi(\lambda) = \int_{C[a,b]} F(\lambda^{-\frac{1}{2}}x) dx$ for all $\lambda > 0$. Clearly $\psi(\lambda)$ is analytic for $\operatorname{Re} \lambda > 0$ and continuous for $\operatorname{Re} \lambda \geq 0$.

Hence $\int_{C[a,b]} F(\lambda^{-\frac{1}{2}}x)dx$ has an analytic extension throughout the right half plane continuous up to the imaginary axis. But for almost all $\rho > 0$, $\int_{C[a,b]} F(\rho^{-\frac{1}{2}}x)dx = \rho^{\frac{1}{2}} \int_0^\infty e^{-\rho s} df(s)$. Thus as $\int_0^\infty e^{-\rho s} |df(s)| < \infty$ for all $\rho > 0$ we know that $\rho^{\frac{1}{2}} \int_0^\infty e^{-\rho s} df(s)$ is analytic in the right half plane and equal to the analytic extension of $\int_{C[a,b]} F(\rho^{-\frac{1}{2}}x)dx$ there. Thus $\lambda^{\frac{1}{2}} \int_0^\infty e^{-\lambda s} df(s)$ approaches a limiting value on the imaginary axis when we approach from the right hand plane and so the limiting M Feynman integral of F exists for $M \geq 1$ which concludes proof of theorem.

If we restrict the bounds on $\text{Var } \phi_i$, $\text{Var } g_i$, and ϕ_i we obtain the following corollary which proves useful in a later section.

COROLLARY TO THEOREM 4. *Assume the hypotheses of Theorem 4 are satisfied where conditions (4.7)–(4.9) are replaced by*

$$\int_{-V}^V |d\phi_i(s)| \leq B[1 + V^d],$$

$$\int_{-V}^V |dg_i(s)| \leq B[1 + V^d], \quad \text{and} \quad |\phi_i(u)| \leq B[1 + |u|^d]$$

for $i = 1, 2, \dots, n-1$, $u \in R_1$, $V > 0$, and $d > 0$. Then the conclusions of Theorem 4 hold and in addition we have $N_\lambda(F) \leq B^n$ and

$$N'_\lambda(1, F) \leq B^n [2\pi(b-a)]^{\frac{1}{2}}$$

$$+ \frac{2^{n-1}B^n(1+3^d)^n}{3[2\pi(t_1-a)]^{1/2}} \left(1 + \left[\frac{2(b-a)}{\lambda} \right]^{dn/2} \Gamma \left[\frac{dn+2}{2} \right] \right)$$

$$+ \frac{2^{n+1}B^n(1+3^d)^n}{3(2\pi)^{1/2}} \left(1 + \left[\frac{2(b-a)}{\lambda} \right]^{(dn+1)/2} \Gamma \left[\frac{dn+3}{2} \right] \right).$$

REMARK. Let us point out at this time that the techniques used in Theorem 4 allow us to strengthen Theorems 7.1 and 7.2 of [2]. For, clearly, if we put the same restrictions on g_n as we put on each g_i , $i = 1, 2, \dots, n-1$, then the conclusions of Theorem 4 hold for $M = 0$. Also if condition (4.6) was relaxed so that

$$\int_{-V}^V |g_i(s)| ds \leq B e^{B^V}, \quad \text{for } i = 1, 2, \dots, n,$$

then one can show, using the techniques of Theorem 4, that $F \in I_0$ and $F \in B(0, \lambda)$ for all $\lambda > 0$. However these same techniques do not imply $F \in \overrightarrow{W}(M, 0)$ since in this case $\psi(\lambda)$ as defined by (4.15) is not necessarily continuous up to $\text{Re } \lambda = 0$.

5. Entire functions of integrals of $\theta(s, x(s))$. Our first theorem

deals with M Ilstow and limiting M Feynman integrability of functionals of the form

$$(5.1) \quad F^*(x) = \Phi \left[\int_a^b \theta(s, x(s)) ds \right] g(x(b)) .$$

THEOREM 5.1. *Assume $\Phi(z)$ is an entire function satisfying $|\Phi(z)| \leq Ae^{B|z|}$. For each $s \in [a, b]$ assume $\theta(s, u)$ is absolutely continuous as a function of u on every bounded interval of R_1 . Also, for each $s \in [a, b]$, assume $\hat{\theta}(s, u) = \text{ess}_{h \rightarrow 0+} \lim \theta_u(s, u + h)$ is of bounded variation as a function of u on every bounded interval of R_1 . Furthermore assume $|\theta(s, V)|$, $\int_{-\infty}^{\infty} |g(w)| dw$, and $\int_{-\infty}^{\infty} |\theta(s, w)| dw$ are bounded by B while $|\hat{\theta}(s, V)|$,*

$$\int_{-V}^V |d_w \theta(s, w)| \quad \text{and} \quad \int_{-V}^V |d_w \hat{\theta}(s, w)|$$

are bounded by $B(1 + |V|^d)$ for each $s \in [a, b]$, $V \in R_1$ and some $d \ni 0 \leq d < 2$. Then for $M = 1, 2, \dots$, the functional $F^(x)$ as defined by (5.1) is M Ilstow and limiting M Feynman integrable for all values of the parameter and $F^*(x) \in B(M, \lambda)$ for all $\lambda > 0$.*

Proof. Since $\Phi(z)$ is an entire function such that $|\Phi(z)| \leq Ae^{B|z|}$ we obtain, by use of Cauchy's inequality, that $\Phi(z) = \sum_{n=0}^{\infty} a_n z^n$ for all z where $a_0 = 1$ and for $n > 0$

$$(5.2) \quad |a_n| \leq A \left(\frac{Be}{n} \right)^n .$$

Thus on $C[a, b]$ we have $F^*(x) = \sum_{n=0}^{\infty} a_n F_n^*(x)$ where

$$F_n^*(x) = g(x(b)) \left[\int_a^b \theta(s, x(s)) ds \right]^n$$

for $n = 0, 1, 2, \dots$. Now for $n > 0$ we have

$$F_n^*(x) = \int_a^b \cdots \int_a^b F_n(x; s) ds_1 \cdots ds_n$$

where

$$F_n(x; s) = F_n(x; s_1, \dots, s_n) = g(x(b)) \prod_{i=1}^n \theta(s_i, x(s_i))$$

when s_1, \dots, s_n are distinct elements of (a, b) but $F_n(x; s) = 0$ when s_1, \dots, s_n are elements of $[a, b]$ but a, s_1, \dots, s_n, b are not distinct.

We note that for each $n > 0$ and each $s \in [a, b]^n$, $F_n(x; s)$ satisfies the hypotheses of the Corollary to Theorem 4. Hence for each s in $[a, b]^n$ we have for $\lambda > 0$

$$\begin{aligned}
N_\lambda(F_n(\cdot; s)) &\leq B^{n+1}, \\
N'_\lambda(1, F_n(\cdot; s)) &\leq H(s) \\
&\equiv B^{n+1}[2\pi(b-a)]^{-\frac{1}{2}} + K_1(n) + K_2(n)[\min\{s_1, \dots, s_n\} - a]^{-\frac{1}{2}}
\end{aligned}$$

where

$$\begin{aligned}
K_1(n) &= 2^{n+2} B^{n+1} (1 + 3^d)^{n+1} 3^{-1} (2\pi)^{-1/2} \\
&\quad \times \left(1 + \left[\frac{2(b-a)}{\lambda} \right]^{[d(n+1)+1]/2} \Gamma \left[\frac{d(n+1)+3}{2} \right] \right)
\end{aligned}$$

and

$$\begin{aligned}
K_2(n) &= 2^n B^{n+1} (1 + 3^d)^{n+1} 3^{-1} (2\pi)^{-1/2} \\
&\quad \times \left(1 + \left[\frac{2(b-a)}{\lambda} \right]^{[d(n+1)]/2} \Gamma \left[\frac{d(n+1)+2}{2} \right] \right).
\end{aligned}$$

Moreover for each s , $F_n(x; s)$ is M Ilstow and limiting M Feynman integrable for each value of the parameter for $M = 1, 2, \dots$, and $F_n(x; s) \in B(1, \lambda)$ for all $\lambda > 0$. By Corollary to Theorem 6.4 of [2] (which can be generalized to apply here) we obtain

$$\begin{aligned}
N'_\lambda(1, F_n^\#) &\leq \int_a^b \cdots \int_a^b H(s) ds_1 \cdots ds_n \\
&< B^{n+1} (b-a)^n [2\pi(b-a)]^{-\frac{1}{2}} + (b-a)^n K_1(n) + 2n(b-a)^n K_2(n)
\end{aligned}$$

where the last inequality follows from (8.12) of [2]. Clearly $N_\lambda(F_n^\#) \leq B^{n+1} (b-a)^n$ so that for $\lambda > 0$

$$\begin{aligned}
N_\lambda \left(1, \sum_{n=0}^{\infty} a_n F_n^\# \right) &\leq \sum_{n=0}^{\infty} N_\lambda(1, a_n F_n^\#) \\
&\leq ([2\pi(b-a)]^{-\frac{1}{2}} + 1) \sum_{n=0}^{\infty} B^{n+1} (b-a)^n a_n \\
&\quad + \sum_{n=0}^{\infty} (b-a)^n a_n [K_1(n) + 2nK_2(n)]
\end{aligned}$$

where $K_1(0) = K_2(0) = 1$. By use of (5.2) the above series converge for $0 \leq d < 2$ by the ratio test and thus the series $\sum_{n=0}^{\infty} a_n F_n^\#$ converges in the $B(1, \lambda)$ topology for each $\lambda > 0$ and since the series also converges uniformly to $F^\#$ it converges in the $B(1, \lambda)$ topology to $F^\#$ and so $F^\# \in B(1, \lambda)$ for all $\lambda > 0$. In particular $F^\#$ is M Ilstow integrable for $M = 1, 2, \dots$.

Next we will show that $F^\#$ is limiting M Feynman integrable for $M = 1$. First apply Corollary to Theorem 4 to the functionals $F_n(x; s)$ and obtain functions $\psi_n(\lambda, s)$ which are analytic for $\operatorname{Re} \lambda > 0$ and continuous for $\operatorname{Re} \lambda \geq 0$ for each $n > 0$ and $s \in [a, b]^n$. In addition for $\operatorname{Re} \lambda \geq 0$ we have

$$\psi_n(\lambda, s) = \lambda^{(n+1)/2} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} g(u_{n+1}) \prod_{i=1}^n \theta(s_i, u_i) E_1^0 E_2^1 \cdots E_n^{n+1} du_{n+1} \cdots du_1$$

so that for $s \in [a, b]^n$ and $\operatorname{Re} \lambda \geq 0$, $|\psi_n(\lambda, s)| \leq |\lambda|^{(n+1)/2} b^{n+1}$. Hence letting

$$\psi_n^{\#}(\lambda) = \int_a^b \cdots \int_a^b \psi_n(\lambda, s) ds_1 \cdots ds_n$$

we have

$$(5.3) \quad |\psi_n^{\#}(\lambda)| \leq |\lambda|^{(n+1)/2} b^{n+1} (b-a)^n$$

for $\operatorname{Re} \lambda \geq 0$ and $n = 1, 2, \dots$. Clearly $\psi_n^{\#}(\lambda)$ is analytic for $\operatorname{Re} \lambda > 0$ and continuous for $\operatorname{Re} \lambda \geq 0$ for $n = 1, 2, \dots$. Similarly we apply Corollary to Theorem 4 to $F_0^{\#}$ and obtain $\psi_0^{\#}(\lambda)$ with the same properties.

Next we see that as the series $\sum_{n=0}^{\infty} a_n F_n^{\#}(x)$ converges uniformly in x over $C[a, b]$ we have for each real $\lambda > 0$

$$(5.4) \quad \begin{aligned} \int_{C[a, b]} F^{\#}(\lambda^{-\frac{1}{2}} x) dx &= \sum_{n=0}^{\infty} a_n \int_{C[a, b]} F_n^{\#}(\lambda^{-\frac{1}{2}} x) dx \\ &= \sum_{n=0}^{\infty} a_n \psi_n^{\#}(\lambda). \end{aligned}$$

But by (5.2) and (5.3) the last member of (5.4) converges uniformly in λ for any closed subset of $\operatorname{Re} \lambda \geq 0$ and hence the first member of (5.4) has an analytic extension throughout $\operatorname{Re} \lambda > 0$. Now let $f_1^{\#}$ denote the M Ilstow integral of $F^{\#}$ for $M = 1$. Then

$$N'_\lambda(1, F^{\#}) = \int_0^{\infty} e^{-\lambda s} |df_1^{\#}(s)|$$

for each $\lambda > 0$ and for almost all $\rho > 0$

$$\int_{C[a, b]} F^{\#}(\rho^{-\frac{1}{2}} x) dx = \rho^{\frac{1}{2}} \int_0^{\infty} e^{-\rho s} df_1^{\#}(s).$$

Hence $\rho^{\frac{1}{2}} \int_0^{\infty} e^{-\rho s} df_1^{\#}(s)$ has an analytic extension in the half plane $\operatorname{Re} \rho > 0$, continuous for $\operatorname{Re} \rho \geq 0$, and so the limiting M Feynman integral of $F^{\#}$ exists for $M = 1$ which concludes proof of theorem.

Next we will show that for appropriate functions θ and σ , the analytic Wiener and limiting M Feynman integrals of the functional

$$(5.5) \quad F(t, \xi, x) = \exp \left[\int_0^t \theta[t-s, x(s) + \xi] ds \right] \sigma[x(t) + \xi]$$

satisfy a certain integral equation.

THEOREM 5.2. Assume $\theta(t, \xi)$ is bounded and continuous on $[0, t_0] \otimes (-\infty, \infty)$ and satisfies for all $t \in [0, t_0]$

$$(5.6) \quad \int_{-\infty}^{\infty} |\theta(t, \xi)| d\xi \leq B.$$

Assume $\sigma(\xi)$ is bounded and continuous on $(-\infty, \infty)$ and satisfies

$$(5.7) \quad \int_{-\infty}^{\infty} |\sigma(\xi)| d\xi \leq B.$$

Then the analytic Wiener integral

$$(5.8) \quad G(t, \xi, \lambda) = \int_{C[0, t_0]}^{aw_\lambda} F(t, \xi, x) dx$$

exists for $(t, \xi, \lambda) \in H = (0, t_0) \otimes (-\infty, \infty) \otimes \{\operatorname{Re} \lambda \geq 0\}$ and satisfies in H the integral equation

$$(5.9) \quad \begin{aligned} G(t, \xi, \lambda) = & \left(\frac{\lambda}{2\pi t} \right)^{1/2} \int_{-\infty}^{\infty} \sigma(u) \exp \left(-\frac{\lambda(\xi - u)^2}{2t} \right) du \\ & + \left(\frac{\lambda}{2\pi} \right)^{1/2} \int_0^t (t-s)^{-1/2} ds \int_{-\infty}^{\infty} \theta(s, u) G(s, u, \lambda) \exp \left(-\frac{\lambda(\xi - u)^2}{2(t-s)} \right) ds \end{aligned}$$

with boundary condition

$$(5.10) \quad \lim_{t \rightarrow 0+} G(t, \xi, \lambda) = \sigma(\xi)$$

holding for all $(\xi, \lambda) \in (-\infty, \infty) \otimes \{\operatorname{Re} \lambda > 0\}$.

Proof. The existence of $G(t, \xi, \lambda)$ for all points of H follows from Theorem 8.2 of [2]. Moreover by (0.11) of [2] and the continuity of the functions involved we have

$$(5.11) \quad G(t, \xi, \lambda) = \int_{C[0, t]} \exp \left[\int_0^t \theta[t-s, \lambda^{-1/2}x(s) + \xi] ds \right] \sigma[\lambda^{-1/2}x(t) + \xi] dx$$

for $\lambda > 0$ and $(t, \xi) \in H_1 = (0, t_0) \otimes (-\infty, \infty)$.

Next we note that by use of Theorem 7 on page 66 of [3] (which can be proved for $n=1$ under weakened conditions on σ^* and θ^*) we have that if $\sigma^*(\xi^*)$ and $\theta^*(t, \xi^*)$ are bounded and continuous for $(t, \xi^*) \in H_2 = [0, t_0] \otimes (-\infty, \infty)$ then

$$(5.12) \quad G^*(t, \xi^*) = \int_{C[0, t]} \exp \left[\int_0^t \theta^*[t-s, x(s) + \xi^*] ds \right] \sigma^*[x(t) + \xi^*] dx$$

exists on H_2 and on H_1 satisfies

$$\begin{aligned}
(5.13) \quad G^*(t, \xi^*) &= (2\pi t)^{-1/2} \int_{-\infty}^{\infty} \sigma^*(u) \exp\left(-\frac{(\xi^* - u)^2}{2t}\right) du \\
&+ (2\pi)^{-1/2} \int_0^t (t-s)^{-1/2} ds \int_{-\infty}^{\infty} \theta^*(s, u) G^*(s, u) \exp\left(-\frac{(\xi^* - u)^2}{2(t-s)}\right) du
\end{aligned}$$

with boundary condition

$$(5.14) \quad \lim_{t \rightarrow 0^+} G^*(t, \xi^*) = \sigma^*(\xi^*) .$$

Thus, if for $\lambda > 0$ we define $\theta^*(t, u) \equiv \theta(t, \lambda^{-1/2}u)$, $\sigma^*(u) \equiv \sigma(\lambda^{-1/2}u)$, and $\xi^* \equiv \lambda^{-1/2}\xi$ then θ^* and σ^* are bounded and continuous and so $G^*(t, \xi^*)$ as defined by (5.12) exists on H_2 and on H_1 satisfies (5.13) and (5.14). But for $\lambda > 0$, $\theta^*[t-s, x(s) + \xi^*] = \theta[t-s, \lambda^{-1/2}x(s) + \xi]$, $\sigma^*[x(s) + \xi^*] = \sigma[\lambda^{-1/2}x(s) + \xi]$, and $G(t, \xi, \lambda) = G^*(t, \lambda^{1/2}\xi)$. Hence substituting into (5.13) we see that for $\lambda > 0$ and $(t, \xi) \in H_1$, $G(t, \xi, \lambda)$ satisfies (5.9). Then by use of Corollary to Theorem 8.2 of [2] $G(t, \xi, \lambda)$ is bounded in any closed bounded subset of $\text{Re } \lambda \geq 0$ for all $(t, \xi) \in H_1$ and so by use of (5.6) and (5.7) we see that both sides of (5.9) are analytic functions of λ for $\text{Re } \lambda > 0$, continuous for $\text{Re } \lambda \geq 0$ and as (5.9) holds for all $\lambda > 0$ it must hold throughout H .

Boundary condition (5.10) is verified by noting that

$$\begin{aligned}
(5.15) \quad \lim_{t \rightarrow 0^+} \left(\frac{\lambda}{2\pi} \right)^{1/2} \int_0^t (t-s)^{-1/2} ds \\
\times \int_{-\infty}^{\infty} \theta(s, u) G(s, u, \lambda) \exp\left(-\frac{\lambda(\xi - u)^2}{2(t-s)}\right) du = 0
\end{aligned}$$

and

$$\begin{aligned}
&\left| \left(\frac{\lambda}{2\pi t} \right)^{1/2} \int_{-\infty}^{\infty} \sigma(u) \exp\left(-\frac{\lambda(\xi - u)^2}{2t}\right) du - \sigma(\xi) \right| \\
&= \left| \left(\frac{\lambda}{2\pi t} \right)^{1/2} \int_{-\infty}^{\infty} [\sigma(u + \xi) - \sigma(\xi)] \exp\left(-\frac{\lambda u^2}{2t}\right) du \right| \\
&\rightarrow 0 \text{ as } t \rightarrow 0^+ .
\end{aligned}$$

THEOREM 5.3. Assume $\theta(t, \xi)$ satisfies the hypotheses of Theorem 5.1 where $[a, b] = [0, t_0]$. Assume $\sigma(\xi)$ satisfies the hypotheses of Theorem 5.2. Then for $M = 1, 2, \dots$, the limiting M Feynman integral

$$(5.16) \quad \hat{G}(t, \xi, q) = \int_{C[0, t]}^{\rightarrow f_q^M} F(t, \xi, x) dx$$

where $F(t, \xi, x)$ is defined by (5.5), exists for

$$(t, \xi, q) \in \hat{H} = (0, t_0) \otimes R_1 \otimes (R_1 - \{0\})$$

and satisfies in \hat{H} the integral equation (1.2). Furthermore if we assume $\text{Var}_{(-\infty, \infty)} \sigma(\xi) \leq B$ then

$$(5.17) \quad \lim_{t \rightarrow 0^+} \hat{G}(t, \xi, q) = \sigma(\xi) .$$

Proof. The existence of (5.16) for $M = 1, 2, \dots$, and all $(t, \xi, q) \in \hat{H}$ follows from Theorem 5.1.. But by use of Theorem 2.2 we see that $\hat{G}(t, \xi, q) = G(t, \xi, -qi)$ where G is defined by (5.8). Then substituting into (5.9) establishes (1.2).

To establish (5.17) first note that

$$\lim_{t \rightarrow 0^+} \left(\frac{-iq}{2\pi} \right)^{1/2} \int_0^t (t-s)^{-1/2} ds \int_{-\infty}^{\infty} \theta(s, u) \hat{G}(s, u, q) \exp \left(\frac{qi(\xi - u)^2}{2(t-s)} \right) du = 0$$

for all $\xi, q \in R_1, q \neq 0$. Also by contour integration and integration by parts one obtains

$$\begin{aligned} & \left(\frac{-iq}{2\pi t} \right)^{1/2} \int_{\xi}^{\infty} \sigma(u) \exp \left(\frac{qi(\xi - u)^2}{2t} \right) du \\ &= \frac{\sigma(\xi)}{2} + \left(\frac{1}{2\pi} \right)^{1/2} \int_{\xi}^{\infty} \left[\int_{-(iq/t)^{1/2}(u-\xi)}^{(-iq/t)^{1/2}\infty} \exp \left(-\frac{z^2}{2} \right) dz \right] d\sigma(u) \\ & \rightarrow \frac{\sigma(\xi)}{2} \quad \text{as } t \rightarrow 0^+ \text{ by dominated convergence.} \end{aligned}$$

Similarly

$$\left(\frac{-iq}{2\pi t} \right)^{1/2} \int_{-\infty}^{\xi} \sigma(u) \exp \left(\frac{qi(\xi - u)^2}{2t} \right) du \rightarrow \frac{\sigma(\xi)}{2} \quad \text{as } t \rightarrow 0^+ .$$

REMARK. In view of the remark at the end of § 4 we see that the conclusions of Theorem 5.1 will hold for $M = 0$ under the additional assumption that g' is of bounded variation on every bounded interval and satisfies

$$\int_{-V}^V |dg'(w)| \leq B[1 + |V|^d] .$$

Also the conclusion of Theorem 5.3 will hold for $M = 0$ under the additional assumption that σ' is of bounded variation on every bounded interval and satisfies

$$\int_{-V}^V |d\sigma'(w)| \leq B[1 + |V|^d] .$$

The author wishes to express his gratitude to Professor Robert Cameron for his encouragement and valuable advise .

BIBLIOGRAPHY

1. R. H. Cameron, *A family of integrals serving to connect the Wiener and Feynman integrals*, J. Math. and Physics **39** (1960), 126-140.
2. ———, *The Ilstow and Feynman integrals*, J. D'Analyse Mathématique **10** (1962-63), 287-361.
3. ———, *Nonlinear Volterra functional equations and linear parabolic differential systems*, J. D'Analyse Mathématique **5** (1956), 136-182.
4. R. A. Kallman, *On using the Ilstow integral to solve a certain partial differential system*, (Submitted for publication)

Received April 28, 1967.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN

Stanford University
Stanford, California

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS

University of Washington
Seattle, Washington 98105

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

Pacific Journal of Mathematics

Vol. 26, No. 1

November, 1968

Efraim Pacillas Armendariz, <i>Closure properties in radical theory</i>	1
Friedrich-Wilhelm Bauer, <i>Postnikov-decompositions of functors</i>	9
Thomas Ru-Wen Chow, <i>The equivalence of group invariant positive definite functions</i>	25
Thomas Allan Cootz, <i>A maximum principle and geometric properties of level sets</i>	39
Rodolfo DeSapio, <i>Almost diffeomorphisms of manifolds</i>	47
R. L. Duncan, <i>Some continuity properties of the Schnirelmann density</i>	57
Ralph Jasper Faudree, Jr., <i>Automorphism groups of finite subgroups of division rings</i>	59
Thomas Alastair Gillespie, <i>An invariant subspace theorem of J. Feldman</i>	67
George Isaac Glauberman and John Griggs Thompson, <i>Weakly closed direct factors of Sylow subgroups</i>	73
Hiroshi Haruki, <i>On inequalities generalizing a Pythagorean functional equation and Jensen's functional equation</i>	85
David Wilson Henderson, <i>D-dimension. I. A new transfinite dimension</i>	91
David Wilson Henderson, <i>D-dimension. II. Separable spaces and compactifications</i>	109
Julien O. Hennefeld, <i>A note on the Arens products</i>	115
Richard Vincent Kadison, <i>Strong continuity of operator functions</i>	121
J. G. Kalbfleisch and Ralph Gordon Stanton, <i>Maximal and minimal coverings of $(k - 1)$-tuples by k-tuples</i>	131
Franklin Lowenthal, <i>On generating subgroups of the Moebius group by pairs of infinitesimal transformations</i>	141
Michael Barry Marcus, <i>Gaussian processes with stationary increments possessing discontinuous sample paths</i>	149
Zalman Rubinstein, <i>On a problem of Ilyeff</i>	159
Bernard Russo, <i>Unimodular contractions in Hilbert space</i>	163
David Lee Skoug, <i>Generalized Ilstow and Feynman integrals</i>	171
William Charles Waterhouse, <i>Dual groups of vector spaces</i>	193