

Pacific Journal of Mathematics

ON MEASURES WITH SMALL TRANSFORMS

RAOUF DOSS

ON MEASURES WITH SMALL TRANSFORMS

RAOUF DOSS

G is a locally compact abelian group whose dual Γ is algebraically ordered, i.e., ordered when considered as a discrete group. Every (Radon) complex measure μ on G has a unique Lebesgue decomposition: $d\mu = d\mu_s + g(x)dx$, where $d\mu_s$ is singular and $g \in L^1(G)$. A measure μ on G is of analytic type if $\hat{\mu}(\gamma) = 0$ for $\gamma < 0$, where $\hat{\mu}$ is the Fourier-Stieltjes transform of μ .

The main result of the paper is that if $\int_{\gamma < 0} |\hat{\mu}(\gamma)|^2 d\gamma < \infty$, or more generally, if, for $\gamma < 0$, $\hat{\mu}(\gamma)$ coincides with the transform $\hat{f}(\gamma)$ of a function f in $L^p(G)$, $1 \leq p \leq 2$, then the singular part $d\mu_s$ is of analytic type and $\hat{\mu}_s(0) = 0$.

Throughout the paper the symbol $M(G)$ denotes the Banach algebra under convolution of all regular complex measures on G . Haar measure will be denoted dx on G and $d\gamma$ on Γ . If the singular part $d\mu_s$, of a $\mu \in M(G)$, vanishes, then $d\mu$ is called absolutely continuous.

We first prove that if $\mu \in M(G)$ and $\hat{\mu} \in L^2(\Gamma)$, then μ is absolutely continuous. This natural statement must have been proved before, but it does not seem to appear in the literature. It is not implied by the L^1 -inversion theorem, which assumes $\mu \in M(G)$ and $\hat{\mu} \in L^1(\Gamma)$, nor by Plancherel's theorem. It is best possible in the sense that $\hat{\mu} \in L^2(\Gamma)$ cannot be replaced by the weaker condition $\hat{\mu} \in L^p(\Gamma)$, $p > 2$; for, as shown by Hewitt and Zuckerman [3], on any nondiscrete locally compact abelian group G , there exists a nonvanishing singular measure μ_s for which $\hat{\mu}_s \in L^p(\Gamma)$, for every $p > 2$.

Next we suppose that the dual Γ is algebraically ordered. This means that there exists a semi-group $P \subset \Gamma$ such that $P \cup (-P) = \Gamma$, $P \cap (-P) = \{0\}$. We do not assume that P is closed in Γ , so that, e.g., R^k , $k \geq 1$, is algebraically ordered. If P is closed in Γ , then Γ is called ordered (Rudin [4]). But then R^k is ordered only if $k = 1$. If Γ is discrete, the two notions of ordered and algebraically ordered coincide. A discrete abelian group Γ can be ordered if and only if its (compact) dual G is connected (Rudin [4], 8.1.2 (a) and 2.5.6 (c)). Thus the dual Γ of a locally compact abelian group G can be algebraically ordered if and only if the Bohr compactification \bar{G} of G is connected.

So suppose Γ is algebraically ordered. A measure $\mu \in M(G)$ is said to be of analytic type if $\hat{\mu}(\gamma) = 0$ for $\gamma < 0$. Helson and Lowdenslager [2] prove that for a compact abelian group G , with ordered dual Γ , if $\mu \in M(G)$ is of analytic type, then the singular part μ_s is of analy-

tic type and moreover $\hat{\mu}_s(0) = 0$. Our main result is a twofold generalization of this theorem, namely:

Let G be a locally compact abelian group with algebraically ordered dual Γ and let $\mu \in M(G)$. If $\int_{\gamma < 0} |\hat{\mu}(\gamma)|^2 d\gamma < \infty$ or more generally if, for $\gamma < 0$, $\hat{\mu}$ coincides with the transform \hat{f} of a function f in $L^p(G)$, $1 \leq p \leq 2$, then μ_s is of analytic type and $\hat{\mu}_s(0) = 0$.

This theorem is new even in the case $G = R$. Combined with the F. and M. Riesz theorem it yields the result: if $\mu \in M(R)$ and $\int_{\gamma < 0} |\hat{\mu}(\gamma)|^2 d\gamma < \infty$ then μ is absolutely continuous.

THEOREM 1. *Let μ be a complex measure on the locally compact abelian group G . If $\hat{\mu} \in L^2(\Gamma)$ then μ is absolutely continuous.*

Proof. (Short and due to the referee.) By Plancherel's theorem there is $f \in L^2(G)$ with $\hat{f} = \hat{\mu}$ almost everywhere. Let g be a continuous function, with compact support in G , such that $\hat{g} \in L^1(\Gamma)$. Then

$$\begin{aligned} \int_G f \bar{g} &= \int_\Gamma \hat{f} \bar{\hat{g}} && \text{(Parseval-Plancherel)} \\ &= \int_\Gamma \bar{\hat{g}}(\gamma) d\gamma \int_G \overline{f(x, \gamma)} d\mu(x) && \text{(since } \hat{f} = \hat{\mu}) \\ &= \int_G \bar{g}(x) d\mu(x) && \text{(Fubini and } L^1\text{-inversion theorem).} \end{aligned}$$

Now every continuous h with compact support C in G , can be uniformly approximated by g 's of the above type, with supports in a fixed compact set C' : choose a fixed compact neighborhood V of 0 and a kernel $k \geq 0$, bounded, with support in V , and put $g = h * k$; then

$$\text{support } g \subset C + V = C', \hat{h}, \hat{k} \in L^2(\Gamma), \hat{g} \in L^1(\Gamma),$$

and g may be chosen uniformly close to h . Hence

$$\int_G \bar{h} f = \int_G \bar{h} d\mu$$

for every continuous h with compact support in G . Therefore

$$\int_G |f| \leq \|\mu\| < \infty, f \in L^1(G);$$

since $\hat{\mu} = \hat{f}$, we conclude, by the uniqueness theorem $d\mu(x) = f(x)dx$ and μ is absolutely continuous.

LEMMA 1. *Suppose G is a locally compact abelian group whose dual Γ is algebraically ordered, $\mu \in M(G)$ and μ is of analytic type. Then the singular part of μ is also of analytic type.*

This lemma has been proved in Doss [1] under the assumption that Γ is ordered. But the proof is valid for an algebraically ordered Γ with the following obvious modifications:

The compact interval $[-\delta, \delta]$ is replaced by a compact symmetric neighborhood V of the origin in Γ . The relation $\gamma < -\delta$ is replaced throughout by $\gamma < 0, \gamma \notin V$.

Finally the function k such that

$$\begin{aligned} (1) \quad & k \in L^1(G) \quad k(x) \geq 0 \\ (2) \quad & \hat{k}(\gamma) \geq 0 \quad \hat{k}(\gamma) = 0 \text{ outside } V \end{aligned}$$

is obtained as follows:

Choose a symmetric compact neighborhood W of 0 in Γ . Let $u(\gamma) = 1/\text{meas } W$ on W , $u(\gamma) = 0$ outside W . Then $u \in L^1(\Gamma)$, $u \in L^2(\Gamma)$, $\hat{u} \in L^2(G)$. Put $v = u * u$. Then

$$(2') \quad v(\gamma) \geq 0, \quad v \text{ vanishes outside the compact (symmetric) set} \\ V = W + W.$$

Also $v \in L^1(\Gamma)$ and

$$(1') \quad \hat{v}(x) = |\hat{u}(x)|^2 \geq 0, \quad \hat{v} \in L^1(G).$$

By the inversion theorem

$$v(\gamma) = \int_G \hat{v}(x)(x, \gamma) dx.$$

Put $k(x) = \hat{v}(x)$. Then, by (1')

$$(1) \quad k \in L^1(G), \quad k(x) \geq 0.$$

Moreover, $\hat{k}(\gamma) = \int_G k(x)\overline{(x, \gamma)} dx = v(-\gamma)$. Hence, by (2')

$$(2) \quad \hat{k}(\gamma) \geq 0, \quad \hat{k}(\gamma) = 0 \text{ outside } V.$$

LEMMA 2. *Let G be a locally compact abelian group whose dual Γ is algebraically ordered. Let*

$$d\sigma = ds + w(x)dx$$

be a positive measure on G , where ds is singular and $w \in L^1(G)$. Let K be a compact set in Γ and denote by Ω the set of trigonometric polynomials $p(x)$ of the type

$$p(x) = \sum a(x, \gamma) \quad \gamma < 0, \quad \gamma \notin K.$$

Let φ be the unique function belonging to the closure of Ω in $L^2(d\sigma)$

and such that

$$\int_G |1 - \varphi|^2 d\sigma = \inf_{p \in \Omega} \int_G |1 - p|^2 d\sigma .$$

Then

$$\int_G |1 - \varphi|^2 d\sigma \leq \int_G w dx .$$

Proof. φ is the unique function belonging to the closure of Ω in $L^2(d\sigma)$, for which

$$(1) \quad \int_G \overline{(x, \gamma)}(1 - \varphi) d\sigma = 0 \quad \text{for } \gamma < 0, \gamma \notin K .$$

We can easily find, by means of an appropriate kernel, an $f \in L^1(G)$ whose transform \hat{f} is equal to the transform of the measure $(1 - \varphi)d\sigma$, for $\gamma < 0$. But then the measure $(1 - \varphi)d\sigma - f(x)dx$ is of analytic type. By Lemma 1, the singular part $(1 - \varphi)ds$ is of analytic type:

$$\int_G \overline{(x, \gamma)}(1 - \varphi) ds = 0 \quad \text{for } \gamma < 0 .$$

By continuity (or by the Helson-Lowdenslager theorem, in case Γ is discrete), the same relation holds for $\gamma = 0$. We conclude

$$\int_G \overline{(x, \gamma)} \overline{(1 - \varphi)}(1 - \varphi) ds = 0 \quad \text{for } \gamma \leq 0 ,$$

and since $|1 - \varphi|^2 ds$ is real, the above relation is true for $\gamma \geq 0$. Hence, by the uniqueness theorem:

$$(2) \quad |1 - \varphi|^2 ds = 0 \quad (1 - \varphi) ds = 0 .$$

Hence (1) reduces to

$$\int_G \overline{(x, \gamma)}(1 - \varphi) w dx = 0 \quad \text{for } \gamma < 0, \gamma \notin K .$$

Since φ belongs to the closure of Ω in $L^2(w)$ we conclude

$$\int_G |1 - \varphi|^2 w dx = \inf_{p \in \Omega} \int_G |1 - p|^2 w dx \leq \int_G w dx .$$

Hence, by (2)

$$\int_G |1 - \varphi|^2 d\sigma \leq \int_G w dx$$

and the lemma is proved.

MAIN THEOREM. *Let G be a locally compact abelian group*

whose dual Γ is algebraically ordered. Let

$$d\mu = d\mu_s + g(x)dx$$

be a complex measure on G , where $d\mu_s$ is singular and $g \in L^1(G)$. If $\int_{\gamma < 0} |\hat{u}(\gamma)|^2 d\gamma < \infty$, or more generally if, for $\gamma < 0$, $\hat{u}(\gamma)$ coincides with the transform $\hat{f}(\gamma)$ of some function $f \in L^r(G)$, $1 \leq r \leq 2$, then $d\mu_s$ is of analytic type and $\hat{u}_s(0) = 0$.

Proof. It is sufficient to prove $\hat{u}_s(0) = 0$, for by translation, we get $\hat{u}_s(\gamma) = 0$ for $\gamma < 0$. By hypothesis there is $f \in L^r(G)$, such that

$$\hat{f}(\gamma) = \hat{u}(\gamma) \quad \text{a.e. for } \gamma < 0.$$

Let $\varepsilon > 0$ be given. There is $k_1 \in L^1(G)$ such that \hat{k}_1 has compact support K_1 and such that

$$\|g - g * k_1\|_1 < \varepsilon.$$

(see e.g. [4], 2.6.6). Also there is $h_1 \in L^1(G)$ such that \hat{h}_1 has compact support H_1 and such that

$$\|f - f * h_1\|_r < \varepsilon^{1/r}$$

(the proof of 2.6.6 in [4] works unchanged). Put

$$g_1 = g - g * k_1, \quad f_1 = f - f * h_1.$$

Then

$$\|g_1\|_1 < \varepsilon, \quad \|f_1\|_r < \varepsilon^{1/r}.$$

Put

$$\begin{aligned} d\lambda &= d\mu_s + g_1(x)dx \\ d\sigma &= d|\mu_s| + |g_1(x)| dx + |f_1(x)|^r dx. \end{aligned}$$

Let V be a symmetric compact neighborhood of the origin independent of ε and the subsequent choice of k_1, h_1, K_1, H_1 . Put

$$K = K_1 + H_1 + V$$

so that K is compact.

By Lemma 2 there is a

$$p(x) = \sum a_n(x, \gamma_n) \quad \gamma_n < 0, \gamma_n \notin K$$

such that

$$(1) \quad \int_G |1 - p|^2 d\sigma \leq \varepsilon + \|g_1\|_1 + \|f_1^r\|_1 \leq 3\varepsilon.$$

Put $p_1 = \frac{2}{r} \frac{1}{p_1} + \frac{1}{q_1} = 1$. By Hölder's inequality and (1)

$$\begin{aligned} \int_G \overline{(1-p)} f_1^r dx &\leq \int_G |1-p|^r d\sigma \\ &\leq \left[\int_G |1-p|^{rp_1} d\sigma \right]^{1/p_1} \left[\int_G d\sigma \right]^{1/p_1} \leq (3\varepsilon)^{1/q_1} \sigma(G)^{1/q_1}. \end{aligned}$$

This, combined with $\|f_1\|_r < \varepsilon^{1/r}$ gives

$$(2) \quad \|\bar{p}f_1\|_r \leq \varepsilon^{1/r} + [(3\varepsilon)^{1/p_1} \sigma(G)^{1/q_1}]^{1/r}.$$

By the Schwarz inequality and (1)

$$\int_G |1-p| d\sigma \leq (3\varepsilon)^{1/2} \sigma(G)^{1/2}.$$

Hence

$$\left| \int_G \overline{(x, \gamma)} \overline{(1-p)} d\lambda \right| \leq (3\varepsilon)^{1/2} (\sigma(G))^{1/2}$$

i.e.,

$$(3) \quad |\hat{\lambda}(\gamma) - (\bar{p}d\lambda)^\wedge(\gamma)| \leq (3\varepsilon)^{1/2} \sigma(G)^{1/2}.$$

Now from the definition of $d\lambda$ and from $\hat{f}(\gamma) = \hat{u}(\gamma)$ a.e. for $\gamma < 0$ we see that

$$(4) \quad \hat{\lambda}(\delta) = \hat{u}(\delta) = \hat{f}(\delta) = \hat{f}_1(\delta) \text{ a.e. for } \delta < 0, \delta \notin K_1 \cup H_1.$$

But $\gamma_n < 0, \gamma_n \notin (K_1 \cup H_1) - V$. Hence, if $\gamma \leq 0, \gamma \in V$ we have

$$\gamma + \gamma_n < 0, \gamma + \gamma_n \notin K_1 \cup H_1.$$

Whence, by (4)

$$\int_G \overline{(x, \gamma)} \overline{(x, \gamma_n)} d\lambda = \hat{f}_1(\gamma + \gamma_n) \text{ a.e. for } \gamma \leq 0, \gamma \in V.$$

Therefore,

$$(\bar{p}d\lambda)^\wedge(\gamma) = (\bar{p}f_1)^\wedge(\gamma) \text{ a.e. for } \gamma \leq 0, \gamma \in V.$$

We deduce, by (3)

$$\begin{aligned} |\hat{\lambda}(\gamma) - (\bar{p}f_1)^\wedge(\gamma)| &\leq (3\varepsilon)^{1/2} \sigma(G)^{1/2} \\ &\text{a.e. for } \gamma \leq 0, \gamma \in V. \end{aligned}$$

Finally

$$(5) \quad \begin{aligned} |\hat{u}_s(\gamma) - (\bar{p}f_1)^\wedge(\gamma)| &< \varepsilon + (3\varepsilon)^{1/2} \sigma(G)^{1/2} \\ &\text{a.e. for } \gamma \leq 0, \gamma \in V. \end{aligned}$$

Now $\varepsilon > 0$ is arbitrary. By (2) and (5) there exists a sequence $\varphi_n \in L^*(G)$ such that

$$(6) \quad \begin{aligned} \|\varphi_n\|_r &\rightarrow 0 \\ \hat{\varphi}_n(\gamma) &\rightarrow \hat{u}_s(\gamma) \quad \text{a.e. for } \gamma \leq 0, \gamma \in V. \end{aligned}$$

We deduce from (6)

$$\|\hat{\varphi}_n\|_{r'} \rightarrow 0 \quad \left(\frac{1}{r} + \frac{1}{r'} = 1 \right).$$

This shows that $\hat{\mu}_s(\gamma) = 0$ a.e. for $\gamma \leq 0, \gamma \in V$.

In particular, by continuity, $\hat{u}_s(0) = 0$ and the theorem is proved.

REFERENCES

1. R. Doss, *On the Fourier-Stieltjes transforms of singular or absolutely continuous measures*, Math. Zeit. **97** (1967), 77-84.
2. H. Helson, and D. Lowdenslager, *Prediction theory and Fourier series in several variables*, Acta Math. **99** (1958), 165-202.
3. H. Hewitt and S. Zuckerman, *Singular measures with absolutely continuous convolution squares*, Proc. Cambridge Phil. Soc. **62** (1966), 399-420.
4. W. Rudin, *Fourier Analysis on Groups*, Interscience, New York, 1962.

Received November 27, 1967, and in revised form March 27, 1968.

STATE UNIVERSITY OF NEW YORK
STONY BROOK, NEW YORK

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN

Stanford University
Stanford, California

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

J. P. JANS

University of Washington
Seattle, Washington 98105

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California 90024.

Each author of each article receives 50 reprints free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners of publishers and have no responsibility for its content or policies.

Seymour Bachmuth and Horace Yomishi Mochizuki, <i>Kostrikin's theorem on Engel groups of prime power exponent</i>	197
Paul Richard Beesack and Krishna M. Das, <i>Extensions of Opial's inequality</i>	215
John H. E. Cohn, <i>Some quartic Diophantine equations</i>	233
H. P. Dikshit, <i>Absolute $(C, 1) \cdot (N, p_n)$ summability of a Fourier series and its conjugate series</i>	245
Raouf Doss, <i>On measures with small transforms</i>	257
Charles L. Fefferman, <i>L_p spaces over finitely additive measures</i>	265
Le Baron O. Ferguson, <i>Uniform approximation by polynomials with integral coefficients. II</i>	273
Takashi Ito and Thomas I. Seidman, <i>Bounded generators of linear spaces</i>	283
Masako Izumi and Shin-ichi Izumi, <i>Nörlund summability of Fourier series</i>	289
Donald Gordon James, <i>On Witt's theorem for unimodular quadratic forms</i>	303
J. L. Kelley and Edwin Spanier, <i>Euler characteristics</i>	317
Carl W. Kohls and Lawrence James Lardy, <i>Some ring extensions with matrix representations</i>	341
Ray Mines, III, <i>A family of functors defined on generalized primary groups</i>	349
Louise Arakelian Raphael, <i>A characterization of integral operators on the space of Borel measurable functions bounded with respect to a weight function</i>	361
Charles Albert Ryavec, <i>The addition of residue classes modulo n</i>	367
H. M. (Hari Mohan) Srivastava, <i>Fractional integration and inversion formulae associated with the generalized Whittaker transform</i>	375
Edgar Lee Stout, <i>The second Cousin problem with bounded data</i>	379
Donald Curtis Taylor, <i>A generalized Fatou theorem for Banach algebras</i>	389
Bui An Ton, <i>Boundary value problems for elliptic convolution equations of Wiener-Hopf type in a bounded region</i>	395
Philip C. Tonne, <i>Bounded series and Hausdorff matrices for absolutely convergent sequences</i>	415