

Pacific Journal of Mathematics

**FRACTIONAL INTEGRATION AND INVERSION FORMULAE
ASSOCIATED WITH THE GENERALIZED WHITTAKER
TRANSFORM**

H. M. (HARI MOHAN) SRIVASTAVA

FRACTIONAL INTEGRATION AND INVERSION FORMULAE ASSOCIATED WITH THE GENERALIZED WHITTAKER TRANSFORM

H. M. SRIVASTAVA

In the present note, we invoke the theories of the Mellin transform as well as fractional integration to investigate a solution of the integral equation

$$(*) \quad \int_0^\infty (xt)^{\sigma-(1/2)} e^{-(1/2)xt} W_{k+(1/2), m}(xt) f(t) dt = W_{k, m}^{(\sigma)}\{f: x\},$$

$x > 0,$

which defines a generalized Whittaker transform of the unknown function $f \in L_2(0, \infty)$ to be determined in terms of its image $W_{k, m}^{(\sigma)}\{f: x\}$.

It is shown that under certain constraints (*) can be reduced to the form of a Laplace integral which is readily solvable by familiar techniques.

Two well-known generalizations of the classical Laplace transform (cf., e.g., [12])

$$(1) \quad L[f: x] = \int_0^\infty e^{-xt} f(t) dt, \quad x > 0,$$

are due to Meijer [6] and Varma [11]. The object of the present note is to investigate a solution of the integral equation

$$(2) \quad \int_0^\infty (xt)^{\sigma-(1/2)} e^{-(1/2)xt} W_{k+(1/2), m}(xt) f(t) dt = W_{k, m}^{(\sigma)}\{f: x\}, \quad x > 0,$$

which defines a generalized Whittaker transform [5, p. 23] of the unknown function $f(t) \in L_2(0, \infty)$ to be determined in terms of its image $W_{k, m}^{(\sigma)}\{f(t): x\}$, so that by appropriately specializing the parameter σ our results would readily enable us to invert the integral transforms of Meijer (cf., [1], [7]) and Varma (cf., [8], [9]).

In what follows we shall make a free use of the existing theories of (i) fractional integration due to Kober [4] and Erdélyi [2], and (ii) the Mellin transform detailed in [10, p. 94]. In the familiar notation, the operator of fractional integration that we need in our analysis is defined as follows:—

$$(3) \quad K_{\zeta, \alpha, n}^{(-)} f(x) = \frac{n}{\Gamma(\alpha)} x^\zeta \int_x^\infty (u^n - x^n)^{\alpha-1} u^{-\zeta-n\alpha+n-1} f(u) du,$$

where $f \in L_p(0, \infty)$, $p^{-1} + q^{-1} = 1$, if $1 < p < \infty$, and q^{-1} or $p^{-1} = 0$

according as p or $q = 1; \alpha > 0, n > 0, \zeta > -p^{-1}$.

Confining ourselves to the L_2 -space theory, for simplicity of the conditions involved, and invoking Fox's lemma (see [3], p. 458), we can establish the following theorems in the usual manner.

THEOREM 1. *Let $f \in L_2(0, \infty)$ be a solution of the integral equation (2). Then*

$$(4) \quad f(x) = L^{-1}[K_{\sigma-k, \sigma+k, 1}^{(-)} W_{k, m}^{(\sigma)}\{f: x\}],$$

provided (i) $x > 0$, (ii) $\sigma + k \geq 0$ and (iii) $1/2 + \sigma - k > 0$.

THEOREM 2. *Let $f(x)$ be a solution of (1) that belongs to $L_2(0, \infty)$. Then*

$$(5) \quad K_{\sigma+m, \alpha, 1}^{(-)} x^{\sigma-m} L[t^{\sigma-m} f(t): x] = W_{-m-\alpha, m}^{(\sigma)}\{f: x\},$$

provided (i) $x > 0$, (ii) $\alpha \geq 0$ and (iii) $1/2 + \sigma + m > 0$.

It may be of interest to remark here that when $\alpha = -k - m$, (5) is reduced to the interesting relationship

$$(6) \quad W_{k, m}^{(\sigma)}\{f: x\} = K_{\sigma+m, -k-m, 1}^{(-)} x^{\sigma-m} L[t^{\sigma-m} f(t): x],$$

which leads us to the construction of a table of generalized Whittaker transforms from that of the classical Laplace transform, provided $k + m \leq 0, 1/2 + \sigma + m > 0$ and $x > 0$.

The author wishes to thank Professor Richard F. Arens and the referee for helpful suggestions.

REFERENCES

1. A. Erdélyi, *On a generalization of the Laplace transformation*, Proc. Edinburgh Math. Soc. (2) **10** (1951), 53-55.
2. ———, *On some functional transformations*, Rend. Sem. Mat. Univ. e Politec. Torino **10** (1950-51), 217-234.
3. C. Fox, *An inversion formula for the kernel $K_\nu(x)$* , Proc. Cambridge Philos. Soc. **61** (1965), 457-467.
4. H. Kober, *On fractional integrals and derivatives*, Quart. J. Math. (Oxford Ser.) **11** (1940), 193-211.
5. V. P. Mainra, *A new generalization of the Laplace transform*, Bull. Calcutta Math. Soc. **53** (1961), 23-31.
6. C. S. Meijer, *Eine neue Erweiterung der Laplace-Transformation*, Proc. Kon. Nederl. Akad. Wetensch. **44** (1941), 727-737 and 831-839.
7. K. M. Saxena, *Inversion and representation theorems for a generalization of Laplace transformation*, Nieuw Arch. v Wisk. (3) **6** (1958), 1-9.
8. ———, *Inversion and representation theorems for a generalized Laplace integral*, Pacific J. Math. **8** (1958), 597-607.

9. R. K. Saxena, *An inversion formula for the Varma transform*, Proc. Cambridge Philos. Soc. **62** (1966), 467-471.
10. E. C. Titchmarsh, *An Introduction to the Theory of Fourier Integrals*, The Oxford University Press, London, 1937.
11. R. S. Varma, *On a generalization of Laplace integral*, Proc. Nat. Acad. Sci. India (A) **20** (1951), 209-216.
12. D. V. Widder, *The Laplace Transform*, Princeton Univ. Press, Princeton, 1941.

Received November 27, 1967.

WEST VIRGINIA UNIVERSITY
MORGANTOWN, WEST VIRGINIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN

Stanford University
Stanford, California

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

J. P. JANS

University of Washington
Seattle, Washington 98105

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California 90024.

Each author of each article receives 50 reprints free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners of publishers and have no responsibility for its content or policies.

Seymour Bachmuth and Horace Yomishi Mochizuki, <i>Kostrikin's theorem on Engel groups of prime power exponent</i>	197
Paul Richard Beesack and Krishna M. Das, <i>Extensions of Opial's inequality</i>	215
John H. E. Cohn, <i>Some quartic Diophantine equations</i>	233
H. P. Dikshit, <i>Absolute $(C, 1) \cdot (N, p_n)$ summability of a Fourier series and its conjugate series</i>	245
Raouf Doss, <i>On measures with small transforms</i>	257
Charles L. Fefferman, <i>L_p spaces over finitely additive measures</i>	265
Le Baron O. Ferguson, <i>Uniform approximation by polynomials with integral coefficients. II</i>	273
Takashi Ito and Thomas I. Seidman, <i>Bounded generators of linear spaces</i>	283
Masako Izumi and Shin-ichi Izumi, <i>Nörlund summability of Fourier series</i>	289
Donald Gordon James, <i>On Witt's theorem for unimodular quadratic forms</i>	303
J. L. Kelley and Edwin Spanier, <i>Euler characteristics</i>	317
Carl W. Kohls and Lawrence James Lardy, <i>Some ring extensions with matrix representations</i>	341
Ray Mines, III, <i>A family of functors defined on generalized primary groups</i>	349
Louise Arakelian Raphael, <i>A characterization of integral operators on the space of Borel measurable functions bounded with respect to a weight function</i>	361
Charles Albert Ryavec, <i>The addition of residue classes modulo n</i>	367
H. M. (Hari Mohan) Srivastava, <i>Fractional integration and inversion formulae associated with the generalized Whittaker transform</i>	375
Edgar Lee Stout, <i>The second Cousin problem with bounded data</i>	379
Donald Curtis Taylor, <i>A generalized Fatou theorem for Banach algebras</i>	389
Bui An Ton, <i>Boundary value problems for elliptic convolution equations of Wiener-Hopf type in a bounded region</i>	395
Philip C. Tonne, <i>Bounded series and Hausdorff matrices for absolutely convergent sequences</i>	415