

Pacific Journal of Mathematics

BASES IN HILBERT SPACE

WILLIAM JAY DAVIS

BASES IN HILBERT SPACE

WILLIAM J. DAVIS

A sequence (x_i) of elements of a Hilbert space, \mathcal{H} , is a *basis* for \mathcal{H} if every $h \in \mathcal{H}$ has a unique, norm-convergent expansion of the form $h = \sum a_i x_i$, where (a_i) is a sequence of scalars. The sequence is *minimal* if there exists a sequence $(y_i) \subset \mathcal{H}$ such that $(x_i, y_j) = \delta_{ij}$. Every basis is minimal, and the sequence (a_i) in the expansion of h (above) is given by $a_i = (h, y_i)$. In this paper, we restrict our attention to real Hilbert space.

We derive, from classical characterizations of bases in B -spaces, criteria for (x_i) to be a basis for \mathcal{H} , as well as for (x_i) to be minimal in \mathcal{H} . We show that the sequence is minimal if and only if there are sequences $(g_i) \subset \mathcal{H}$ whose Gram matrices have a prescribed form. Similar conditions are obtained for (x_i) to be a basis for \mathcal{H} .

Let (x_i) be a linearly independent sequence of elements of \mathcal{H} . Using the Gram-Schmidt process, one finds an orthonormal basis, (w_i) , for the closed span, $[x_i]$ of the sequence (x_i) . We assume throughout that $[x_i] = \mathcal{H}$. Then, we may write

$$x_i = \sum_{j=0}^i p_{ij} w_j,$$

and

$$w_i = \sum_{j=0}^i q_{ij} x_j.$$

If we let P and Q denote the matrices (p_{ij}) and (q_{ij}) , respectively, then each is lower triangular, and $PQ = QP = I = (\delta_{ij})$. It is a classical result that Q is the unique inverse of P .

For (x_i) to be minimal, we need a sequence (y_i) such that $(x_i, y_j) = \delta_{ij}$. It is easy to see that, formally, $y_i = \sum_{j=i}^{\infty} q_{ji} w_j$. Further, the sequence is minimal if and only if the distance from x_k to $[x_j]$, $j \neq k$ is positive. Using these facts, we get the following theorem. The second part is similar to the characterization of minimality due to Foias and Singer [2].

THEOREM 1. *Let $H = (h_{ij})$ denote the Gram matrix of (x_i) , i.e., $h_{ij} = (x_i, x_j)$. Then the sequence is minimal if and only if any of the following conditions holds:*

- (a) *The matrix $R = Q^T Q$ exists.*
- (b) *There exists a sequence, (δ_i) , with $\delta_i > 0$ for all i , such*

that for all real vectors $A = (a_0, a_1, \dots, a_n, 0, \dots)$, $AHA^T \geq \sum \delta_i a_i^2$.

(c) There exists a sequence (ε_i) with $\varepsilon_i > 0$ for all i such that $ARA^T \geq \sum \varepsilon_i a_i^2$, with A as in (b).

Proof. (a) Follows from the formal relation $y_i = \sum q_j w_j$. For (b), notice that $AHA^T \geq \|\sum a_i x_i\|^2$. If (x_i) is minimal, then $AHA^T \geq \lambda_i \|x_i\|^2 a_i^2$, where $\lambda_i^{1/2}$ is the distance from $x_i/\|x_i\|$ to $[x_j]$, $j \neq i$. Therefore, for each permutation (n_i) of the nonnegative integers,

$$AHA^T \geq \sum a^{-(n_i+1)} \lambda_i a_i^2 \|x_i\|^2 .$$

So $\delta_i = 2^{-(n_i+1)} \lambda_i \|x_i\|^2$ works. On the other hand, if $AHA^T \geq \sum \delta_i a_i^2$, then $AHA^T \geq \delta_i a_i^2 = \lambda_i \|x_i\|^2 a_i^2$ for each i . Part (c) follows since (y_i) is minimal if and only if (x_i) is minimal.

2. Here we derive further criteria, for minimal and basic sequences, which depend upon the existence of certain Gram matrices. First, we recall that a fundamental sequence (x_i) in a B -space is minimal if and only if, for each n , there exists a constant $K_n \geq 1$ such that, for all m and all sequences (a_j) ,

$$\left\| \sum_{j=0}^n a_j x_j \right\| \leq K_n \left\| \sum_{j=0}^{n+m} a_j x_j \right\| .$$

Further, such a sequence is basic if and only if (K_n) is bounded (that is, if and only if a bounded sequence (K_n) can be chosen) [1]. In either case, K_n is to be chosen in such a way that

$$\left\{ K_n^2 \left\| \sum_{j=0}^{n+m} a_j x_j \right\|^2 - \left\| \sum_{j=0}^n a_j x_j \right\|^2 \right\}$$

defines a positive definite form on the collection of all finite real sequences. Associated with this form is the matrix $S = S(n, K_n)$, defined as follows:

$$S_{ij} = \begin{cases} (K_n^2 - 1)(x_i, x_j); & 1 \leq i, j \leq n \\ K_n^2(x_i, x_j); & \text{otherwise} . \end{cases}$$

The positive definiteness of the form ASA^T will be achieved over the finite vectors $A = (a_1, a_2, \dots, a_n, 0, \dots)$ if and only if each principal $k \times k$ submatrix, $S^{(k)}$ of S is positive definite. Each $S^{(k)}$ is positive definite if and only if there exists a real, nonsingular, lower triangular matrix T such that $S^{(k)} = T^{(k)} T^{(k)T}$. A routine calculation shows that

$$T_{ij} = \begin{cases} \sqrt{K_n^2 - 1} p_{ij}; & 1 \leq i, j \leq n \\ \frac{K_n^2}{\sqrt{K_n^2 - 1}} p_{ij}; & i > n, 1 \leq j \leq n . \end{cases}$$

Thus, we must solve, in the reals, the equations

$$\sum_{j=n+1}^i T_{ij} T_{kj} = K_n^2(x_i, x_k) - \frac{K_n^4}{K_n^2 - 1} (\pi_n x_i, x_k),$$

where $\pi_n x_i = \sum_{j=1}^n p_{ij} w_j$. If these equations are solvable, then S is positive definite (over finite A), if and only if $T_{ii} \neq 0$. Now let $(f_i)_{i=n+1}^\infty$ be any linearly independent sequence in \mathcal{H} for which

$$(f_i, f_j) = K_n^2(x_i, x_j) - \left(\frac{K_n^4}{K_n^2 - 1} \right) (\pi_n x_i, x_j),$$

if it exists. If we orthonormalize (f_i) , we get a sequence $(g_i)_{i=n+1}^\infty$ and

$$f_i = \sum_{j=n+1}^i T_{ij} g_j.$$

Linear independence of (f_i) gives $T_{ii} \neq 0$. On the other hand, if the equations above are solvable, for (T_{ij}) , we may set $f_i = \sum_{j=n+1}^i T_{ij} w_j$.

We have the following theorem:

THEOREM 2. *The sequence (x_i) is*

(a) *minimal if and only if, for each n , there exists $K_n \geq 1$ and a linearly independent sequence $(f_i)_{i=n+1}^\infty$ such that*

$$(f_i, f_j) = K_n^2(x_i, x_j) - \frac{K_n^4}{K_n^2 - 1} (\pi_n x_i, x_j).$$

(b) *a basis if and only if it is minimal, and the sequence (K_n) may be chosen so that it is bounded.*

The sequence (x_j) is minimal if and only if, for each n , there exists $C_n \geq 1$ such that, for all m and sequences (a_i) ,

$$\left\| \sum_{i=n+1}^{n+m} a_i x_i \right\| \leq C_n \left\| \sum_{i=1}^{n+m} a_i x_i \right\|.$$

It is basic if and only if (C_n) may be chosen as a bounded sequence (see, e.g., [4]). Using these facts, and arguments similar to those for Theorem 2, we obtain,

THEOREM 3. *The sequence (x_i) is*

(a) *minimal if and only if, for each n , there exists $C_n \geq 1$ and a linearly independent sequence $(g_i)_{i=n+1}^\infty$ such that, for $i, j > n$,*

$$(g_i, g_j) = (C_n^2 - 1)(x_i, x_j) - C_n^2(\pi_n x_i, x_j) ,$$

and

(b) *basic if and only if it is minimal and (C_n) may be chosen as a bounded sequence.*

In deriving Theorem 3, one must determine the positive definiteness of the matrices S defined by

$$S_{ij} = \begin{cases} C_n^2(x_i, x_j); & 1 \leq i \leq n \text{ or } 1 \leq j \leq n \\ (C_n^2 - 1)(x_i, x_j); & i, j > n . \end{cases}$$

An interesting characterization of minimal sequences and bases is the following.

PROPOSITION. *The sequence (x_i) is*

(a) *minimal if its Gram matrix, H , is strictly diagonally dominant, and*

(b) *a basis if its Gram matrix is uniformly diagonally dominant.*¹

Proof. If H is strictly diagonally dominant, for each n there exists $\gamma_n \in (0, 1)$ such that $\gamma_n | (x_n, x_n) | < \sum_{j \neq n} | (x_n, x_j) |$. Then, for $C_n^2 = 1/\gamma_n$, the matrix S is strictly diagonally dominant, and hence positive definite over finite $A[5]$. Part (b) follows in the same manner.

Using the same method of proof, Theorems 3 and 4, and the fact that the positive definite $n \times n$ matrices define a cone in the linear space of all $n \times n$ matrices, we obtain the most general form of our characterization of minimal sequences and bases in \mathcal{H} .

THEOREM 4. *The sequence (x_i) is*

(a) *minimal if and only if, some (and hence all) $\alpha, \beta > 0$ and all n , there exist $K_n, C_n \geq 1$ and $(g_i)_{i=n+1}^\infty$ such that, for $i, j > n$,*

$$(g_i, g_j) = (\alpha K_n^2 + \beta C_n^2 - \beta)(x_i, x_j) - \left(\frac{(\alpha K_n^2 + \beta C_n^2)^2}{\alpha K_n^2 + \beta C_n^2 - \alpha} \right) (\pi_n x_i, x_j) ,$$

and

A symmetric matrix A is *strictly diagonally dominant* [5] if $|a_{ii}| > \sum_{j \neq i} |a_{ij}|$ for all i , and is *uniformly diagonally dominant* if there exists $\gamma \in (0, 1)$ such that $\gamma |a_{ii}| \geq \sum_{j \neq i} |a_{ij}|$ for each i .

(b) *a basis if and only if (K_n) and (C_n) may be chosen as bounded sequences.*

REFERENCES

1. S. Banach, *Theorie des operations lineaires*, Warsaw, 1932.
2. C. Foias and I. Singer, *Some remarks on strongly linearly independent sequences and bases in Banach spaces*, *Revue de math pures et appl.* **6** (1961), 589-594.
3. B. Kacymarz and H. Steinhaus, *Theorie der orthogonalreihen*, Warsaw, 1935.
4. I. Singer, *Baze in spatiu Banach, I*, *Studii si Cercetari Matematice*, R. P. R. **14** (1963), 533-585.
5. R. Varga, *Matrix iterative analysis*, Princeton, N. J., 1962.

Received December 13, 1966. This work was supported by National Science Foundation Grant number GP-6152.

THE OHIO STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN

Stanford University
Stanford, California

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS

University of Washington
Seattle, Washington 98105

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California 90024.

Each author of each article receives 50 reprints free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners of publishers and have no responsibility for its content or policies.

Leonard Asimow, <i>Universally well-capped cones</i>	421
Lawrence Peter Belluce, William A. Kirk and Eugene Francis Steiner, <i>Normal structure in Banach spaces</i>	433
William Jay Davis, <i>Bases in Hilbert space</i>	441
Larry Lee Dornhoff, <i>p-automorphic p-groups and homogeneous algebras</i>	447
William Grady Dotson, Jr. and W. R. Mann, <i>A generalized corollary of the Browder-Kirk fixed point theorem</i>	455
John Brady Garnett, <i>On a theorem of Mergelyan</i>	461
Matthew Gould, <i>Multiplicity type and subalgebra structure in universal algebras</i>	469
Marvin D. Green, <i>A locally convex topology on a preordered space</i>	487
Pierre A. Grillet and Mario Petrich, <i>Ideal extensions of semigroups</i>	493
Kyong Taik Hahn, <i>A remark on integral functions of several complex variables</i>	509
Choo Whan Kim, <i>Uniform approximation of doubly stochastic operators</i>	515
Charles Alan McCarthy and L. Tzafriri, <i>Projections in \mathcal{L}_1 and \mathcal{L}_∞-spaces</i>	529
Alfred Berry Manaster, <i>Full co-ordinals of RETs</i>	547
Donald Steven Passman, <i>p-solvable doubly transitive permutation groups</i>	555
Neal Jules Rothman, <i>An L^1 algebra for linearly quasi-ordered compact semigroups</i>	579
James DeWitt Stein, <i>Homomorphisms of semi-simple algebras</i>	589
Jacques Tits and Lucien Waelbroeck, <i>The integration of a Lie algebra representation</i>	595
David Vere-Jones, <i>Ergodic properties of nonnegative matrices. II</i>	601
Donald Rayl Wilken, <i>The support of representing measures for $R(X)$</i>	621
Abraham Zaks, <i>Simple modules and hereditary rings</i>	627