

Pacific Journal of Mathematics

A GENERALIZED COROLLARY OF THE BROWDER-KIRK FIXED POINT THEOREM

WILLIAM GRADY DOTSON, JR. AND W. R. MANN

A GENERALIZED COROLLARY OF THE BROWDER-KIRK FIXED POINT THEOREM

By W. G. DOTSON, JR. AND W. ROBERT MANN

This paper generalizes a corollary, due to W. A. Kirk, of the F. E. Browder-W. A. Kirk fixed point theorem for non-expansive self-mappings of closed, bounded, convex sets in uniformly convex Banach spaces.

F. E. Browder [1] and W. A. Kirk [4] have independently proved that if F is a closed, bounded, convex subset of a uniformly convex Banach space, and if T is a nonexpansive mapping from F into F , then T has a fixed point in F . The following corollary was proved by Kirk [4] and also by Browder and Petryshyn [2]: If E is a uniformly convex Banach space, and $T: E \rightarrow E$ is a nonexpansive mapping, and if for some $x_1 \in E$ the sequence $\{T^n x_1\}$ of Picard iterates of T is bounded, then T has a fixed point in E . Browder and Petryshyn also observed that if the nonexpansive mapping T has a fixed point in E , then for any $x_1 \in E$ the sequence $\{T^n x_1\}$ will be bounded. Outlaw and Groetsch [6] have recently announced the following extension of this corollary: If E is a uniformly convex Banach space, and $T: E \rightarrow E$ is a nonexpansive mapping, and $S_\lambda = \lambda I + (1 - \lambda)T$ for a given λ , $0 < \lambda < 1$, then T has a fixed point in E if and only if the sequence $\{S_\lambda^n x_1\}$ of Picard iterates of S_λ is bounded for each $x_1 \in E$. The purpose of this note is to show that this corollary and its extension are both special cases of a considerably more general corollary of the Browder-Kirk theorem.

W. R. Mann [5] introduced the following general iterative process: Suppose $A = [a_{np}]$ is an infinite real matrix satisfying (1) $a_{np} \geq 0$ for all n, p , and $a_{np} = 0$ for $p > n$; (2) $\sum_{p=1}^n a_{np} = 1$ for each n ; (3) $\lim_n a_{np} = 0$ for each p . If F is a closed convex subset of a Banach space E , and $T: F \rightarrow F$ is a continuous mapping, and $x_1 \in F$, then the process $M(x_1, A, T)$ is defined by

$$v_n = \sum_{p=1}^n a_{np} x_p, \quad x_{n+1} = T v_n, \quad n = 1, 2, 3, \dots$$

Various choices of the matrix A yield many interesting iterative processes as special cases. With A the infinite identity matrix, one gets the Picard iterates of $T: v_{n+1} = x_{n+1} = T v_n$, whence $v_{n+1} = T^n v_1 = T^n x_1$. With $0 < \lambda < 1$ and $A = [a_{np}]$ defined by $a_{np} = \lambda^{n-1}$ if $p = 1$, $a_{np} = \lambda^{n-p}(1 - \lambda)$ if $1 < p \leq n$, $a_{np} = 0$ if $p > n$, $n = 1, 2, 3, \dots$, one gets $v_{n+1} = \lambda v_n + (1 - \lambda)T v_n = S_\lambda v_n$, whence $v_{n+1} = S_\lambda^n v_1 = S_\lambda^n x_1$. If T is linear then an appropriate choice of A yields

$$v_{n+1} = (x_1 + Tx_1 + \dots + T^n x_1)/(n + 1) ,$$

thus providing a connection with mean ergodic theorems for linear operators. Another choice of A yields an iterative process recently investigated by Halpern [3], provided $x_1 = 0$. Many other choices are possible, of course. Our main theorem is as follows.

THEOREM 1. *If E is a uniformly convex Banach space, and if $T: E \rightarrow E$ is a nonexpansive mapping, and if there exist $x_1 \in E$ and a process $M(x_1, A, T)$ such that either of the sequences $\{x_n\}, \{v_n\}$ is bounded, then T has a fixed point in E .*

To prove this, we will make use of the following lemma which is a straightforward consequence of uniform convexity.

LEMMA 1. *Suppose E is a uniformly convex Banach space, and suppose $r > 0$. For each $\varepsilon > 0$ let $p_\varepsilon = \sup \{s : s = \|u - v\| \text{ where } u, v \in E, \|u\| = 2r, 2r < \|v\| \leq 2r + \varepsilon, \text{ and } \|(1 - t)u + tv\| > 2r \text{ for all } t \in (0, 1)\}$. Given any $c > 0$, there exists $\varepsilon > 0$ such that $p_\varepsilon \leq c$.*

Proof of Theorem 1. We first observe that if either of the sequences $\{x_n\}, \{v_n\}$ in the process $M(x_1, A, T)$ is bounded, then the other is also bounded. For if $\|x_n\| \leq b$ for all n , then

$$\|v_n\| = \left\| \sum_{p=1}^n \alpha_{np} x_p \right\| \leq \sum_{p=1}^n \alpha_{np} \|x_p\| \leq b \sum_{p=1}^n \alpha_{np} = b$$

for all n ; and if $\|v_n\| \leq b$ for all n , then

$$\|x_{n+1} - T(0)\| = \|T(v_n) - T(0)\| \leq \|v_n - 0\| \leq b$$

for all n . So, given $x_1 \in E$ and a process $M(x_1, A, T)$ in which both of the sequences $\{x_n\}, \{v_n\}$ are bounded, we wish to show that T has a fixed point. This will be done by showing that T maps a certain bounded, closed, convex set into itself. We use the notation $D_r(p) = \{x : \|x - p\| \leq r\}$, $r > 0, p \in E$. Let $r > 0$ be such that $x_n \in D_r(0)$ and $v_n \in D_r(0)$ for all n . For each $i = 1, 2, 3, \dots$, define sets C_i and G_i by

$$C_i = \bigcap_{n=i}^{\infty} D_{2r}(x_n), \quad G_i = \bigcap_{n=i}^{\infty} \{D_{2r}(x_n) \cap D_{2r}(v_n)\} .$$

For each i , we have

$$D_r(0) \subset G_i \subset C_i \subset D_{2r}(x_i) \subset D_{3r}(0) .$$

Each C_i and each G_i is a nonempty bounded, closed, convex set, and it is clear that $C_i \subset C_{i+1}$ and $G_i \subset G_{i+1}$. We now show $T(G_i) \subset C_{i+1}$: $x \in G_i$ implies $\|x - v_n\| \leq 2r$ for all $n \geq i$, which gives $\|Tx - Tv_n\| \leq$

$\|x - v_n\| \leq 2r$ for all $n \geq i$; but, since $x_{n+1} = Tv_n$, this can be written $\|Tx - x_{n+1}\| \leq 2r$ for all $n \geq i$, so that $Tx \in C_{i+1}$. Define sets C and G by

$$C = \bigcup_{i=1}^{\infty} C_i, \quad G = \bigcup_{i=1}^{\infty} G_i .$$

Clearly, $D_r(0) \subset G \subset C \subset D_{3r}(0)$; and \bar{G}, \bar{C} are bounded, closed, convex sets. Since $T(G_i) \subset C_{i+1}$ for each i , we have $T(G) \subset C$. Since T is continuous, $T(\bar{G}) \subset \overline{T(\bar{G})} \subset \bar{C}$. The proof will be completed by showing $C \subset \bar{G}$, so that $T(\bar{G}) \subset \bar{C} \subset \bar{G}$ (i.e., T maps the bounded, closed, convex set \bar{G} into itself). Since $C = \bigcup_{i=1}^{\infty} C_i$, it suffices to show that for each $i, C_i \subset \bar{G}$. Suppose i is a given positive integer, and $x \in C_i$. We wish to show that $x \in \bar{G}$. The first step toward this end is set off as the following lemma.

LEMMA 2. *For each $\varepsilon > 0$ there exists a positive integer $j_\varepsilon \geq i$ such that $x \in \bigcap_{n=j_\varepsilon}^{\infty} \{D_{2r}(x_n) \cap D_{2r+\varepsilon}(v_n)\} = F_{j_\varepsilon}$.*

Proof of Lemma 2. Since $x \in C_i$ we have $\|x - x_p\| \leq 2r$ for all $p \geq i$. For all $n \geq i$ we have

$$\|x - v_n\| = \left\| \sum_{p=1}^n a_{np}x - \sum_{p=1}^n a_{np}x_p \right\| = \left\| \sum_{p=1}^n a_{np}(x - x_p) \right\|$$

so that

$$\|x - v_n\| \leq \sum_{p=1}^n a_{np} \|x - x_p\| = \sum_{p=1}^{i-1} a_{np} \|x - x_p\| + \sum_{p=i}^n a_{np} \|x - x_p\| ,$$

whence, for all $n \geq i$,

$$\|x - v_n\| \leq \left(\sum_{p=1}^{i-1} a_{np} \right) \cdot \max_{1 \leq p \leq i-1} \|x - x_p\| + 2r .$$

Since i and x are fixed, and since $\lim_n a_{np} = 0$ for each $p = 1, 2, \dots, i-1$, it is clear that for any $\varepsilon > 0$ there exists a positive integer $j_\varepsilon \geq i$ such that $\|x - v_n\| \leq 2r + \varepsilon$ for all $n \geq j_\varepsilon$. But $n \geq j_\varepsilon \geq i$ also implies $\|x - x_n\| \leq 2r$ since $x \in C_i$. Hence $n \geq j_\varepsilon$ implies

$$x \in D_{2r}(x_n) \cap D_{2r+\varepsilon}(v_n) ,$$

and so $x \in \bigcap_{n=j_\varepsilon}^{\infty} \{D_{2r}(x_n) \cap D_{2r+\varepsilon}(v_n)\} = F_{j_\varepsilon}$.

Proof of Theorem 1 continued. We return now to the final problem of showing $x \in \bar{G}$ (see immediately before Lemma 2). Given any $c > 0$, choose $\varepsilon > 0$ such that $p_\varepsilon \leq c$ (this can be done by Lemma 1, in which $r > 0$ is taken as the r we are using in this proof). For

this ε , there exists a positive integer $j_\varepsilon \geq i$ such that $x \in F_{j_\varepsilon}$ (by Lemma 2). We will show $G_{j_\varepsilon} \cap D_c(x) \neq \phi$. Since c is arbitrary, this will show $x \in \bar{G} = (\bigcup_{i=1}^\infty G_i)^-$. We suppose $G_{j_\varepsilon} \cap D_c(x) = \phi$ and obtain a contradiction. Since $0 \in D_r(0) \subset G_{j_\varepsilon}$, $0 \notin D_c(x)$, and so $0 < c/\|x\| < 1$. Let $t_1 = 1 - (c/\|x\|)$. Then $0 < t_1 < 1$ and $\|t_1x - x\| = (1 - t_1)\|x\| = c$. Since $t_1x \in D_c(x)$, we have $t_1x \notin G_{j_\varepsilon}$. Now $x \in F_{j_\varepsilon} \subset \bigcap_{n=j_\varepsilon}^\infty D_{2r}(x_n) = C_{j_\varepsilon}$, and since C_{j_ε} also contains 0 and is convex, $t_1x \in C_{j_\varepsilon}$. Since $t_1x \notin G_{j_\varepsilon}$ and $t_1x \in C_{j_\varepsilon}$, we have $t_1x \notin \bigcap_{n=j_\varepsilon}^\infty D_{2r}(v_n)$. Let n be a positive integer, $n \geq j_\varepsilon$, such that $t_1x \notin D_{2r}(v_n)$. Let

$$t_2 = \sup \{t: 0 < t < 1 \text{ and } tx \in D_{2r}(v_n)\}.$$

This set of t 's is nonempty since $D_r(0) \subset D_{2r}(v_n)$. Since $D_{2r}(v_n)$ is closed, we have $t_2x \in D_{2r}(v_n)$; and it is easily seen from the definition of t_2 that we must have $\|t_2x - v_n\| = 2r$. If $t_2 \geq t_1$, then, since 0 and t_2x are in the convex set $D_{2r}(v_n)$, we would have $t_1x \in D_{2r}(v_n)$ which is not true. Hence $t_2 < t_1$. Similarly we have $\|x - v_n\| > 2r$, since 0 is in the convex set $D_{2r}(v_n)$ and t_1x is not. Since $x \in F_{j_\varepsilon}$ and since $n \geq j_\varepsilon$, $x \in D_{2r+\varepsilon}(v_n)$, so we have $2r < \|x - v_n\| \leq 2r + \varepsilon$. Next we observe that if $t \in (0, 1)$

$$\|(1 - t)(t_2x - v_n) + t(x - v_n)\| = \|[(1 - t)t_2 + t \cdot 1]x - v_n\| > 2r$$

since $t_2 < (1 - t)t_2 + t \cdot 1 < 1$ so that $[(1 - t)t_2 + t \cdot 1]x \notin D_{2r}(v_n)$. With $u = t_2x - v_n$ and $v = x - v_n$ we now have $\|u\| = 2r$, $2r < \|v\| \leq 2r + \varepsilon$, and $\|(1 - t)u + tv\| > 2r$ for all $t \in (0, 1)$. Hence $\|u - v\| = \|t_2x - x\| \leq p_\varepsilon$ (see Lemma 1). But ε was chosen so that $p_\varepsilon \leq c$. So we have

$$\|t_2x - x\| = (1 - t_2)\|x\| \leq p_\varepsilon \leq c.$$

This gives $t_2 \geq 1 - (c/\|x\|) = t_1$, which is a contradiction.

For completeness, we include the following theorem which is somewhat stronger than the converse of Theorem 1.

THEOREM 2. *If E is a normed linear space, and if $T: E \rightarrow E$ is a nonexpansive mapping, and if T has a fixed point $p \in E$, then for any $x_1 \in E$ and any process $M(x_1, A, T)$, the sequences $\{x_n\}$, $\{v_n\}$ are bounded.*

Proof. For each $n = 1, 2, 3, \dots$, we have

$$\begin{aligned} \|x_{n+1} - p\| &= \|Tv_n - Tp\| \leq \|v_n - p\| = \left\| \sum_{j=1}^n a_{nj}(x_j - p) \right\| \\ &\leq \sum_{j=1}^n a_{nj} \|x_j - p\| \leq \max_{j=1, \dots, n} \|x_j - p\|. \end{aligned}$$

Thus $\|x_2 - p\| \leq \|x_1 - p\|$, $\|x_3 - p\| \leq \max_{j=1,2} \|x_j - p\| = \|x_1 - p\|$, etc., so that we have $\|x_j - p\| \leq \|x_1 - p\|$ for all $j = 1, 2, 3, \dots$; and hence with $b = \|x_1 - p\| + \|p\|$ we get $\|x_j\| = \|(x_j - p) + p\| \leq \|x_j - p\| + \|p\| \leq b$ for all $j = 1, 2, 3, \dots$. Finally,

$$\|v_n\| = \left\| \sum_{j=1}^n a_{nj} x_j \right\| \leq \sum_{j=1}^n a_{nj} \|x_j\| \leq b \cdot \sum_{j=1}^n a_{nj} = b$$

for all $n = 1, 2, 3, \dots$.

BIBLIOGRAPHY

1. F. E. Browder, *Nonexpansive nonlinear operators in a Banach space*, Proc. Nat. Acad. Sci. U.S.A. **54** (1965), 1041-1044.
2. F. E. Browder and W. V. Petryshyn, *The solution by iteration of nonlinear functional equations in Banach spaces*, Bull. Amer. Math. Soc. **72** (1966), 571-575.
3. B. Halpern, *Fixed points of nonexpanding maps*, Bull. Amer. Math. Soc. **73** (1967), 957-961.
4. W. A. Kirk, *A fixed point theorem for mappings which do not increase distances*, Amer. Math. Monthly **72** (1965), 1004-1006.
5. W. Robert Mann, *Mean value methods in iteration*, Proc. Amer. Math. Soc. **4** (1953), 506-510.
6. C. L. Outlaw and C. W. Groetsch, *Averaging iteration in a Banach space*, Notices Amer. Math. Soc. **15** (1968), 180.

Received March 12, 1968.

NORTH CAROLINA STATE UNIVERSITY AND
UNIVERSITY OF NORTH CAROLINA, CHAPEL HILL

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN

Stanford University
Stanford, California

J. DUGUNDJI

Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS

University of Washington
Seattle, Washington 98105

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California 90024.

Each author of each article receives 50 reprints free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners of publishers and have no responsibility for its content or policies.

Leonard Asimow, <i>Universally well-capped cones</i>	421
Lawrence Peter Belluce, William A. Kirk and Eugene Francis Steiner, <i>Normal structure in Banach spaces</i>	433
William Jay Davis, <i>Bases in Hilbert space</i>	441
Larry Lee Dornhoff, <i>p-automorphic p-groups and homogeneous algebras</i>	447
William Grady Dotson, Jr. and W. R. Mann, <i>A generalized corollary of the Browder-Kirk fixed point theorem</i>	455
John Brady Garnett, <i>On a theorem of Mergelyan</i>	461
Matthew Gould, <i>Multiplicity type and subalgebra structure in universal algebras</i>	469
Marvin D. Green, <i>A locally convex topology on a preordered space</i>	487
Pierre A. Grillet and Mario Petrich, <i>Ideal extensions of semigroups</i>	493
Kyong Taik Hahn, <i>A remark on integral functions of several complex variables</i>	509
Choo Whan Kim, <i>Uniform approximation of doubly stochastic operators</i>	515
Charles Alan McCarthy and L. Tzafriri, <i>Projections in \mathcal{L}_1 and \mathcal{L}_∞-spaces</i>	529
Alfred Berry Manaster, <i>Full co-ordinals of RETs</i>	547
Donald Steven Passman, <i>p-solvable doubly transitive permutation groups</i>	555
Neal Jules Rothman, <i>An L^1 algebra for linearly quasi-ordered compact semigroups</i>	579
James DeWitt Stein, <i>Homomorphisms of semi-simple algebras</i>	589
Jacques Tits and Lucien Waelbroeck, <i>The integration of a Lie algebra representation</i>	595
David Vere-Jones, <i>Ergodic properties of nonnegative matrices. II</i>	601
Donald Rayl Wilken, <i>The support of representing measures for $R(X)$</i>	621
Abraham Zaks, <i>Simple modules and hereditary rings</i>	627