

Pacific Journal of Mathematics

ON A THEOREM OF MERGELYAN

JOHN BRADY GARNETT

ON A THEOREM OF MERGELYAN

JOHN GARNETT

The following is a theorem of S. N. Mergelyan.

THEOREM. *If X is a compact plane set with finitely many complementary components, then every continuous function on X which is analytic on the interior is a uniform limit on X of rational functions.*

Here two short proofs of the theorem are given. Both proofs are obtained by considering the measures on X orthogonal to the algebra of rational functions.

Let X be a compact plane set and suppose its complement $C \setminus X$ has finitely many components. In 1951 Mergelyan [9] proved:

MERGELYAN'S THEOREM. *Every function continuous on X and analytic on the interior is uniformly approximable by rational functions.*

By examining the measures on X orthogonal to the rational functions, Glicksberg and Wermer [6] have found an elegant proof of Mergelyan's Theorem in the case $C \setminus X$ connected. Their argument has recently been extended by Ahern and Sarason [1] and by Glicksberg [5] to give functional analytic proofs of the general Mergelyan result. However those proofs are quite long. Here we give two shorter proofs of the theorem. The first proof begins with the known fact that the question of approximation is a local one, and the theorem is thereby reduced to the simply connected case. The second proof follows the reasoning of [6]. Here we use some results from [1] and [5], but avoid the more lengthy arguments in those papers. Though longer than the first, this second proof is included because the ideas therein yield related results not accessible via our first proof.

Denote by $A(X)$ the Banach algebra of functions continuous on X and analytic on X° , the interior. Let $R(X)$ be the closed subalgebra of $A(X)$ spanned by the rational functions analytic on X . By a measure on X we mean a finite complex Borel measure on X . Write $\mu \perp R(X)$ if μ is a measure on X orthogonal to $R(X)$, and $\mu \perp A(X)$ if μ annihilates $A(X)$ as well. As have several previous authors, we prove the Mergelyan Theorem by showing $\mu \perp R(X)$ implies $\mu \perp A(X)$.

2. **The first proof.** We quote the following lemma of Bishop [3, p. 40]. This lemma was stated for polynomials, but Bishop's proof is valid for rational functions.

LEMMA 2.1. (*Bishop*). Let X be a compact subset of \mathbb{C} . For x a real number, set

$$L_x = X \cap \{z: \text{Real } z \leq x\}, \quad R_x = X \cap \{z: \text{Real } z \geq x\}.$$

Let $\mu \perp R(X)$. Then for almost all x there are measures ν_1 on L_x and ν_2 on R_x such that $\nu_1 \perp R(L_x)$, $\nu_2 \perp R(R_x)$ and $\mu = \nu_1 + \nu_2$.

This lemma gives the known result that questions concerning $R(X)$ are local ones.

COROLLARY 2.2. Let f be a continuous function on the compact set X . Assume each point in X has a closed neighborhood K such that the restricted function $f|_K$ is in $R(K)$. Then f is in $R(X)$.

Proof. Let μ be a measure orthogonal to $R(X)$ and let $\{U_1, U_2, \dots, U_n\}$ be any finite open cover of X . It follows from repeated applications of 2.1 that there are finitely many compact rectangles $\{T_1, \dots, T_m\}$ and measures ν_j on T_j such that

$$(i) \quad T_j \subset U_i \text{ for some } i$$

$$(ii) \quad \nu_j \perp R(T_j)$$

$$(iii) \quad \mu = \sum_{j=1}^m \nu_j.$$

If f satisfies the hypotheses, then using such a decomposition we see that $\int f d\mu = 0$. This implies $f \in R(X)$.

Actually we can localize measures as in the proof of 2.2 without appealing to Bishop's lemma. Instead we use an observation shown us by A. Browder, who credits it to K. Hoffman. For a measure μ on X define its *Cauchy transform*

$$\hat{\mu}(\alpha) = \int (z - \alpha)^{-1} d\mu(z).$$

Then $\hat{\mu}$ is locally integrable with respect to plane Lebesgue measure. Also $\hat{\mu} = 0$ almost everywhere if and only if $\mu = 0$. See [11]. The observation, whose proof we only sketch, is this:

(a) $\mu \perp R(X)$ if and only if $\hat{\mu} = 0$ on $\mathbb{C} \setminus X$. This is seen by taking partial fractions and by "pushing poles together."

(b) An application of Green's theorem yields, for φ a C^∞ function with compact support,

$$\varphi \hat{\mu} = \widehat{\varphi \mu} + \hat{\sigma} \quad \text{where} \quad \sigma = \frac{1}{2\pi i} \frac{\partial \varphi}{\partial \bar{z}} \hat{\mu}(z) dz \wedge d\bar{z}.$$

Now if $\mu \perp R(X)$ and K is the closed support of φ , then by (a) and (b),

$$\varphi \mu + \sigma \perp R(X \cap K).$$

Letting φ run through a finite partition of unity $\{\varphi_j\}_{j=1}^m$ and setting $\nu_j = \varphi_j \mu + 1/2\pi i \partial\varphi_j/\partial\bar{z} \hat{\mu}(z) dz \wedge d\bar{z}$ we obtain a decomposition of μ satisfying (i), (ii) and (iii) in the proof of 2.2.

Proof of theorem. Let H_1, H_2, \dots, H_n be the bounded components of $C \setminus X$ and let $f \in A(X)$. For each point in X choose a compact neighborhood K with diameter less than the diameter of each H_j . Then $C \setminus K$ is connected and $f|K$ is in $A(K)$. Thus by [6], $f|K$ is in $R(K)$. Hence f is in $R(X)$ by 2.2.

We note that this proof is still valid when $C \setminus X$ has infinitely many components but the diameters of these components are bounded below. Examples of such sets are easily constructed.

3. **The second proof.** By the maximum modulus principle $A(X)$ can be identified with the algebra of boundary values of functions in $A(X)$. We will view $A(X)$ as a closed algebra of continuous functions in ∂X , the boundary of X , and we consider now only measures supported on $\partial(X)$.

We begin with a theorem of Walsh [10] and Lebesgue [8]. A more recent proof is contained in the first three lemmas of [4]. $C^R(\partial X)$ is the space of real continuous functions on ∂X , and Real $R(X)$ is the space (restrictions to ∂X) of real parts of functions in $R(X)$.

WALSH-LEBESGUE THEOREM. *Let H_1, H_2, \dots, H_n be the bounded components of $C \setminus X$ and let $a_j \in H_j$. Then Real $R(X) \oplus \text{Span} \{ \log |z - a_j| : 1 \leq j \leq n \}$ is uniformly dense in $C^R(\partial X)$.*

This means the space of real measures on ∂X orthogonal to $R(X)$ has dimension at most n . It also means the Dirichlet problem is solvable on X ; if $u \in C^R(\partial X)$ then there is a unique \hat{u} in $C^R(X)$ harmonic on X^0 with $u = \hat{u}$ on ∂X . Hence for $z \in X$ there is a unique probability measure λ_z on ∂X such that $\hat{u}(z) = \int_{\partial X} u d\lambda_z$ for all $u \in C^R(\partial X)$. It follows directly that λ_z is a *representing measure* for $A(X)$, as λ_z is positive and

$$f(z) = \int_{\partial X} f d\lambda_z \quad \text{for all } f \in A(X).$$

It also follows that λ_z is an *Arens-Singer measure* for $A(X)$,

$$\log |f(z)| = \int_{\partial X} \log |f| d\lambda_z \quad \text{for } f \in A(X) \quad \text{and} \quad \frac{1}{f} \in A(X).$$

Moreover the Walsh-Lebesgue theorem implies that λ_z is the unique

Arens-Singer measure for $R(X)$. Consequently the assumptions of the paper [1] of Ahern and Sarason apply to $R(X)$. We now state a lemma of theirs and some immediate consequences. All except Corollary 3.4 are proven on pp. 126–128 of [1].

LEMMA 3.1. (*Ahern-Sarason*). *Let $(v_n)_{n=1}^{\infty}$ be a sequence of non-negative functions in $C^R(\partial X)$ with $\int v_n d\lambda_z \rightarrow 0$. Then there is a subsequence $(u_n)_{n=1}^{\infty}$ and a sequence $(f_n)_{n=1}^{\infty}$ of functions in $R(X)$ such that $|f_n| \leq e^{-u_n}$ and $f_n \rightarrow 1$ almost everywhere λ_z .*

COROLLARY 3.2. (*F. and M. Riesz Theorem*). *If $z \in X$ and $\mu \perp R(X)$, let $\mu = \mu_z + \tilde{\mu}_z$ be the Lebesgue decomposition of μ with respect to λ_z . Then $\mu_z \perp R(X)$ and $\tilde{\mu}_z \perp R(X)$.*

COROLLARY 3.3. *For $z \in X$, every representing measure on ∂X for z is absolutely continuous with respect to λ_z . In particular, when $z \in \partial X$, the only representing measure is the point mass at z .*

The next corollary is a consequence of the F. and M. Riesz theorem, and is Theorem 2.8 of Glicksberg's paper [5].

COROLLARY 3.4. *If $\mu \perp R(X)$, then there is a sequence $(z_n)_{n=1}^{\infty}$ in X such that*

$$(1) \quad \mu = \mu_s + \sum_{n=1}^{\infty} \mu_{z_n}$$

where the series converges in norm, each term is orthogonal to $R(X)$, μ_{z_n} is absolutely continuous with respect to λ_{z_n} and μ_s is singular with respect to all representing measures.

Proof. First take the “simultaneous Lebesgue decomposition of μ with respect to all λ_z .” That is, let $c = \sup \{ \|\mu_z\| : z \in X \}$ and choose z_1 so that $\|\mu_{z_1}\| > c/2$. Take the Lebesgue decomposition $\mu = \tilde{\mu}_{z_1} + \mu_{z_1}$. Repeating the argument with $\tilde{\mu}_{z_1}$ and continuing by induction yields a decomposition (1), in which μ_s is singular with respect to all λ_z .

By Corollary 3.2, $\mu_{z_1} \perp R(X)$, and so by induction every μ_{z_n} (and hence μ_s as well) is orthogonal to $R(X)$. Finally by Corollary 3.3, μ_s is singular with respect to every representing measure.

From Corollary 3.4 we see that to show any $\mu \in (R(X))^\perp$ annihilates $A(X)$ it suffices to consider the case when $\mu = \mu_z$ for some z and the case when μ is singular with respect to all representing measures. The latter case is settled by the following argument of Wilken [12],

valid for any compact plane set.

LEMMA 3.5. (*Wilken*). *Let X be any compact plane set and let $\mu \perp R(X)$. If μ is singular with respect to all representing measures then $\mu = 0$.*

Proof. As in [12], there is a point $z \in X$ such that

$$\int \frac{d|\mu|(\zeta)}{|z - \zeta|} < \infty \quad \text{and} \quad \int \frac{d\mu(\zeta)}{z - \zeta} = c \neq 0.$$

Then the measure $\nu = 1/c \mu(\zeta)/z - \zeta$ is a complex measure representing z ; so that by [7] there is a positive representing measure m for z absolutely continuous with respect to $|\nu|$. Then m is absolutely continuous with respect to $|\mu|$ and the assertion obtains.

For our analysis of the absolutely continuous orthogonal measures we need two more lemmas. For $z \in X$, write $\lambda = \lambda_z$, and let $H^2(R(X))$ be the closure of $R(X)$ in $L^2(\lambda)$. Set $H^\infty(R(X)) = H^2(R(X)) \cap L^\infty(\lambda)$. Define $H^2(A(X))$ and $H^\infty(A(X))$ analogously. The first lemma is from Ahern and Sarason [1] and follows directly from Lemma 3.1 by an argument of Hoffman and Wermer. (See also [5] and [11].)

LEMMA 3.6. *If h is in $H^\infty(R(X))$, then there is a sequence $(h_n)_{n=1}^\infty$ in $R(X)$ with $\|h_n\|_\infty \leq \|h\|_\infty$ and $h_n \rightarrow h$ almost everywhere λ .*

The second lemma is a minor variation of Glicksberg's Lemma 3.16 of [5]. The proof is due to T. W. Gamelin and the author.

LEMMA 3.7. $H^2(R(X)) = H^2(A(X))$.

Proof. Let $\overline{H_0^2(R(X))}$ be the space of complex conjugates of functions in $H^2(R(X))$ having integral zero. Since λ is multiplicative on $R(X)$, $L^2(\lambda)$ has the following orthogonal decomposition:

$$L^2(\lambda) = H^2(R(X)) \oplus \overline{H_0^2(R(X))} \oplus E.$$

The space E is spanned by the real functions in L^2 annihilating $R(X)$ and thus by the Walsh-Lebesgue theorem E is finite dimensional. Also λ is multiplicative on $A(X)$, so an orthogonality argument gives

$$H^2(R(X)) \subset H^2(A(X)) \subset H^2(R(X)) \oplus E.$$

This means $H^2(A(X)) = H^2(R(X)) \oplus N$ where N is finite dimensional. Let G_1, G_2, \dots, G_p be an orthonormal basis of N and for an appropriately small ε choose g_1, g_2, \dots, g_p in $A(X)$ with $\|g_j - G_j\|_2 < \varepsilon$. Let J be the span of $\{g_1, \dots, g_p\}$. Then

$$H^2(A(X)) = H^2(R(X)) + J$$

where the sum is direct but not necessarily orthogonal. Let P_J be the projection of $H^2(A(X))$ onto J which is zero on $H^2(R(X))$. For $f \in R(X)$ and $g \in J$ define $T_f(g) = P_J(fg)$. Then T is a representation of $R(X)$ on J . Since $R(X)$ is commutative and J is finite dimensional there is a common eigenvector $g \in J$. Therefore for $f \in R(X)$ we have

$$(2) \quad fg = \varphi(f) \cdot g + r$$

where $r \in H^2(R(X))$ and $\varphi(f) \in \mathbb{C}$.

Clearly φ is a multiplicative linear functional on $R(X)$, so that there is a point $z_1 \in X$ such that $\varphi(f) = f(z_1)$ for all $f \in R(X)$. Also because g is bounded φ extends continuously to $H^2(R(X))$ and so there is an $F \in L^2(\lambda)$ such that $f(z_1) = \int fF d\lambda$. Hence by (7) z_1 has a positive representing measure absolutely continuous with respect to λ . By Corollary 3.3 then $z_1 \notin \partial X$.

Taking $f = z$ in (2), we have $h = (z - z_1)g \in H^2(R(X))$. Hence there is a sequence $(h_n)_{n=1}^\infty$ in $R(X)$ converging to h in $L^2(\lambda)$. Let $c = \int hF d\lambda = \lim_{n \rightarrow \infty} \int h_n F d\lambda$. Then since $z_1 \in X^0$, $r_n = h_n - h_n(z_1)/z - z_1$ is in $R(X)$ and r_n converges to $g - c(z - z_1)^{-1}$ in $L^2(\lambda)$. Thus

$$(3) \quad g - c(z - z_1)^{-1} \in H^2(R(X)).$$

If $c = 0$, then $g \in H^2(R(X))$ and we are done. On the other hand if $c \neq 0$, then $(z - z_1)^{-1} \in H^2(A(X))$. Being bounded on ∂X , $(z - z_1)^{-1}$ is therefore in $H^\infty(A(X))$. Now $H^\infty(A(X))$ is an algebra, and consequently $(z - z_1)^{-n} \in H^2(A(X))$ for all n . But $H^2(R(X))$ has finite codimension in $H^2(A(X))$; whence some polynomial in $(z - z_1)^{-1}$ is in $H^2(R(X))$. That is

$$\frac{a_1}{z - z_1} + \frac{a_2}{(z - z_1)^2} + \dots + \frac{a_n}{(z - z_1)^n} \in H^2(R(X)). \quad (a_n \neq 0).$$

Multiplying by $(z - z_1)^{n-1}$ we have $(z - z_1)^{-1} \in H^2(R(X))$. By (3), then $g \in H^2(R(X))$. Therefore $J = \{0\}$ and

$$H^2(R(X)) = H^2(A(X)).$$

To conclude the proof of Mergelyan's theorem, let $\mu \perp R(X)$. We can assume μ is absolutely continuous with respect to some λ_z . Let $h \in A(X)$. Then by 3.7 $h \in H^2(R(X))$ and so by 3.6 there is a sequence $(h_n)_{n=1}^\infty$ in $R(X)$ converging boundedly pointwise to h almost everywhere λ_z . As $\int h_n d\mu = 0$, dominated convergence implies $\int h d\mu = 0$ and μ is orthogonal to $A(X)$.

We should remark that the proof of Lemma 3.7 actually shows that $H^\infty(R(X))$ is maximal among the subalgebras of $L^\infty(\lambda)$ on which λ is multiplicative. This means that one can compute the defect of Real $R(X)$ in $C^R(\partial X)$ and obtain the other results that Ahern and Sarason do in [2] without using the deeper theorems of their earlier paper [1].

Added in proof. After a closer reading of the literature we have found that the proof of Mergelyan's theorem in § 2 was essentially given by Laura Kodama in "Boundary measures of analytic differentials" Pacific J. Math., 15 (1965), 1261-1277.

We wish to thank T. W. Gamelin and K. Hoffman for helpful conversations.

REFERENCES

1. P. R. Ahern and Donald Sarason, *The H^p spaces of a class of function of algebras*, Acta. Math. **117** (1967), 123-163.
2. P. R. Ahern and Donald Sarason, *On some hypodichlet algebras of analytic functions*, Amer. J. Math. **89** (1967), 932-941.
3. E. Bishop, *Subalgebras of functions on a Riemann surface*, Pacific J. Math. **8** (1958), 29-50.
4. L. Carleson, *Mergelyan's theorem on uniform polynomial approximation*, Math. Scand. **15** (1964), 167-175.
5. I. Glicksberg, *Dominant representing measures and rational approximation* Trans. Amer. Math. Soc. **130** (1968), 425-462.
6. I. Glicksberg and J. Wermer, *Measures orthogonal to a Dirichlet algebra*, Duke Math. J. **30** (1963), 661-666.
7. K. Hoffman and H. Rossi, *On the extension of positive weak continuous functionals*, Duke Math. J. **34** (1967), 453-466.
8. H. Lebesgue, *Sur le problem de Dirichlet*, Rend. Palermo, **29** (1907), 371-402.
9. S. N. Mergelyan, *Uniform approximation to functions of a complex variable*, Amer. Math. Soc. Translation No. 101.
10. J. L. Walsh, *The approximation of harmonic functions by harmonic polynomials and harmonic rational functions*, Bull. Amer. Math. Soc. **35** (1929), 499-544.
11. J. Wermer, *Seminar über Funktionen-Algebren*, Springer, Berlin, 1964.
12. D. R. Wilken, *Lebesgue measure for parts of $R(X)$* , Proc. Amer. Math. Soc. **18** (1967), 508-512.

Received May 25, 1967. Research sponsored by the Air Force Office of Scientific Research, Office of Aerospace Research, United States Air Force, under AFOSR Grant No. 335-63.

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California 90024.

Each author of each article receives 50 reprints free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners of publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 26, No. 3

BadMonth, 1968

Leonard Asimow, <i>Universally well-capped cones</i>	421
Lawrence Peter Belluce, William A. Kirk and Eugene Francis Steiner, <i>Normal structure in Banach spaces</i>	433
William Jay Davis, <i>Bases in Hilbert space</i>	441
Larry Lee Dornhoff, <i>p-automorphic p-groups and homogeneous algebras</i>	447
William Grady Dotson, Jr. and W. R. Mann, <i>A generalized corollary of the Browder-Kirk fixed point theorem</i>	455
John Brady Garnett, <i>On a theorem of Mergelyan</i>	461
Matthew Gould, <i>Multiplicity type and subalgebra structure in universal algebras</i>	469
Marvin D. Green, <i>A locally convex topology on a preordered space</i>	487
Pierre A. Grillet and Mario Petrich, <i>Ideal extensions of semigroups</i>	493
Kyong Taik Hahn, <i>A remark on integral functions of several complex variables</i>	509
Choo Whan Kim, <i>Uniform approximation of doubly stochastic operators</i>	515
Charles Alan McCarthy and L. Tzafriri, <i>Projections in \mathcal{L}_1 and \mathcal{L}_∞-spaces</i>	529
Alfred Berry Manaster, <i>Full co-ordinals of RETs</i>	547
Donald Steven Passman, <i>p-solvable doubly transitive permutation groups</i>	555
Neal Jules Rothman, <i>An L^1 algebra for linearly quasi-ordered compact semigroups</i>	579
James DeWitt Stein, <i>Homomorphisms of semi-simple algebras</i>	589
Jacques Tits and Lucien Waelbroeck, <i>The integration of a Lie algebra representation</i>	595
David Vere-Jones, <i>Ergodic properties of nonnegative matrices. II</i>	601
Donald Rayl Wilken, <i>The support of representing measures for $R(X)$</i>	621
Abraham Zaks, <i>Simple modules and hereditary rings</i>	627