

# Pacific Journal of Mathematics

**COMMUTATIVITY THEOREMS FOR NONASSOCIATIVE  
RINGS WITH A FINITE DIVISION RING HOMOMORPHIC  
IMAGE**

EUGENE CARLYLE JOHNSEN, DAVID LEWIS OUTCALT  
AND ADIL MOHAMED YAQUB

## COMMUTATIVITY THEOREMS FOR NONASSOCIATIVE RINGS WITH A FINITE DIVISION RING HOMOMORPHIC IMAGE

E. C. JOHNSEN, D. L. OUTCALT, AND ADIL YAQUB

Wedderburn's Theorem, asserting that a finite associative division ring is necessarily commutative, has been extended to

**THEOREM 1.** Let  $R$  be a noncommutative Jordan ring of characteristic not 2, and let  $I$  be an ideal in  $R$  such that  $R/I$  is a finite division ring of characteristic  $p > 5$  with exactly  $q$  elements. Suppose that (i)  $I$  is commutative and every associator contained in the ideal generated by  $I^2$  vanishes, and (ii)  $x \equiv y \pmod{I}$  implies  $x^q = y^q$  or both  $x$  and  $y$  commute with all elements of  $I$ . Then  $R$  is commutative.

The object of this paper is to extend Theorem 1 in two directions. First we replace the assumption that  $R$  is a noncommutative Jordan ring by the weaker assumption that  $R$  is power-associative. Next we assume that  $R$  is a flexible power-associative ring but replace the hypothesis that every associator in the ideal generated by  $I^2$  vanishes with the weaker assumption that  $I$  is associative. In each case we drop the assumption that  $R$  is of characteristic not 2.

The proof of Theorem 1 appears in [2].

By a noncommutative Jordan ring is meant a ring in which the associative law is replaced by the weaker identities

$$(1.1) \quad (x, y, x) = 0,$$

and

$$(1.2) \quad (x^2, y, x) = 0;$$

where the associator  $(a, b, c)$  is defined by  $(a, b, c) = (ab)c - a(bc)$ . A ring is flexible in case only (1.1) is assumed, and a ring is power-associative provided

$$(1.3) \quad x^m x^n = x^{m+n}$$

holds in the ring for all positive integers  $m, n$ . It is known that a noncommutative Jordan ring of characteristic not 2 is power-associative [4], but there are flexible rings which are not power-associative. A ring  $R$  is said to be of characteristic not 2 if  $2x = 0$  implies  $x = 0$  in  $R$ .

### 2. Main results.

**THEOREM 2.** *Let  $R$  be a power-associative ring and let  $I$  be an ideal in  $R$  such that  $R/I$  is a finite division ring of characteristic  $p > 5$  with exactly  $q$  elements. Suppose that (i)  $I$  is commutative and every associator in the ideal generated by  $I^2$  vanishes, and (ii)  $x \equiv y \pmod{I}$  implies  $x^q = y^q$  or both  $x$  and  $y$  commute with all elements of  $I$ . Then  $R$  is commutative.*

*Proof.* We first note that since  $R/I$  is a finite power-associative division ring of characteristic  $p > 5$ ,  $R/I$  is a finite field [1; Th. 5]. Hence for every  $\bar{x} \in R/I$ ,  $\bar{x}^q = \bar{x}$ , whence for every  $x \in R$ ,  $x^q \equiv x \pmod{I}$ . Now let  $a_0 \in I, b \in R$ . We first wish to show that  $a_0 b = b a_0$ . Suppose not. Let  $a \in I$ . Since  $b + a \equiv b \pmod{I}$  and  $a_0 b \neq b a_0$ , we have that

$$(2.1) \quad (b + a)^q = b^q .$$

Now by the power-associativity of  $R$

$$(2.2) \quad ((b + a)^{q^l}, (b + a)^{q^m}, (b + a)) = 0, \quad l, m \text{ positive integers.}$$

Hence, by (2.1),  $(b^{q^l}, b^{q^m}, b + a) = 0$ , whence

$$(2.3) \quad (b^{q^l}, b^{q^m}, a) = 0 .$$

Similarly

$$(2.4) \quad (b^{q^l}, a, b^{q^m}) = (a, b^{q^l}, b^{q^m}) = 0 .$$

Since  $b^{q^l} \equiv b^l \pmod{I}$  and  $b^{q^m} \equiv b^m \pmod{I}$ , (2.3) and the vanishing of every associator in the ideal generated by  $I^2$  imply

$$(2.5) \quad (b^l, b^m, a) = 0 .$$

Similarly, from (2.4),

$$(2.6) \quad (a, b^l, b^m) = (b^l, a, b^m) = 0 .$$

We now show that the subring  $R_{a_0, b}$  of  $R$  generated by  $a_0$  and  $b$  is associative. It is sufficient to show that

$$(2.7) \quad (\langle a_0, b \rangle_1 \langle a_0, b \rangle_2 \langle a_0, b \rangle_3 = \langle a_0, b \rangle_1 (\langle a_0, b \rangle_2 \langle a_0, b \rangle_3)$$

where  $\langle a_0, b \rangle_i$  denotes a finite product of  $a_0$ 's and  $b$ 's. If no  $a_0$  appears in the left side of (2.7), then the equation holds by the power-associativity of  $R$ . If exactly one  $a_0$  appears in the left side then it holds by either (2.5) or (2.6). Finally, if more than one  $a_0$  appears in the left side, then (2.7) holds since every associator in the ideal generated by  $I^2$  vanishes (also by (2.5) or (2.6) in some cases).

Now by the associativity of  $R_{a_0, b}$  and the commutativity of  $I$ , we easily compute that

$$(a_0b + b)^q = b^q + a_0b^q + \sum_{i=1}^{q-1} b^i a_0 b^{q-i} + \sum_{i=2}^q \binom{q}{i} a_i^i b^q,$$

and

$$(ba_0 + b)^q = b^q + b^q a_0 + \sum_{i=1}^{q-1} b^i a_0 b^{q-i} + \sum_{i=2}^q \binom{q}{i} a_i^i b^q.$$

By the commutativity of  $I$  and  $a_0b \neq ba_0$ ,  $(a_0b + b)a_0 \neq a_0(a_0b + b)$ . Then, since  $a_0b + b \equiv ba_0 + b \pmod{I}$ , we have by (ii) that

$$(2.8) \quad 0 = (a_0b + b)^q - (ba_0 + b)^q = a_0b^q - b^q a_0.$$

Now  $b^q \equiv b \pmod{I}$ , hence by the commutativity of  $I$

$$(2.9) \quad a_0b^q - b^q a_0 = a_0b - ba_0.$$

But (2.8) and (2.9) imply that  $a_0b = ba_0$ , a contradiction. Hence  $ab = ba$  for all  $a \in I, b \in R$ .

To complete the proof of the theorem, let  $x, y \in R$ . Since all elements of  $I$  commute with all elements of  $R$  we may assume that  $x, y \notin I$ . Since  $R/I$  is a finite field, the multiplicative group of nonzero elements of  $R/I$  is cyclic. Let  $\bar{\xi}$  be a generator of this group and  $\bar{\xi} = \xi + I, \xi \in R$ . Then, for some integers  $i, j$  and some  $a_1, a_2 \in I$ ,  $x = \xi^i + a_1$  and  $y = \xi^j + a_2$ . By an easy computation we get  $xy = yx$ . Hence  $R$  is commutative.

**THEOREM 3.** *Let  $R$  be a flexible power-associative ring and let  $I$  be an ideal in  $R$  such that  $R/I$  is a finite division ring of characteristic  $p > 5$  with exactly  $q$  elements. Suppose that (i)  $I$  is commutative and associative, and (ii)  $x \equiv y \pmod{I}$  implies  $x^q = y^q$  or both  $x$  and  $y$  commute with all elements of  $I$ . Then  $R$  is commutative.*

*Proof.* Assume that  $a_0b \neq ba_0$  for some  $a_0 \in I, b \in R$ . Let  $a \in I$  be arbitrary. We note that the proof of Theorem 2 is still valid through equation (2.4). Now since  $b$  does not commute with all elements of  $I$ , we have by (ii),  $x \equiv b \pmod{I}$  implies  $x^q = b^q = \eta$  where  $\eta$  is the common  $q^{\text{th}}$  power of all the elements  $x \equiv b \pmod{I}$ . We observe that since  $\eta^q \equiv \eta \equiv b^q \equiv b \pmod{I}$ ,  $\eta^q = b^q = \eta$ . Equations (2.3) and (2.4) become

$$(2.10) \quad (\eta^l, \eta^m, a) = (\eta^l, a, \eta^m) = (a, \eta^l, \eta^m) = 0, \quad l > 0, m > 0, \text{ integers.}$$

Now

$$(2.11) \quad (\eta + a, \eta + a, (\eta + a)^{q^l}) = 0, \quad l > 0 \text{ an integer.}$$

Since  $\eta + a \equiv \eta \pmod{I}$ ,  $(\eta + a)^q = \eta^q = \eta$ , hence by (2.10), (2.11)

becomes

$$(2.12) \quad (a, a, \gamma^l) = 0 .$$

Similarly

$$(2.13) \quad (a, \gamma^l, a) = (\gamma^l, a, a) = 0 .$$

Linearizing (2.12) and (2.13) we get

$$(2.14) \quad \begin{aligned} (\alpha_1, \alpha_2, \gamma^l) + (\alpha_2, \alpha_1, \gamma^l) &= (\alpha_1, \gamma^l, \alpha_2) + (\alpha_2, \gamma^l, \alpha_1) \\ &= (\gamma^l, \alpha_1, \alpha_2) + (\gamma^l, \alpha_2, \alpha_1) \\ &= 0 , \quad \alpha_1, \alpha_2 \in I . \end{aligned}$$

We now wish to show by induction that

$$(2.15) \quad (a^m, a, \gamma^l) = (a, a^m, \gamma^l) = 0, \quad m > 0, l > 0 \text{ integers.}$$

For  $m = 1$  (2.15) is true by (2.12). Assume (2.15) is true for  $m = k$ . Now it is readily verified that in an arbitrary nonassociative ring, the Teichmüller identity holds:

$$\begin{aligned} 0 &= h(w, x, y, z) \\ &= (wx, y, z) - (w, xy, z) + (w, x, yz) - w(x, y, z) - (w, x, y)z . \end{aligned}$$

Expanding  $0 = h(a, a^k, a, \gamma^l)$  we obtain by induction and the associativity of  $I$  that

$$0 = (a^{k+1}, a, \gamma^l) - (a, a^{k+1}, \gamma^l)$$

whence by (2.14),  $0 = 2(a^{k+1}, a, \gamma^l)$ . But then since  $p\gamma^l \in I$  and  $I$  is associative,

$$0 = (a^{k+1}, a, p\gamma^l) = p(a^{k+1}, a, \gamma^l) .$$

Therefore, since  $p > 2$ ,  $0 = (a^{k+1}, a, \gamma^l)$ . Hence (2.15) follows using (2.14).

Linearizing (1.1) yields

$$(2.16) \quad 0 = (x, y, z) + (z, y, x) .$$

Hence by (2.16), (2.15) yields

$$(2.17) \quad (\gamma^l, a, a^m) = (\gamma^l, a^m, a) = 0, \quad m > 0, l > 0 \text{ integers.}$$

The Jacobi identity

$$(xy, z) + (yz, x) + (zx, y) = (x, y, z) + (y, z, x) + (z, x, y)$$

can easily be shown to hold in an arbitrary nonassociative ring, where  $(u, v) = uv - vu$ . Now, in the Jacobi identity let  $x = y = a$ ,  $z = \gamma^l$

and use (2.12), (2.13), and the commutativity of  $I$  to obtain

$$(2.18) \quad (a^2, \eta^l) = 0 .$$

We now show by induction that

$$(2.19) \quad (a^m, \eta^l) = 0 , \quad m \geq 2, l > 0 \text{ integers.}$$

By (2.18), (2.19) is true for  $m = 2$ . Assume (2.19) is true for  $m = k$ . Now, by (2.15), the induction assumption, the commutativity of  $I$ , and (2.17) we obtain

$$a^{k+1}\eta^l = (aa^k)\eta^l = a(a^k\eta^l) = a(\eta^l a^k) = (\eta^l a^k)a = \eta^l(a^k a) = \eta^l a^{k+1} .$$

Hence (2.19) follows.

By (2.15), (2.10) and (2.14), we easily obtain

$$(2.20) \quad (a, a^m \eta^n, \eta^l) = (a^m \eta^n, a, \eta^l) = 0 , \quad m > 0, n > 0, l > 0 \text{ integers.}$$

Hence by (2.16)

$$(2.21) \quad (\eta^l, a^m \eta^n, a) = (\eta^l, a, a^m \eta^n) = 0 , \quad m > 0, n > 0, l > 0 \text{ integers.}$$

The anti-isomorphic copy  $R'$  or  $R$  satisfies the hypotheses of the theorem, hence (2.20) and (2.21) hold in  $R'$ . Therefore, in  $R$ , we have

$$(2.22) \quad \begin{aligned} 0 &= (\eta^l, \eta^n a^m, a) = (\eta^l, a, \eta^n a^m) \\ &= (a, \eta^n a^m, \eta^l) = (\eta^n a^m, a, \eta^l) , \quad m > 0, n > 0, l > 0 \text{ integers.} \end{aligned}$$

We wish to show by induction that

$$(2.23) \quad (a\eta)^m = a^m \eta^m , \quad m > 0 \text{ an integer.}$$

Indeed, using in order (2.20), the commutativity of  $I$ , (2.15), and (2.10), we compute

$$(a^k \eta^k)(a\eta) = ((a^k \eta^k)a)\eta = (a(a^k \eta^k))\eta = (a^{k+1} \eta^k)\eta = a^{k+1} \eta^{k+1} ,$$

hence the result. Considering the anti-isomorphic copy  $R'$  of  $R$  again yields, because of (2.23),

$$(2.24) \quad (\eta a)^m = \eta^m a^m , \quad m > 0 \text{ an integer.}$$

Next we show by induction that

$$(2.25) \quad (\eta + a)^m = \eta^m + \sum_{i=0}^{m-1} (\eta^i a) \eta^{m-i-1} + \sum_{i=2}^m \binom{m}{i} a^i \eta^{m-i} , \quad m \geq 2,$$

where by convention  $\eta^0 x = x \eta^0 = x$  for all  $x \in R$ . Clearly this is true for  $m = 2$ . Assume it is true for  $m = k$ . Then

$$\begin{aligned}
(\eta + a)^{k+1} &= \left( \eta^k + \sum_{i=0}^{k-1} (\eta^i a) \eta^{k-i-1} + \sum_{i=2}^k \binom{k}{i} a^i \eta^{k-i} \right) (\eta + a) \\
(2.26) \quad &= \eta^{k+1} + \sum_{i=0}^{k-1} ((\eta^i a) \eta^{k-i-1}) \eta + \sum_{i=2}^k \binom{k}{i} (a^i \eta^{k-i}) \eta \\
&\quad + \eta^k a + \sum_{i=0}^{k-1} ((\eta^i a) \eta^{k-i-1}) a + \sum_{i=2}^k \binom{k}{i} (a^i \eta^{k-i}) a .
\end{aligned}$$

By (2.10),

$$((\eta^i a) \eta^{k-i-1}) \eta = (\eta^i a) \eta^{k-i}, \quad i = 0, \dots, k-1,$$

and

$$(a^i \eta^{k-i}) \eta = a^i \eta^{k+1-i}, \quad i = 2, \dots, k.$$

Next, using in order the commutativity of  $I$ , (2.22), commutativity of  $I$  again, (2.13), (2.18), and (2.10), we compute for  $i = 0, \dots, k-1$

$$\begin{aligned}
((\eta^i a) \eta^{k-i-1}) a &= a((\eta^i a) \eta^{k-i-1}) = (a(\eta^i a)) \eta^{k-i-1} = ((\eta^i a) a) \eta^{k-i-1} \\
&= (\eta^i a^2) \eta^{k-i-1} = (a^2 \eta^i) \eta^{k-i-1} = a^2 \eta^{k-1}.
\end{aligned}$$

Finally, using the commutativity of  $I$  and (2.15) we compute for  $i = 2, \dots, k$

$$(a^i \eta^{k-1}) a = a(a^i \eta^{k-i}) = a^{i+1} \eta^{k-i}.$$

Therefore, (2.26) becomes

$$\begin{aligned}
(\eta + a)^{k+1} &= \eta^{k+1} + \sum_{i=0}^{k-1} (\eta^i a) \eta^{k-i} + \sum_{i=2}^k \binom{k}{i} a^i \eta^{k+1-i} \\
&\quad + \eta^k a + \sum_{i=0}^{k-1} a^2 \eta^{k-1} + \sum_{i=2}^k \binom{k}{i} a^{i+1} \eta^{k-i}.
\end{aligned}$$

Hence upon collecting terms we have

$$(\eta + a)^{k+1} = \eta^{k+1} + \sum_{i=0}^k (\eta^i a) \eta^{k-i} + \sum_{i=2}^{k+1} \binom{k+1}{i} a^i \eta^{k+1-i},$$

which completes the proof of (2.25).

Now, in (2.25) replace  $a$  by  $a\eta$  and by  $\eta a$  to obtain, respectively

$$(2.27) \quad (\eta + a\eta)^m = \eta^m + \sum_{i=0}^{m-1} (\eta^i (a\eta)) \eta^{m-i-1} + \sum_{i=2}^m \binom{m}{i} (a\eta)^i \eta^{m-i},$$

and

$$(2.28) \quad (\eta + \eta a)^m = \eta^m + \sum_{i=0}^{m-1} (\eta^i (\eta a)) \eta^{m-i-1} + \sum_{i=2}^m \binom{m}{i} (\eta a)^i \eta^{m-i}.$$

But by (2.19), (2.23), and (2.24),  $(a\eta)^i = (\eta a)^i$ ,  $i \geq 2$ . Moreover, by (2.10)

$$(\eta^i(a\eta))\eta^{m-i-1} = (\eta^i a)\eta^{m-i}$$

and

$$(\eta^i(\eta a))\eta^{m-i-1} = (\eta^{i+1}a)\eta^{m-i-1}, \quad i = 0, \dots, m - 1.$$

Therefore, upon subtracting (2.28) from (2.27) we obtain

$$(2.29) \quad (\eta + a\eta)^m - (\eta + \eta a)^m = a\eta^m - \eta^m a.$$

Now, since  $\eta + a\eta \equiv \eta + \eta a \equiv b \pmod{I}$  and since  $a_0 b \neq b a_0$ ,  $(\eta + a\eta)^a = (\eta + \eta a)^a$  which implies by (2.29) that  $a\eta^a = \eta^a a$ . However, since  $b^a = \eta^a$ ,  $b^a - b \in I$ , and  $I$  is commutative,

$$\begin{aligned} 0 &= a_0(b - b^a) - (b - b^a)a_0 = a_0 b - b a_0 - a_0 b^a + b^a a_0 \\ &= a_0 b - b a_0 - a_0 \eta^a + \eta^a a_0 = a_0 b - b a_0, \end{aligned}$$

a contradiction. Hence  $ab = ba$  for all  $a \in I$ ,  $b \in R$ .

The proof of Theorem 3 is completed exactly as that of Theorem 2.

3. Remarks. The following example is a model for our theorems which is not associative. First, we define  $R_1, I_1$  by

$$R_1 = \left\{ \begin{pmatrix} a & b \\ 0 & 0 \end{pmatrix} \mid a, b \in GF(p) \right\}; \quad I_1 = \left\{ \begin{pmatrix} 0 & b \\ 0 & 0 \end{pmatrix} \mid b \in GF(p) \right\};$$

$p > 5$  a prime.

Now, let “ $\circ$ ” denote the Jordan product in  $R_1$ :

$$\begin{pmatrix} a & b \\ 0 & 0 \end{pmatrix} \circ \begin{pmatrix} c & d \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} ac & \frac{ad + bc}{2} \\ 0 & 0 \end{pmatrix}.$$

Since  $p \neq 2$ ,  $ad + bc/2 \in GF(p)$ . Let  $R = R_1(+, \circ)$ , and let  $I = I_1(+, \circ)$ . It is readily verified that  $R$  is a Jordan ring which satisfies all the hypotheses of our theorems. Moreover,  $R$  is not associative. Other models for our theorems appear in [2, 3]. Also, examples are exhibited in those papers which show that the theorems fail in case  $R$  fails to satisfy all of the hypotheses of these theorems.

It can be shown that for the proof of Theorem 2, the condition that every associator contained in the ideal generated by  $I^2$  vanishes may be replaced by the more technical condition

$$(x, a_1, a_2) = (a_1, x, a_2) = (a_1, a_2, x) = 0 \quad \text{for all } x \in R, a_1, a_2 \in I.$$

Finally, an examination of [1, Th. 5] and the proofs of our theorems will reveal that they also hold when the characteristic of  $R/I$  is 3 or 5 providing the center of  $R/I$  has more than five elements.



## REFERENCES

1. A. A. Albert, *On nonassociative division algebras*, Trans. Amer. Math. Soc. **72** (1952), 296-309.
2. E. C. Johnsen, D. L. Outcalt, and Adil Yaqub, *A commutativity theorem for non-commutative Jordan rings*, Math. Jap. **11** (1967), 167-176.
3. D. L. Outcalt and Adil Yaqub, *A generalization of Wedderburn's Theorem*, Proc. Amer. Math. Soc. **18** (1967), 175-177.
4. R. D. Schafer, *Noncommutative Jordan algebras of characteristic 0*, Proc. Amer. Math. Soc. **6** (1955), 472-475.

Received October 3, 1967, and in revised form April 5, 1968. The first two authors were supported by Air Force Office of Scientific Research Grant AFOSR 698-67, and the third author by National Science Foundation Grant GP-5929.

UNIVERSITY OF CALIFORNIA, SANTA BARBARA

UNIVERSITY OF HAWAII

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. ROYDEN  
Stanford University  
Stanford, California

J. DUGUNDJI  
Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. R. PHELPS  
University of Washington  
Seattle, Washington 98105

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL WEAPONS CENTER

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California 90024.

Each author of each article receives 50 reprints free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners of publishers and have no responsibility for its content or policies.

# Pacific Journal of Mathematics

Vol. 27, No. 2

February, 1968

Leonard E. Baum and George Roger Sell, <i>Growth transformations for functions on manifolds</i> . . . . .	211
Henry Gilbert Bray, <i>A note on CLT groups</i> . . . . .	229
Paul Robert Chernoff, Richard Anthony Rasala and William Charles Waterhouse, <i>The Stone-Weierstrass theorem for valuable fields</i> . . . . .	233
Douglas Napier Clark, <i>On matrices associated with generalized interpolation problems</i> . . . . .	241
Richard Brian Darst and Euline Irwin Green, <i>On a Radon-Nikodym theorem for finitely additive set functions</i> . . . . .	255
Carl Louis DeVito, <i>A note on Eberlein's theorem</i> . . . . .	261
P. H. Doyle, III and John Gilbert Hocking, <i>Proving that wild cells exist</i> . . . . .	265
Leslie C. Glaser, <i>Uncountably many almost polyhedral wild <math>(k - 2)</math>-cells in <math>E^k</math> for <math>k \geq 4</math></i> . . . . .	267
Samuel Irving Goldberg, <i>Totally geodesic hypersurfaces of Kaehler manifolds</i> . . . . .	275
Donald Goldsmith, <i>On the multiplicative properties of arithmetic functions</i> . . . . .	283
Jack D. Gray, <i>Local analytic extensions of the resolvent</i> . . . . .	305
Eugene Carlyle Johnsen, David Lewis Outcalt and Adil Mohamed Yaqub, <i>Commutativity theorems for nonassociative rings with a finite division ring homomorphic image</i> . . . . .	325
André (Piotrowsky) De Korvin, <i>Normal expectations in von Neumann algebras</i> . . . . .	333
James Donald Kuelbs, <i>A linear transformation theorem for analytic Feynman integrals</i> . . . . .	339
W. Kuich, <i>Quasi-block-stochastic matrices</i> . . . . .	353
Richard G. Levin, <i>On commutative, nonpotent archimedean semigroups</i> . . . . .	365
James R. McLaughlin, <i>Functions represented by Rademacher series</i> . . . . .	373
Calvin R. Putnam, <i>Singular integrals and positive kernels</i> . . . . .	379
Harold G. Rutherford, II, <i>Characterizing primes in some noncommutative rings</i> . . . . .	387
Benjamin L. Schwartz, <i>On interchange graphs</i> . . . . .	393
Satish Shirali, <i>On the Jordan structure of complex Banach* algebras</i> . . . . .	397
Earl J. Taft, <i>A counter-example to a fixed point conjecture</i> . . . . .	405
J. Roger Teller, <i>On abelian pseudo lattice ordered groups</i> . . . . .	411